

**Zeitschrift:** Archives des sciences et compte rendu des séances de la Société  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 50 (1997)  
**Heft:** 3: Archives des Sciences

**Artikel:** Effect of UV-C on two cell lines from sugarbeet  
**Autor:** Notaris, Deniela / Crespi, Pierre / Greppin, Hubert  
**DOI:** <https://doi.org/10.5169/seals-740285>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Archs Sci. Genève	Vol. 50	Fasc. 3	pp. 223-232	Décembre 1997
-------------------	---------	---------	-------------	---------------

Communication présentée à la séance du 11 décembre 1997

## EFFECT OF UV-C ON TWO CELL LINES FROM SUGARBEET

BY

**Daniela NOTTARIS\*, Pierre CRESPI\*, Hubert GREPPIN\* & Claude PENEL\***

### ABSTRACT

**Effect of UV-C on two cell lines from sugarbeet.-** Two different sugarbeet calli were exposed to UV-C for short times. The first one, a hormone-dependent, non-organogenic callus (N), produced significant amounts of hydrogen peroxide even after short irradiations. The second one, a habituated, hormone-independent non-organogenic callus (H), appeared to be much less reactive and produced only weak amounts of H<sub>2</sub>O<sub>2</sub> after rather long exposures to UV-C. It was possible to isolate a substance as yet unidentified, which was less abundant in N than in H cells and which disappeared partly after UV-C treatments. Peroxidases from different cell compartments were measured 24 hours after UV-C irradiations, using guaiacol, ferulic acid or ascorbic acid as electron donors. In most cases, UV-C produced a decrease of activity. The most important effect was observed on ascorbate peroxidase activity which decreased strongly in every compartments of both cell types. This effect of UV-C could be explained by the presence of increased amount of H<sub>2</sub>O<sub>2</sub> in the apoplast.

**Key-words:** Tissue culture, Habituation, Hydrogen peroxide, Chemiluminescence, Peroxidases, Bilins.

### INTRODUCTION

Exposure of plants or cultured plant cells to ultraviolet radiation (UV) of various wavelengths brings about many physiological effects (CALDWELL, 1971; ROZEMA *et al.*, 1997). These effects include changes in ion permeability (MURPHY & WILSON, 1982; KARCZ & STOLAREK, 1988) or gene transcription (STRID, 1993), and inhibition of photosynthesis (BORNMAN, 1989; CALDWELL, 1971). The production of activated oxygen species is also a consequence of an irradiation with UV. It may be observed in animals (BERTLING *et al.*, 1996) and in plants (MURPHY & HUERTA, 1990) in which it induces an oxidative stress (PENEL, 1997). The accumulation of hydrogen peroxide has been reported to occur in whole plants, but also in suspension cultures of cells, following UV irradiation (MURPHY & HUERTA, 1990; RAO *et al.*, 1996). The question arises to know whether each type of cells respond to UV irradiation by such a production of H<sub>2</sub>O<sub>2</sub>, or if there are differences according to the physiological state or the redox status of the cells. To address this problem, we have submitted two different

---

\*Laboratoire de Physiologie végétale, Université de Genève, place de l'Université 3, CH-1211 Genève 4, Switzerland.

cell lines from sugarbeet to short irradiations with UV-C. One is a normal non-organogenic callus (N), the other one is a fully habituated, non-organogenic callus (H) growing without the supply of exogenous hormones. Both were established from leaf pieces of sugarbeet almost twenty years ago (DE GREEF & JACOBS, 1979) and kept their main properties since that time. Although they have been derived from the same plant and exhibit similar protein composition (TACCHINI *et al.*, 1995), these two calli are quite different in their ultrastructure (CREVECOEUR *et al.*, 1992), metabolisms and physiology (LE DILY *et al.*, 1993; GASPAR *et al.*, 1988). N callus is hormone-dependent for its growth and has a rather common behaviour. H callus, in addition to its complete independence with respect to exogenously supplied hormones, exhibits many of the characteristics of a cancerous state (GASPAR *et al.*, 1991). One of the most striking differences between these two calli concerns their redox properties. Whereas N cells produce much apoplastic  $H_2O_2$  and reduce only weakly extracellular ferricyanide, N cells synthesize little  $H_2O_2$  and exhibit a strong reducing capacity (CARRIÉ *et al.*, 1994). It appeared interesting to study the response of these two cell lines to an exposure to UV radiation, since their redox status is quite different. An explanation of the difference of response observed between N and H cells was tentatively looked for in their different tetrapyrrole metabolism (HAGEGE *et al.*, 1992).

## MATERIAL AND METHODS

All the experiments were performed on cells from sugarbeet (*Beta vulgaris* L., *altissima*) N and H calli. The experimental conditions to obtain and subculture these calli were already described in detail (KEVERS *et al.*, 1981). Forteen-day old calli were used for the experiments.

Ultraviolet irradiations (UV-C) were performed with a 15 W low pressure tube (50  $\mu W/cm^2$  at 254 nm). Petri dishes containing the calli were opened aseptically during UV exposure and then closed until cells were used. Stock solutions of biliverdin and bilirubin (Sigma) were prepared in methanol and diluted in water before addition on cells, which were then irradiated.

Apoplastic hydrogen peroxide was quantified by chemiluminescence in the presence of luminol, as described previously (CARRIÉ *et al.*, 1994), using 500 mg of cells dispersed in 1 ml Tris buffer at pH 8.6. Peroxidase activities were determined in three different fractions prepared as follows. One gram of callus was dispersed in 1 ml of 20 mM Hepes buffer pH 7. After a few min of incubation, the liquid which contained the soluble extracellular peroxidases was separated by centrifugation at 1'000 g for 5 min (fraction 1). Then the cells were ground in Hepes buffer pH 7, containing 1 M NaCl. After a centrifugation at 8'000 g for 10 min, the supernatant was used as fraction 2, containing mainly the ionically bound proteins. The pellet was then resuspended in a buffer containing 2% cellulase and 1% macerozyme (R10, Yakult Pharmaceutical, Japan). After an incubation overnight, the preparation was centrifuged at 8'000 g for 10

min. The supernatant contained the covalently bound proteins (fraction 3). Peroxidase activity was assayed in the three fractions using 8 mM guaiacol (2-methoxyphenol) and 2 mM  $\text{H}_2\text{O}_2$  in 50 mM phosphate buffer pH 6.1 ( $\Delta\text{A}470/\text{min.mg}$  protein) (KIEFER *et al.*, 1985), or 2 mM ferulic acid (4-hydroxy-3-methoxycinnamic acid) and 5 mM  $\text{H}_2\text{O}_2$  in acetate buffer pH 4.5 ( $\Delta\text{A}310/\text{min.mg}$  protein) (TAKAHAMA, 1995), or 0.4 mM ascorbic acid and 2 mM  $\text{H}_2\text{O}_2$  in phosphate buffer pH 6.1 ( $\Delta\text{A}265/\text{min.mg}$  protein) (CASTILLO *et al.*, 1986).

Samples of N and H calli were ground in methanol (1 gr/1 ml LiChrosolv, Merck). The extracts were centrifuged at 1'000 rpm for 5 min. Methanol was evaporated under nitrogen. The resulting dry residue was dissolved in 50  $\mu\text{l}$  methanol. After centrifugation at 1'000 rpm for 3 min, the sample was ready for HPLC separation. Five  $\mu\text{l}$  were injected into a HPLC column (Res Elut-BD5 C18, Varian). Methanol was used as eluent (flow rate 0.1 ml/min).

## RESULTS AND DISCUSSION

Samples of N and H calli were exposed to UV-C during various times to determine their sensitivity to this treatment (Fig. 1). When measured one hour after the irra-




FIG.1.

Effect of different durations of exposure to UV-C on the chemiluminescence level of N and H calli, measured 60 min after the treatment.

diations, the chemiluminescence of N callus was maximum after an exposure of 2-3 minutes and somewhat less important following longer treatments. H callus exhibited a small response only after 7-8 minutes. In the conditions used in the present work, the basal level of H callus chemiluminescence was near zero. Values for N cells in the absence of UV were comprised between 800 and 6000 cpm. These data are in accordance with those previously published (CARRIE *et al.*, 1994). The addition of catalase (10 units/ml) reduced the chemiluminescence by 65 per cent (data not shown), confirming that it is mainly due to the presence of  $H_2O_2$ .

The level of chemiluminescence produced by sugarbeet cells in the presence of luminol was measured after exposure to UV-C (Fig. 2). After a strong increase during the first hour following UV, N cells exhibited a fluctuating response. H callus showed a very slow and regular increase of chemiluminescence, which remained at a low level. In the days after a single UV-C treatment, N cells did not exhibit macroscopic signs of injury, whereas H callus presented many necroses. The data shown in Figs. 1 and 2 indicated that UV-C applied to plant cells for a short time induced the formation of  $H_2O_2$ , as already reported (MURPHY & HUERTA, 1990). The data indicated also that the extent of the response differed according to the type of cells which was irradiated. The resistance was also dependent on the cell type. Surprisingly, N cells, which accumulated much more  $H_2O_2$  than H cells, were less injured, suggesting that the level of this reduced oxygen species is not the only factor explaining the appearance of necrotic lesions.




FIG. 2.

Levels of chemiluminescence measured at different times after an exposure to UV-C of 2 min (N callus) or 8 min (H callus).

It is known that UV irradiation induces the formation of activated oxygen species in mammalian cells, and that bilirubin or biliverdin are produced to protect cells against oxidative stress (KEYSE & TYRRELL, 1989). Phycobiliproteins of cyanobacteria were also affected by an exposure to UV radiations (SINHA *et al.*, 1995). On the other hand, it has been reported that N and H calli differ by their tetrapyrrole metabolism (HAGEGE *et al.*, 1992). H cells exhibit a general deficiency in hemoproteins and chlorophylls. This particularity may be explained by the abnormal structure of their chloroplasts (CREVECOEUR *et al.*, 1992). Both types of cells use the Shemin pathway for their tetrapyrrole biosynthesis, which is rather unusual for plant cells (BISBIS *et al.*, 1997). The possibility exists also that this disturbed porphyrin metabolism results in the accumulation of bilin type compounds, which could play a role in the response of H cells to UV irradiations. In order to test this hypothesis, N cells were exposed to UV-C in the presence of bilirubin and biliverdin (Fig. 3), two products of heme degradation which are known to play a protective role in animal cells. These two molecules reduced




FIG. 3.

Effect of bilirubin (BR, 50 and 5  $\mu$ M), biliverdin (BV, 50 and 5  $\mu$ M) and methanol (met, 1%) on the level of chemiluminescence of N callus exposed to 2 min UV-C. Measurements were performed 60 min after irradiations.

significantly the extent of the chemiluminescence response, suggesting that they decreased the formation of  $H_2O_2$  upon exposure to UV-C, maybe by scavenging the newly formed oxygen species. Methanol, used to solubilize these two bilins had also an effect. These observations were in accordance with the known antioxidant properties of bilins (STOCKER, 1987).

Attempts to detect either biliverdin or bilirubin in sugarbeet calli, using commercial detection kits, or to find a heme oxygenase activity remained unsuccessful. The research of similar compounds was undertaken. Substances extracted with methanol from N and H cells were separated by HPLC, using a detection at 327 nm. In the

conditions used a major peak was detected during the elution with a retention time of 6.3-6.4 minutes. Biliverdin in the same conditions came out at about 6 minutes. The substance separated by this procedure was quantified in N and H calli at various times after exposure to UV-C (Fig. 4). H cells contained a significantly higher amount of this unknown substance. UV-C reduced rapidly its amount, down to a level similar to




FIG. 4.

Effect of an exposure of N and H calli to UV-C on the amount of the unknown substance extracted from one gram of callus at various times after the treatment. The substance separated by HPLC had a retention time of 6.3-6.4 min, and was detected at 327 nm. Peak area was expressed in arbitrary units.

the level of N cells within 90 minutes. The rather low amount of substance present in N cells was not much affected by UV-C. This substance has not been identified. It presented two absorption maxima at 276 and 327 nm (Fig. 5).

Apoplastic  $H_2O_2$  formed in response to a short UV-C irradiation is the substrate of peroxidases, enzymes which are known to be mainly present in cell wall and extracellular spaces (GASPAR *et al.*, 1982). An analysis of peroxidase activity was therefore performed in order to see if there was an effect of UV-C on these enzymes. As peroxidases are known to be able to use various electron donors to reduce  $H_2O_2$ , guaiacol, ferulic acid and ascorbic acid were tested with enzymes prepared from extracellular spaces and cells (Fig. 6). The general effect of UV-C on peroxidases was a decrease of activity in almost each case. This was particularly obvious when ascorbic acid was used as electron donor. Twenty four hours after an exposure to 2 min UV-C, the ascorbate peroxidase activity was decreased by 80 to 88 per cent in all fractions except H3. When determined with guaiacol or ferulic acid, the peroxidase activity exhibited also a decrease in UV-C irradiated N cells. The effect of the treatment was less important on these peroxidase activities in H cells. This difference between the two types of calli






FIG. 5.

Absorption spectra of the unknown substance separated by HPLC from 5 gr of N and H calli.

could be explained by the level of  $H_2O_2$  produced in reaction to UV-C. A one- or two-minute irradiation induced an important formation of  $H_2O_2$  in N callus apoplast, but not in H callus (Fig. 1). This would be consistent with the report showing that an increase of  $H_2O_2$  concentration in apoplast induced a degradation of proteins and an inactivation of many enzymes (GOMEZ *et al.*, 1995). The most striking differences were observed when ascorbic acid was used as electron donor. This particular peroxidase activity is normally due to ascorbate peroxidases (E.C. 1.11.1.11) which are distinct from the secreted plant peroxidases (E.C. 1.11.1.7), but the latter are also able to oxidise ascorbate in certain conditions (MELHORN *et al.*, 1996). The present results do not allow to know which kind of enzymes was measured, but it appeared clearly that this activity was the most sensitive to an exposure to UV-C, which could act directly on the protein or indirectly through the formation of  $H_2O_2$ . The main or unique function of ascorbate peroxidase is  $H_2O_2$  scavenging (PENEL, 1997). In addition, the results presented in Fig. 6 provided confirmation that peroxidase activity is greater in N than in H callus (KEVERS *et al.*, 1981).

In conclusion, the level of formation of  $H_2O_2$  in cells exposed to UV-C appeared to be different in different cells. This variability could result from the presence of an unknown substance which would have a protective effect. Further work will be necessary to identify the chemical nature of this product.






FIG. 6.

Peroxidase activities in fractions 1, 2 and 3 (see material and methods) measured with guaiacol, ferulic acid and ascorbic acid as electron donors. Control calli (0 min) and calli irradiated during 1 or 2 min were used 24 hours after UV-C.

## RÉSUMÉ

### EFFET DE L'UV-C SUR DEUX LIGNÉES CELLULAIRES DE BETTERAVE

Deux cals de betterave différents ont été exposés à un rayonnement UV-C pendant des temps courts. L'un deux, non-organogène croissant seulement en présence d'un apport d'hormones (N), produit des quantités importantes de peroxyde d'hydrogène en réponse à de courtes expositions aux UV-C. L'autre, cal habitué également non-organogène dont la croissance est indépendante d'hormones exogènes (H), ne produit que de faibles quantités de  $H_2O_2$  en réponse à une exposition relativement plus longue à UV-C. Une substance inconnue, dont la quantité diminue très nettement dans le cal H après exposition à UV-C, a été isolée dans les deux types de cals. L'activité peroxydasique extraite de différents compartiments sub-cellulaires a été mesurée 24 heures après une exposition à UV-C en utilisant le gaïacol, l'acide férulique et l'acide ascorbique comme donneurs d'électrons. Il ressort des mesures que l'UV-C a provoqué une diminution d'activité dans presque tous les cas. L'effet le plus important a été observé

sur l'activité ascorbate peroxydase, qui décroît dans les divers compartiments des deux types de cellules. Cet effet de l'UV-C pourrait être dû à l'augmentation de H<sub>2</sub>O<sub>2</sub> dans l'apoplaste.

**Mots-clés:** Culture de tissu, habitude, peroxyde d'hydrogène, chemiluminescence, peroxydases, bilines.

## REFERENCES

- ARNOTT, T. & T.M. MURPHY. 1991. A comparison of the effects of a fungal elicitor and ultraviolet radiation on ion transport and hydrogen peroxide synthesis by rose cells. *Environ. Experim. Bot.* 31: 209-216.
- BERTLING, C.J., F. LIN & A.W. GIROTTI. 1996. Role of hydrogen peroxide in the cytotoxic effects of UVA/B radiation on mammalian cells. *Photochem. Photobiol.* 64: 137-142.
- BISBIS, B., C. KEVERS, C. PENEL, H. GREPPIN & TH. GASPAR. 1997. Coexistence of the Beale and the Shemin pathways for the biosynthesis of tetrapyrrole-containing compounds, including peroxidases, in a normal sugarbeet callus. *Plant Perox. Newslett.* 9: 13-20.
- BORNMAN, J.F. 1989. Target sites of UV-B radiation in photosynthesis of higher plants. *J. Photochem. Photobiol. B*, 4: 145-158.
- CALDWELL, M.M. 1971. Solar UV irradiation and the growth and development of higher plants, pp. 131-178. In: *Photophysiology*, VI (A.C. Giese, ed.), Academic Press, New York.
- CARRIE, B., TH. GASPAR, H. GREPPIN & C. PENEL. 1994. Redox characteristics of normal and habituated cell lines of sugarbeet. *Plant Cell Environ.* 17: 457-461.
- CASTILLO, F.J., G. COTTON, C. KEVERS, H. GREPPIN, TH. GASPAR & T. VANDEN DRIESCHE. 1986. Changes in ascorbic acid content and ascorbate peroxidase activity during the development of *Acetabularia mediterranea*. *Differentiation* 33: 17-23.
- CREVECOEUR, M., D. HAGEGE, A.-M. CATESSON, H. GREPPIN & TH. GASPAR. 1992. Ultrastructure characteristics of cells from normal and habituated sugar beet calli. *Plant Physiol. Biochem.* 30: 87-95.
- DE GREEF, W. & M. JACOBS. 1979. In vitro culture of the sugarbeet: description of a cell line with high regeneration capacity. *Plant Sci. Lett.* 17: 55-61.
- GASPAR, TH., D. HAGEGE, C. KEVERS, C. PENEL, M. CREVECOEUR, I. ENGELMANN, H. GREPPIN & J.-M. FOIDART. 1991. When plant teratomas turn into cancers in the absence of pathogens. *Physiol. Plant.* 83: 696-701.
- GASPAR, TH., C. KEVERS, C. PENEL, M. CREVECOEUR & H. GREPPIN. 1988. Biochemical characterization of normal and habituated sugarbeet calli. Relationship with anatomy, habituation and organogenesis. *Potsdamer Forsch.* B57: 20-30.
- GASPAR, TH., C. PENEL, T. THORPE & H. GREPPIN 1982. *Peroxidases 1970-1980. A Survey of Their Biochemical and Physiological Roles In Higher Plants*. Université de Genève-Centre de Botanique.
- GOMEZ, L.D., L.M. CASANO & V.S. TRIPPI. 1995. Effect of hydrogen peroxide on degradation of cell wall associated proteins in growing bean hypocotyls. *Plant Cell Physiol.* 36: 1259-1264.
- HAGEGE, D., D. WERCK-REICHART, P. SCHMITT & TH. GASPAR. 1992. Deficiency in tetrapyrrole-containing compounds in a non-organogenic habituated sugarbeet cell line. *Plant Physiol. Biochem.* 30: 649-654.
- KARCZ, W. & J. STOLAREK. 1988. Effects of u.v.-C radiation on extension growth, H<sup>+</sup>-extrusion and transmembrane electric potential in maize coleoptile segments. *Physiol. Plant.* 74: 770-774.
- KEVERS, C., M. COUMANS, W. DE GREEF, M. HOFINGER & TH. GASPAR. 1981. Habituation in sugarbeet callus: auxin content, auxin protectors, peroxidase pattern and inhibitors. *Physiol. Plant.* 51: 281-286.

- KEYSE, S.M. & R.M. TYRRELL. 1989. Heme oxygenase is the major 32-kDa stress protein induced in human skin fibroblasts by UVA radiation, hydrogen peroxide, and sodium arsenite. *Proc. Natl. Acad. Sci. USA*, 86: 99-103.
- KIEFER, S., C. PENEL & H. GREPPIN. 1985.  $\text{Ca}^{2+}$  and  $\text{Mn}^{2+}$  mediated binding of the glycoprotein peroxidase to membranes of *Pharbitis* cotyledons. *Plant Sci.* 39: 37-43.
- LE DILY, F., J.-P. BILLARD, TH. GASPARD & C. HUAULT. 1993. Disturbed nitrogen metabolism associated with the hyperhydric status of fully habituated callus of sugarbeet. *Physiol. Plant.* 88: 129-134.
- MELHORN, H., M. LELANDAIS, H.G. KORTH & C.H. FOYER. 1996. Ascorbate is the natural substrate for plant peroxidases. *FEBS Lett.* 378: 203-206.
- MURPHY, T.M. & A.J. HUERTA. 1990. Hydrogen peroxide formation in cultured rose cells in response to u.v. radiation. *Physiol. Plant.* 78: 247-253.
- PENEL, C. 1997. Production and roles of hydrogen peroxide, pp. 219-238. In: *Travelling Shot on Plant Development* (H. Greppin, C. Penel and P. Simon, eds.) University of Geneva.
- RAO, M. V., G. PALIYATH & D.P. OMROD. 1996. Ultraviolet-B- and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiol.* 110: 125-136.
- ROZEMA, J., J. VAN DE STAAL, L.O. BJÖRN & M. CALDWELL. 1997. UV-B as an environmental factor in plant life: stress and regulation. *Trends Ecol. Evol.* 12: 22-28.
- STOCKER, R., Y. YAMAMOTO, A.F. McDONAGH, A.N. GLAZER & B.N. AMES. 1987. Bilirubin is an antioxidant of possible physiological importance. *Science* 235: 1043-1046.
- STRID, A. 1993. Alteration in expression of defence genes in *Pisum sativum* after exposure to supplementary ultraviolet-B radiation. *Plant Cell Physiol.* 34: 949-953.
- TACCHINI, P., A. FINK, G.X. XUE, TH. GASPARD & H. GREPPIN. 1995. Analysis of proteins of a fully habituated non-organogenic sugarbeet callus and a hormone-dependent one by high performance 2-D gel electrophoresis. *Plant Physiol. Biochem.* 33: 361-366.
- TAKAHAMA, U. 1995. Oxidation of hydroxycinnamic acid and hydroxycinnamyl alcohol derivatives by laccase and peroxidase. Interactions among *p*-hydroxyphenyl, guaiacyl and syringyl groups during the oxidation reactions. *Physiol. Plant.* 93: 61-68.