

The lichen genus *Parmentaria* from Indian subcontinent

Autor(en): **Upreti, D.K. / Singh, Ajay**

Objekttyp: **Article**

Zeitschrift: **Candollea : journal international de botanique systématique = international journal of systematic botany**

Band (Jahr): **43 (1988)**

Heft 1

PDF erstellt am: **31.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-879726>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The lichen genus *Parmentaria* from Indian subcontinent

D. K. UPRETI
&
AJAY SINGH

RÉSUMÉ

UPRETI, D. K. & A. SINGH (1988). Le genre de lichens *Parmentaria* dans le subcontinent indien. *Candollea* 43: 109-121. En anglais, résumés français et anglais.

Onze espèces du genre *Parmentaria* sont décrites en détail pour le subcontinent indien. *Parmentaria anamalaiense*, *P. ceylonense* et *P. indica* sont décrites comme de nouvelles espèces.

ABSTRACT

UPRETI, D. K. & A. SINGH (1988). The lichen genus *Parmentaria* from Indian subcontinent. *Candollea* 43: 109-121. In English, French and English abstracts.

Eleven species of the genus *Parmentaria* are described in detail from the Indian subcontinent. *Parmentaria anamalaiense*, *P. ceylonense* and *P. indica* are described as new species.

Introduction

The first report of *Parmentaria* from the Indian subcontinent in form of *Pyrenastrum americanum* Sprengel (= *Parmentaria astroidea* Fée) from Carnatic, South India came from BÉLANGER (1834). LEIGHTON (1869) in his enumeration of lichens from Sri Lanka (Ceylon) included the same taxon as *Verrucaria aspistea* var. *astroidea* Fée (= *Parmentaria astroidea* Fée) and STIRTON (1876) added one more species, *Astrothelium interlatens* (Nyl.) Stirton (= *Parmentaria interlatens* (Nyl.) Müll. Arg.) to the list of lichens of that country. Lastly, in the 19th century, STIRTON (1881) discovered *Verrucaria assamiensis* Stirton, in a collection made by A. Watt from Assam, which was re-named by ZAHLBRUCKNER (1922) as *Parmentaria* (?) *assamiensis* (Stirton) Zahlbr. It was, however, transferred to *Anthracotheceum* by SINGH (1982), proposing a new combination *Anthracotheceum assamiensis* (Stirton) A. Singh. Till 1983, thus, only two species of *Parmentaria* from this vast region were on record.

On the basis of investigations of old as well as recent collections of lichens from the Indian subcontinent, it has been realized that this region is after all not that poor in the representation of *Parmentaria* species as projected by the literature.

This work includes eleven species of *Parmentaria* from the Indian subcontinent, that are keyed and described in detail.

Taxonomic account

Parmentaria a stromatoid pyrenocarpous genus was placed in *Astrotheliaceae* by ZAHLBRUCKNER (1926). This family was characterized by radially arranged, \pm horizontally or obliquely placed perithecia opening laterally by their individual lateral ostiolar canals into a common central ostiole of the stroma or through individual separate ostioles converging to the centre of the stroma. This family could not sustain its identity in light of later works and the five rather

unrelated genera included in it were distributed to other families, to which they show affinity. Stroma, discarded as an important character at family level by Zahlbruckner (l.c.) seems to have lost its significance but remains the uniting factor of species at the generic level.

In recent classification, HALE (1967) placed *Parmentaria* along with *Anthracotheceum* in family *Arthopyrenaceae*, the only family belonging to order *Pleosporales*, which is characterised by well defined pseudothecia, resembling perithecia and asci \pm regularly arranged in stromatic layer. VĚZDA (1968), POELT (1973) and HESSEN & JAHNS (1974) placed *Parmentaria* in *Pyrenulaceae*, that also includes *Anthracotheceum*. *Parmentaria* and *Anthracotheceum* exhibit similar basic character of ascocarps, paraphysoid threads, asci and spores and show close affinity to each other. It is also evident from the fact that they have been placed in the same family by later workers.

Key to the *Parmentaria* species from the Indian subcontinent

1. Ostioles of different ascocarps in a stroma converging and uniting into one single ostiole of the stroma before communicating with the exterior 2
- 1a. Ostioles of different ascocarps in a stroma converging at the centre of stroma but remaining separate and each ascocarp communicating independently with the exterior 5
2. Asci 8-spored **3. *P. astroidea***
- 2a. Asci 2-spored 3
3. Stroma deeply embedded in thallus **8. *P. keralensis***
- 3a. Stroma situated in \pm hemispherical thalline verrucae 4
4. Spore cells parenchymatously arranged, i.e., primary septa indistinct.... **6. *P. indica***
- 4a. Spores with distinct 12-18 primary septa **7. *P. interlatens***
5. Spores 60 μ m or more long 6
- 5a. Spores 50 μ m or less long 7
6. Stroma deeply embedded in thallus, spores with 14-18 transverse tiers of cells
1. *P. anamalaiense*
- 6a. Stroma situated in convex thalline verrucae, spores with 8-10 transverse tiers of cells
10. *P. oligocarpa*
7. Ostiolar plate mamillate to conico-depressed 8
- 7a. Ostiolar plate flat 10
8. Ascocarps covered with thallus and of thalline colour or covered with thick corticiform layer and blackish **5. *P. immersa***
- 8a. Ascocarp naked and black 9
9. Ascocarps 1.0-1.2 mm in diam. **4. *P. ceylonense***
- 9a. Ascocarps up to 0.7 mm in diam. **9. *P. mamillata***
10. Stroma 2(-3)-carpous **2. *P. andamanica***
- 10a. Stroma mostly 4-7-carpous **11. *P. pluricarpa***

1. *Parmentaria anamalaiense* Upreti & A. Singh, spec. nov.

Type: India: Kerala, Anamalai hills, Thenmalai, 24.1.1976, Patwardhan & Nagarkar 76487B (AMH! Holotype) (growing along the holotype of *Anthracotheceum megaspermum*, leg. Patwardhan & Nagarkar 76487A) (Figs. 1, 12 & 23).

Thallus corticolous, endophloedus, stramineus, laevigatus. Stromata 1-3 carpa, omnia thallina immersa, distingubilis 1-3 brunnene punctatus ostiolis in minutuae papillae in pagina thallina. Ascocarpia oblique ampulliformis, perithecium fuliginosum, asci clavatae, 8-spori; sporae brunneolae, ellipsoidae vel nonnihil oblongo-ellipsoidae, murali divisae, cells 14-18 serie transversatis dispositis, 5-8-septatae, 60-90 μm longae, 30-35 μm crassae.

Thallus corticolous, endophloedal, straw-coloured, smooth, 60-80 μm thick, K-, C-, KC-, P-, hypothallus indistinct.

Stromata 1-3-carpous, ca. 1.5 mm in diameter, immersed, discernible by 1-3 brown punctate ostioles situated at the centre of minute papillae on the thalline surface. Ascocarps 900 μm long, 400 μm broad, obliquely flask-shaped with a slightly curved neck, each opening independently, wall black and carbonaceous, 80-100 μm thick at sides, colourless near ostiole and with heavy deposition of colourless crystals in this region; nucleus I + vinose red, without oil globules; paraphysoid threads simple; asci clavate, 8-spored, 290-320 \times 45-50 μm ; spores uni- or sometimes biserial in ascus, brown, ellipsoid to somewhat oblong-ellipsoid, multicelled-muriform, cells arranged in 14-18 transverse tiers, with up to 7 cells in each, with 5-8 primary septa, 60-90 \times 30-35 μm .

Remarks

P. anamalaiense resembles *P. oligocarpa* in thallus colour, independent ostiolar opening for each ascocarp in the stroma and spore size. The latter, however, has superficial and dull black ascocarps in contrast to those of the former that are deeply immersed in thallus and thus \pm vertical in orientation. Number of transverse tiers of cells in *P. oligocarpa* is 8-10 whereas in *P. anamalaiense* are 14-18 in number.

Known from the type collection only.

2. *Parmentaria andamanica* A. Singh, Feddes Repert. 96: 266. 1985.

Type: India: Andaman Islands, South Andaman Island, Port Blair, on twigs, *Singh 78802* (LWG! Holotype) (Figs. 2, 13 & 24).

Thallus corticolous, endophloedal, buff to straw-coloured, smooth, 80-100 μm thick, K-, C-, KC-, P-, hypothallus as a black-line, bordering the thallus.


Stromata up to 1.5 mm in diameter, 2(-3)-carpous, ascocarps radially arranged, 0.6-0.7 mm in diameter, ca. 700 μm long from bottom to top, immersed, later emerging and convex, becoming naked and black, ostioles converging at the centre of stroma but opening separately and situated on a plain to slightly raised, pale or brown to blackish, circular platform; perithecium black and carbonaceous, with or without moderate deposition of colourless crystals, 60-150 μm thick at top and sides, 200-250 μm thick at bottom (i.e. opposite ostiole); nucleus I + vinose red, without oil globules; paraphysoid threads simple; asci clavate to cylindrical, 8-spored, 174-200 \times 30-33 μm ; spores uniseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8-transverse tiers, with 1-3 cells in each, with 7 primary septa, 35-48 \times 15-20 μm .

Remarks

Parmentaria andamanica resembles *Parmentaria ravenelii*, but the latter differs in having larger number (up to 5) of ascocarps in a stroma, that are invariably covered with corticiform layer of thallus, and in larger spores, measuring up to 60 μm long.


Additional specimen examined

India: Andaman Islands, Long Island, *Singh 52954* (LWG); South Andaman Island, Port Blair, *Singh 68983* (LWG).




Figs. 1-7. — Habit (scale 1 division = 1 mm).

1, *Parmentaria anamalaiense*; 2, *P. andamanica*; 3, *P. astroidea*; 4, *P. ceylonense*; 5, *P. immersa*; 6, *P. indica*; 7, *P. interlatens*.




Figs. 8-11. — Habit (scale 1 division = 1 mm).
8, *P. keralensis*; 9, *P. mamillata*; 10, *P. oligocarpa*; 11, *P. pluricarpa*.



Figs. 12-22. — Vertical section of ascocarps.

12, *Parmentaria anamalaiense*; 13, *P. andamanica*; 14, *P. astroidea*; 15, *P. ceylonense*; 16, *P. immersa*; 17, *P. indica*; 18, *P. interlatens*; 19, *P. keralensis*; 20, *P. mamillata*; 21, *P. oligocarpa*; 22, *P. pluricarpa*.



Figs. 23-32. — Spores.

23 *Parmentaria anamalaiense*; 24, *P. andamanica*; 25, *P. astroidea*; 26, *P. ceylonense*; 27, *P. indica*; 28, *P. interlatens*; 29, *P. keralensis*; 30, *P. mamillata*; 31, *P. oligocarpa*; 32, *P. pluricarpa*.

3. *Parmentaria astroidea* Fée, Essai sur les Cryptogames des écorces officinales: xci et 70. 1824.

Type: not seen (Figs. 3, 14 & 25).

Thallus corticolous, endophloedal, buff, smooth, 97-150 μm , thick, K-, C-, KC-, P-, hypothallus indistinct.

Stromata up to 2 mm in diameter (2-)4-6-carpous; ascocarps radially arranged, 0.5-0.7 mm in diameter, 700-800 μm from bottom to ostiole, naked and black occasionally partly to \pm completely covered with thallus, obliquely placed and an obliquely running ostiolar canal leading to a common cavity (where other ostiolar canals of the stroma also open) communicating with the exterior by an apical pore; peritheciium black and carbonaceous, 50-100 μm thick all-round; nucleus I + vinose red, without oil globules; asci clavate, 8-spored, 195-210 \times 36-40 μm ; spores biseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8(-12) transverse tiers, 37-45 \times 15-20 μm .

Remarks

P. astroidea seems to be close to *P. baileyi* Müll. Arg. but the latter is reported to have the maximum of 4 ascocarps in a stroma as compared to up to ten (TUCKERMAN, 1958) in the former. *P. astroidea* has slightly larger spores than *P. baileyi*. Another species that shows superficial resemblance to *P. astroidea* is *P. subastroidea* Müll. Arg., but the latter is characterised by 2-spored asci.

Specimen examined

Sri Lanka (Ceylon); Central Province, *Thwaites CL 93C* (BM).

4. *Parmentaria ceylonense* Upreti & A. Singh, spec. nov.

Type: Sri Lanka (Ceylon), no precise locality, *Thwaites s.n. (Leighton 187)* (BM: holotype) annotated as *Verrucaria aspistea* Fée (non Ach.) var. *astroidea* and *Anthracotheccium libricolum* Müll. Arg. (Figs. 4, 15 & 26).

Thallus corticolous, endophloedus, bubalinus, ochraceus vel brunneolus, laevigatus. Stromata 1.5-1.8 mm diam., 2-6-carpa, convexa. Ascocarpia 1-1.2 mm diam., \pm pyriformis, convexa, denudata, nigra, ostiolis a unusquisque ascocarpum e stroma convergens sed remanens discretis; peritheciium fuliginosum, asci clavati, 8-spori; sporae fuscae, ellipsoidae, muraliter divisae, cellulis 8-seriatis transversaliter dispositis, 27-36 μm longae, 9-14 μm crassae.

Thallus corticolous, endophloedal, straw-coloured, ochre to brownish, smooth, 110-130 μm thick, K-, C-, KC-, P-, hypothallus indistinct.

Stromata 1.5-1.8 mm in diameter, 2-6-carpous, convex; ascocarps 1-1.2 mm in diameter, \pm pyriform, narrow towards ostiole and broad outward, up to 1100 μm from bottom to ostiole, \pm immersed when young, emerging later, convex, naked and black, each with its separate ostiole, ostioles converging and situated on central, slightly raised, concolorous to thallus or of darker shaded platform; peritheciium black and carbonaceous, with deposition of colourless crystals at non-carbonized ostiolar region, 35-50 μm thick at sides, 100-120 μm thick at bottom (opposite ostiole); nucleus I + vinose red, without oil globules; paraphysoid threads simple; asci clavate, 8-spored, 110-135 \times 27-30 μm ; spores uni- or biseriate in ascus, brown, ellipsoid, multi-celled muriform, with 8-transverse tiers of cells, with 2-3 cells in each, 27-36 \times 9-14 μm .

Remarks

P. ceylonense differs from *P. astroidea* Fée to which it was earlier annotated by means of its ostioles which like the latter species converge to the centre of the stroma but unlike it open separately to the exterior. In this respect *P. ceylonense* resembles *P. ravenelii* (Tuck.) Müll. Arg., *P. andamanica* and *P. mamillata*, but differs from them in its larger ascocarps which in *P. andamanica* and *P. mamillata* are up to 0.7 mm in diameter and in *P. ravenelii*, is up to 0.9 mm in diameter.

Known from the type collection only.

5. *Parmentaria immersa* (Patw. & Makhija) A. Singh, Nova Hedwigia 44(3-4): 373-375. 1987
= *Anthracothecium immersum* Patw. & Makhija, Kavaka 8: 22. 1980.

Type: India: Karnataka, Shimoga-dist., Sagar-Shimoga Road, Kalkoppa Forest, *Patwardhan & Kulkarni* 78-2872 (AMH! Holotype) annotated as *Anthracothecium immersum* by Patwardhan & Makhija (Figs. 5 & 16).

Thallus endophloedal, yellow-brown to brownish, smooth to verrucose, 60-150 μ m thick, verruca present in association with stroma, K-, C-, KC-, P-, hypothallus indistinct.

Thalline verruca always fertile, each enclosing a single stroma; stromata 1.5-1.75 mm in diameter, 3-4-carpous; ascocarps 0.5-0.75 mm in diameter, ca. 750 μ m long from bottom to ostiole, completely embedded in thalline verruca and concolorous to thallus or covered with thick corticiform layer and blackish, ostioles rather indistinct, converging but opening separately on a prominent protuberance of thalline colour; perithecium black and carbonaceous, 80-160 μ m thick at sides and 40-70 μ m thick at top; nucleus I —, without oil globules; paraphysoid threads simple; asci cylindrical, 8-spored, 205-210 \times 20-25 μ m, spores uniseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8(-4) transverse tiers, 1-3 cells in each, with 7 primary septa, 25-40 \times 10-18 μ m.

Remarks

P. immersa closely resembles *P. mamillata* in many characters including the ostiolar plate, which is raised as a protuberance or as a mamillate structure. The stromata in *P. immersa*, however, are completely immersed in the thalline wart and are concolorous to thallus or are sometimes covered with thick corticiform layer of thallus and are dull black, which in *P. mamillata* are naked and black.

Known from the type collection only.

6. *Parmentaria indica* Upreti & A. Singh, spec. nov.

Type: India: Arunachal Pradesh, Dibang Valley dist., Shantipur, alt. ca. 500 m, 6.6.1984, *Upreti* L81777 (LWG! Holotype) (Figs. 6, 17 & 27).

Thallus corticolous, endophloedus, laevigatus, flavide-fuscescentus. Stromata 1-3-carpa, 1.8-2 mm diam., convexa, orbiculata vel 2-3 labata, omnia strata thallina velata et thallina concolorata, interdum thallina non algifera obducta et nigra; ascocarpia 0.8-1 mm diam., pyriformia, ostiolum commune fuscum; perithecium fuliginosum, asci 2-spore; sporae fuscae, oblong-ellipsoidae, muraliter divisae, cellulia parenchymaticodispositis, 135-175 μ m longae, 40-50 μ m crassae.

Thallus corticolous, endophloedal, yellow-brown, smooth, 60-130 μ m, K-, C-, KC-, P-, hypothallus indistinct.

Stromata embedded in thalline warts, 1.8-2 mm in diam., 1-3-carpous, circular in outline or 2-3 lobate, ostioles one for each stroma, centrally located, laterally located in 1-carpous stroma, prominent, generally brown, ascocarps radially arranged, 0.8-1 mm in diam., \pm pyriform, with their narrow end towards the ostiole, completely embedded in and concolours with thallus, sometimes covered with corticiform layer only and then \pm dull black, each ascocarp opening laterally through a canal to the common ostiole; perithecium black and carbonaceous, 40-70 μ m thick at top and 60-120 μ m thick at sides, indistinct at bottom; nucleus I —, without oil globules; asci clavate, 2-spored, 170-220 \times 45-50 μ m; spores brown, oblong-ellipsoid, multi-celled muriform, cells parenchymatously arranged, 135-175 \times 40-50 μ m.

Remarks

P. indica is characterized by lobed, completely covered stromata with a single ostiole, 2-spored asci and spores measuring 135-175 \times 40-50 μ m, with cells parenchymatously arranged. In genus *Parmentaria*, six species possess \pm similar characters but they differ from *P. indica* in the following manners. The spores of *P. gregalis* Müll. Arg., *P. grossa* Müll. Arg. and *P. subalpina* Müll. Arg. are smaller and their length ranges between 75 and 120 μ m. *P. subastroidea* and *P. toowoombensis*

Müll. Arg. with 2-spored asci and spores with parenchymatously arranged cells, have 4-6-carpous stromata and the ascocarps are naked and black. In *P. lyoni* Zahlbr. the stroma, though 2-3-carpous, are larger (2.5 mm in diam.) and the ascocarps are naked, black and nitidus, *P. interlatens*, the only other species with 2-spored asci reported from the Indian subcontinent (Sri Lanka) possesses 1-3-carpous stromata like *P. indica* but the spores of *P. interlatens*, though of the comparable size with those of *P. indica* have the cells arranged in numerous transverse tiers of cells and are with distinct 12-18 primary septa.

Known from the type collection only.

7. *Parmentaria interlatens* (Nyl.) Müll. Arg., Flora 66: 244. 1883.

= *Astrothelium interlatens* Nyl., Bull. Soc. Linn. Normandie, Ser. 2: 95: 1968.

Type: not seen (Figs. 7, 18 & 28).

Thallus, corticolous, endophloedal, ochre, smooth, up to 136 µm thick, nitidous, K-, C-, KC-, P-, hypothallus indistinct.

Stromata 1.5-2.5 mm in diameter, 1-3-carpous, ± round and unlobed or slightly bi- to tri-lobed (depending on the number of ascocarps present), completely covered with and concolorous to thallus or at places blackish due to corticiform covering only, with a common ostiole at centre in the form of a brown to blackish depression; ascocarps 1-1.5 mm in diameter, 700-800 µm from bottom to ostiole, each communicating by a lateral ostiolar canal with the central cavity opening to the exterior by an apical pore; perithecium black and carbonaceous, 125-250 µm thick, nucleus I —, without oil globules; asci clavate, 2-spored, 230-260 × 55-65 µm; spores brown, ellipsoid to oblong-ellipsoid, multi-celled muriform, cells arranged in numerous transverse tiers, with 12-18 prominent transverse primary septa, 90-160 × 32-50 µm.

Remarks

An annotation on the specimen sheet reads "Sp. 2, interdum 4 nae, fusif. ellip. fusae vel fusco nigrae, murali 160-200 × 38-42 µ". The spore size as per observations is smaller. *P. interlatens* is similar to *P. gregalis* Müll. Arg. in the character of stroma, being concolorous to thallus, but spores in the latter are characterized by 21-24 primary septa. *P. interlatens* is close to *P. pyrioides* Müll. Arg. in outward resemblance, but the latter has 1-spored asci.

Specimen examined

Sri Lanka (Ceylon); Point de Galle, leg. illegible s.n. (BM).

8. *Parmentaria keralensis* (Patw. & Makhija) A. Singh, Nova Hedwigia 44(3-4): 374. 1987.

= *Anthracotheceum keralense* Patw. & Makhija, Kavaka 8: 23. 1980.

Type: India: Kerala, Udabanchola, Cardamom hills, in evergreen rain forest, *Nagarkar & Gole* 76827 (AMH! Holotype) annotated as *Anthracotheceum keralense* by Patwardhan & Makhija (Figs. 8, 19 & 29).

Thallus epiphloedal, ochre, smooth, 50-110 µm thick, undulate to irregularly verrucose (the last character to be attributed to the nature of the substratum and verrucae thus not associated with stroma), K-, C-, KC-, P-, hypothallus indistinct.

Stromata 1.5-2 mm in diameter, 2-4-carpous, completely embedded in thallus and not discernible externally, their position in the thallus being located by means of their ostioles, which are like minute brown to brownish-black circular, slightly depressed spots on the thalline surface; ascocarps 0.5-0.8 mm broad, 1200 µm from bottom to the lateral opening, deeply embedded in thallus; perithecium black and carbonaceous, 40-80 µm thick at top and 160-240 µm thick at sides; nucleus I —, with abundant oil globules; paraphysoid threads simple; asci clavate, 2-spored, 265-320 × 55-65 µm, spores brown, oblong-ellipsoid, multicelled muriform, cells ± parenchymatously arranged, 135-200 × 42-50 µm.

Remarks

P. keralensis shows affinity to *P. subastroidea* Müll. Arg. and *P. lyoni* in ascus and spore characters but the ascocarps in both the latter species are naked, black and nitidus.

Known from the type collection only.

9. *Parmentaria mamillata* A. Singh, Feddes Repert. 96: 266. 1985.

Type: Andaman Islands, South Andaman Island, Port Blair, on bark of *Mangifera indica*, Singh 78815 (LWG! Holotype) (Figs. 9, 20 & 30).

Thallus corticolous endophloedal, yellow-brown, smooth, 70-90 μm thick, \pm nitidus, K-, C-, KC-, P-, hypothallus indistinct.

Stromata up to 1.5 mm in diameter, 2-4-carpous, ascocarps radially arranged, up to 0.7 mm in diameter, ca. 700 μm long from bottom to ostiole, immersed, later emerging, convex, naked and black, each with separate ostiole, ostioles converging and situated on a central brown mamillate platform; perithecium black and carbonaceous, ca. 50 μm thick at top and sides, ca. 150 μm thick opposite the ostiole; nucleus I —, without oil globules; paraphysoid threads simple; asci clavate, 8-spored, 153-177 \times 28-40 μm ; spores uniseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8-10 transverse tiers, with 1-3 cells in each, with 7 primary septa, 20-40 \times 10-20 μm .

Remarks

P. mamillata is easily identified by the presence of the mamillate structure at the centre of the stroma where all the converging ostioles open.

Additional specimen examined

India: Andaman Islands, South Andaman Islands, Port Blair, on bark of *Mangifera indica*, Singh 68988b (LWG).

10. *Parmentaria oligocarpa* A. Singh, Candollea 38: 460. 1983.

Type: India: Tamil Nadu, Madurai district, Shola near 9th mile on Kodaikanal-Berijam Road, alt. 2290 m, Foreau & Awasthi 4192 (AWAS! Holotype) (Figs. 10, 21 & 31).

Thallus corticolous, endophloedal, buff to straw-coloured, smooth, 125-160 μm thick, K-, C-, KC-, P-, hypothallus indistinct.

Stromata (1-)2(-3)-carpous, up to 2 mm long, 1-1.5 mm broad, convex, \pm completely covered with and concolorous to thallus, only top part of individual ascocarps with a corticiform covering, blackish; ascocarps up to 1 mm in diameter, flask-shaped, each opening to the exterior independently through obliquely upward running ostiolar canal, all of which from a stroma, converging and each ending in an independent ostiole, ostioles situated on a thallus coloured or slightly darker-coloured, \pm circular, flat plate in form of (1-)2(-3) fine dots (depending on the number of ascocarps in the stroma); perithecium black and carbonaceous, with heavy deposition of colourless crystals, 200-225 μm thick at top and sides, 100-150 μm thick at bottom; nucleus I + vinose red, without oil globules; paraphysoid threads simple; asci clavate, 8-spored, 310-345 \times 55-70 μm ; spores uni- or biseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8-10 transverse tiers, 1-6 cells in each, with 7-9 primary septa, 70-92 \times 30-46 μm .

Remarks

P. oligocarpa resembles *P. chilensis* Swinscow in thallus colour as well as in stromata and ostiolar characters, the two species differ in other details nonetheless. In *P. chilensis* the number of ascocarps in a stroma is 1-5. The asci are 4-8-spored, a character by virtue of which, as stated by IMSHAUG & HARRIS (1969), *P. chilensis* is well isolated in the genus.

Known from the type collection only.

11. *Parmentaria pluricarpa* A. Singh, Candollea 38: 462. 1983.

Type: India: Kerala, Mallapuram district, Calicut University campus, 23rd Jan. 1975, *Singh & Ranjan 102233b* (LWG! Holotype) (Figs. 11, 22 & 32).

Thallus corticolous, endophloedal, buff, smooth, 60-70 μ m thick, K-, C-, KC-, P-, hypothallus in the form of black line, partly surrounding the thallus.

Stromata (2-)4-7-carpous, up to 2 mm in diameter, immersed to emergent; ascocarps stellately arranged in the stroma, ca. 1 mm in diameter, immersed to emergent and convex to hemispherical, covered by thallus or completely naked, dull to shining black, opening through obliquely upward running ostiolar canals, all converging to the centre of stroma and each ending in an independent ostiole, ostioles situated on a pale or black, \pm circular plate, in form of (2-)4-7 dots (depending on the number of ascocarps in the stroma); perithecium black and carbonaceous, with moderate deposition of colourless crystals, 40-130 μ m thick; nucleus I + vinose red, without oil globules; paraphysoid threads unbranched; asci cylindrical to clavate, 8-spored, 155-180 \times 26-32 μ m; spores uniseriate in ascus, brown, ellipsoid, multi-celled muriform, cells arranged in 8-transverse tiers, with 1-4 cells in each and with 7-primary septa, 28-38 \times 12-18 μ m.

Remarks

P. pluricarpa, somewhat resembles the new world species *P. ravenelii*. The two, however, differ in the following characters. The stroma in *P. pluricarpa* is a compact structure because of larger number of bigger ascocarps in it. That of *P. ravenelii* the stroma is (1-)2-3(-5)-carpous and presents rather a lax appearance. The ascocarps in the latter always remain covered with corticiform layer, which imparts a dull black coloration to them. The ostiolar plate in *P. pluricarpa* is brown in young stroma but turns black later on, but in *P. ravenelii* it always remains brown. The spores in the latter are larger, measuring 43-60 \times 15-22 μ m.

Known from the type collection only.

ACKNOWLEDGEMENTS

We are grateful to the Directors of M.A.C.S. Research Institute, Pune (AMH) and British Museum (Natural History), London for providing material on loan for this study; to Dr. D. D. Awasthi for making available to us the species of *P. oligocarpa* from his personal herbarium (AWAS); to the Director, National Botanical Research Institute, Lucknow for providing facilities for this work; and to Shri Murari Ranjan for helping in laboratory work.

REFERENCES

- BÉLANGER, M. C. (1834). Lichens. In: *Voyage aux Indes Orientales, ... Zoologie et botanique*: 113-144. (Botanique, partie II: Cryptogamie, par Ch. Bélanger & M. Bory de Saint-Vincent). A. Bertrand, Paris.
- HALE, M. E. Jr. (1967). *The biology of lichens* (Ed. 2). Ed. Arnold, London.
- HENSSEN, A. L. & H. M. JAHNS (1974). *Lichenes. Eine Einführung in die Flechtenkunde*, 467 pp. Thieme, Stuttgart.
- IMSHHAUG, H. A. & R. C. HARRIS (1969). *Parmentaria chilensis* Fée. *Lichenol.* 4: 77-82.
- LEIGHTON, W. A. (1869). The lichens of Ceylon, collected by G. H. K. Thwaites. *Trans. Linn. Soc. London* 25: 161-185.
- POELT, J. (1973). Classification. Appendix. In: AHMADJIAN, V. & M. E. HALE (Eds.), *The Lichens*: 599-632. Acad. Press, New York & London.
- SINGH, A. (1982). Observations on some pyrenocarpous lichens. 1. *Nova Hedwigia* 36: 237-239.
- SINGH, A. (1983). Two new species of the lichen genus *Parmentaria* from South India. *Candollea* 38: 459-463.
- SINGH, A. (1985). Two new species of the lichen genus *Parmentaria* from Andaman Islands, India. *Feddes Repert.* 96: 265-268.
- SINGH, A. (1987). Observations on some pyrenocarpous lichens. 3. *Nova Hedwigia* 44(3-4): 373-375.
- STIRTON, J. (1876). Descriptions of recently discovered lichens. *Proc. Roy. Philos. Soc. Glasgow* 10: 150-164.
- STIRTON, J. (1881). On the vegetable parasites on tea plant, more especially that of Assam. *Proc. Roy. Philos. Soc. Glasgow* 13: 181-193.
- TUCKERMAN, E. (1958). Supplement to an enumeration of North American lichens. Part first, containing brief diagnoses of new species. *Amer. J. Sci. & Arts, Ser. 2*, 25: 422-430.

- VĚZDA, A. (1968). Taxonomische Revision der Gattung *Thelopsis* Nyl. (Lichenisierte Fungi). *Folia Geobot. Phytotax. Pratheia* 3: 363-406.
- ZAHLBRUCKNER, A. (1922). *Catalogus lichenum universales*, Vol 1. Bornträger, Leipzig.
- ZAHLBRUCKNER, A. (1926). Lichens (Flechten). In: ENGLER, A. & K. PRANTL, *Die natürlichen Pflanzenfamilien* (Ed. 2, Band. 8). Engelmann, Leipzig.

