

Some new and interesting records for the flora of the United Arab Emirates

Autor(en): **Karim, Fawzi M.**

Objekttyp: **Article**

Zeitschrift: **Candollea : journal international de botanique systématique =
international journal of systematic botany**

Band (Jahr): **50 (1995)**

Heft 1

PDF erstellt am: **03.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-879455>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Some new and interesting records for the flora of the United Arab Emirates

FAWZI M. KARIM

ABSTRACT

KARIM, F. M. (1995). Some new and interesting records for the flora of the United Arab Emirates. *Candollea* 50: 25-31. In English, English and French abstracts.

Recent extensive floristic collections from U.A.E. especially the Al-Ain, Masafi and Dibba districts in the east and north east of the country, revealed the presence of 11 new taxa for the Flora of U.A.E. These are: *Peristrophe paniculata* (Acanthaceae), *Limeum humile* (Aizoaceae), *Amaranthus spinosus* (Amaranthaceae), *Halothamnus bottae* (Chenopodiaceae), *Lactuca serriola*, *Scorzonera tortuosissima* (Compositae), *Convolvulus glomeratus*, *Evolvulus alsinoides* (Convolvulaceae), *Fimbristylis ferruginea* (Cyperaceae), *Plantago cylindrica* (Plantaginaceae), *Calligonum cirnatum* (Polygonaceae).

RÉSUMÉ

KARIM, F. M. (1995). Quelques nouveautés pour la flore des Emirats Arabes Unis. *Candollea* 50: 25-31. En anglais, résumés français et anglais.

Des recherches récentes dans les districts de Al-Ain, Masafi et Dibba des Emirats Arabes Unis ont permis la découverte de 11 taxons nouveaux pour la flore des E.A.U. Il s'agit de: *Peristrophe paniculata* (Acanthaceae), *Limeum humile* (Aizoaceae), *Amaranthus spinosus* (Amaranthaceae), *Halothamnus bottae* (Chenopodiaceae), *Lactuca serriola*, *Scorzonera tortuosissima* (Compositae), *Convolvulus glomeratus*, *Evolvulus alsinoides* (Convolvulaceae), *Fimbristylis ferruginea* (Cyperaceae), *Plantago cylindrica* (Plantaginaceae), *Calligonum cirnatum* (Polygonaceae).

KEY-WOROS: Flora — United Arab Emirates — Arabian Gulf.

This study is based on collections made during many visits to the east and north-east regions of the United Arab Emirates, bordering the Sultanate of Oman (Fig. 1). This area does not seem to have been well explored botanically and the author was fortunate enough to make records in 1990-1992 of many taxa new for the region but well known however in the Mediterranean region and some other parts of the world.

Topographically the area comprises oasis plains, hills, wadis and mountains. The wadis of these regions mostly crossing sea mail Ophiolite of possible middle Cretaceous age, volcanics, metamorphic rocks and metasediments.

The climate of this region is characterised by low rainfalls and high temperatures; the mean annual rainfall being 70-150 mm in January, the mean maximum air temperature 43.6°C in June while the mean minimum air temperature is 8.5°C in January and the mean annual relative humidity is over 70%.

The assessment of the "new records" is based on "Flora of United Arab Emirates; an Introduction" (WESTERN, 1989), "The Living Desert" (JONGBLOED, 1987) and other Floras of the adjacent countries, e.g. Flora of Saudi Arabia (MIGAHID, 1978), Flora of Iraq (TOWNSEND & GUEST, 1974-1980) and Ecology & Flora of Qatar (BATANOUNY, 1981).




Fig. 1. — United Arab Emirates map showing three districts: 1, Al-Ain; 2, Masafi; 3, Dippa.

Specimens were deposited in the National Herbarium of United Arab Emirates University, The Desert and Marine Environment Research Center in Al-Ain (UAEUH) and duplicates were delivered to the Herbarium of the Royal Botanic Gardens in Edinburgh (E).

The following is an alphabetical list of the species indicated as "new" in the flora of the United Arab Emirates. The abbreviations used in the text are:

fl. = flowering specimen
fr. = fruiting specimen

Acanthaceae

***Peristrophe paniculata* (Forssk.) Brummitt (Fig. 2)**

Syn.: *Peristrophe bicalyculata* (Retz.) Nees
Dianthera bicalyculata Retz.

General distribution. — Mostly found in Tropical Africa, Egypt, India (BHANDARI, 1978), Oman (MILLER & MORRIS, 1988), also found in Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl.). Al-Ain district, Al-Oha, 350 m, in the Al-Oha farm, 14.12.1990, F. M. Karim 6717.




Fig. 2. — *Peristrophe paniculata* (Forssk.) Brummitt
a, flower; b, root.

*Aizoaceae***Limeum humile** Forsk.Syn.: *Limeum obovatum* Vicary

General distribution. — Tropical and S Africa, India, Pakistan and Saudi Arabia (NASIR, 1973), Egypt-Nubia (BHANDARI, 1978), Oman (GHAZANFAR, 1992), Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl. & fr.). Al-Ain district, Al-Oha, 350 m, on road side, 22.10.1990, F. M. Karim 4774.

*Amaranthaceae***Amaranthus spinosus** L.

General distribution. — It is of Tropical American origin, now a cosmopolitan weed in the warmer regions of the world and also occurring as a casual in some temperate regions; in Pakistan as elsewhere it occurs as a weed of cultivation, roadsides, waste places (TOWNSEND, 1974 and ZOHARY, 1973), also found in Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl.). Al-Ain district, near Jebal Hafit, 350 m, Saruje farm, 19.4.1992, F. M. Karim 6857.

*Chenopodiaceae***Halothamnus bottae** Jaub. & SpachSyn.: *Salsola bottae* (Jaub. & Spach) Boiss.

General distribution. — It is a West Irano-Turanian element often around rocky limestone hills with thin sand cover (ZOHARY, 1966), Kuwait (AL-RAWI, 1983), Oman (GHAZANFAR, 1992), Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fr.). Al-Ain district, near Hafit mountain, 380 m, on hill side, sandy gravelly soil, 31.12.1990, F. M. Karim 4989.

*Compositae***Lactuca serriola** L. (Fig. 3)Syn.: *Lactuca scariola* L.

General distribution. — A weed of abandoned farms, fields and waste ground around agricultural areas, chiefly in Europe and Mediterranean region (ZOHARY, 1973 and SHERING, 1989), Iraq (AL-RAWI, 1968), Jordan (AL-EISAWI, 1982) Lebanon, Palestine, Sinai (POST & DINSMORE, 1932-33), Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl. & fr.). 10 km. North of Masafi, 500 m, in farm, 20.5.1991, F. M. Karim 5310.

Scorzonera tortuosissima Boiss.Syn.: *Scorzonera intricata* Boiss. var. *petraea* Nabélek

General distribution. — Mediterranean and West Irano-Turanian element often around sandy rocky ground (FEINBRUN-DOTHAN, 1977) Iraq (AL-RAWI, 1968), Jordan (AL-EISAWI, 1982), Kuwait (AL-RAWI, 1983), Palestine (POST & DINSMORE, 1932-33), Saudi Arabia (COLLENETTE, 1985).




Fig. 3. — *Lactuca serriola* L.

Local distribution. — (fl. & fr.). Masafi to Dibba road, on hill side, among rocks, 25.2.1990, *F. M. Karim* 6707.

Convolvulaceae

Convolvulus glomeratus Choisy

Syn.: *C. congestus* R. Br.

C. arabicus Hochst. ex Hall. f.

Ipomoea auricoma Rich.

General distribution. — NE Africa (N Sudan, Egypt), Saudi Arabia, Iran, India, Pakistan and Afghanistan (AUSTIN & GHAZANFAR, 1979), Jordan (AL-EISAWI, 1982), Oman (MILLER & MORRIS, 1988), Qatar (BATANOUNY, 1981), Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl.). Al-Ain district, Kuwaitat farm, 350 m, salty place in the farm, 5.4.1990, *F. M. Karim* 9645.

Evolvulus alsinoides (L.) L.

Syn.: *Convolvulus alsinoides* L.

General distribution. — Nearly all warm countries, Egypt (TÄCKHOLM & DRAR, 1973), Southern USA through Central and South America, Africa, Iran, India, Pakistan, Ceylon and Malaysia (AUSTIN & GHAZANFAR, 1979), also found in Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl.). Al-Ain district, Al-Maqam farm, 350 m, in the farm, 3.2.1990, *F. M. Karim* 9644.

Cyperaceae

Fimbristylis ferruginea (L.) Vahl

Syn.: *Fimbristylis arvensis* Vahl

Scirpus ferrugineus L.

Scirpus arvensis Retz.

General distribution. — Mostly found in warm temperate marshy regions in the world, Australia, Polynesia, Malaysia and India (BHANDARI, 1978), Egypt, Sudan, Eritrea, Ethiopia, Sokotra, Tropical Africa, Palestine, Syria, Anatolia, Rhodes, Transjordan, Saudi Arabia, Iraq, Iran throughout India to Hongkong, China, Australia, Central and South America (TÄCKHOLM & DRAR, 1973), Bahrain (CORNES & CORNES, 1989), Lebanon (POST & DINSMORE, 1932-33) and Oman (MILLER & MORRIS, 1988).

Local distribution. — (fl.). Al-Ain district, Sueyhan, 120 m, on road side, salty place, 3.3.1991, *F. M. Karim* 4735.

Plantaginaceae

Plantago cylindrica Forssk.

General distribution. — It is a Saharo Arabian element found in Egypt, Palestine, Southern Iran to Pakistan (KAZMI, 1974 and ZOHARY, 1973), Jordan (AL-EISAWI, 1982), Iraq (AL-Rawi, 1968), Kuwait (DAOUD & AL-Rawi, 1985) and Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fl.). Al-Ain district, Al-Oha, 350 m, sandy gravelly soil, 12.2.1990, *F. M. Karim* 3505.

Polygonaceae

Calligonum cirnatum Boiss. subsp. **arabicum** (Sosk.) Sosk.

Syn.: *Calligonum arabicum* Sosk.

General distribution. — It is found in Asian desert areas, Beluchistan, Iran, India and Syria (BHANDARI, 1978), Oman (MILLER & MORRIS, 1988), Saudi Arabia (COLLENETTE, 1985).

Local distribution. — (fr.). Al-Faqa, 75 km North West of the Al-Ain, 120 m, sandy soil, on road side, 29.3.1991, F. M. Karim 4564.

ACKNOWLEDGEMENTS

The research for this paper was made possible through kind help and support by the Director of the Desert and Marine Environment Research Centre, U.A.E. University and by Dr. A. Khader, Head of Department, and by Dr. A. Miller of the Herbarium of the Royal Botanical Gardens, Edinburgh. Finally I am indebted to Prof. Dr. A. W. Lulua and Mr. J. A. Anter (Graphic Artist) for their kind help.

REFERENCES

- AL-EISAWI, D. M. (1982). List of Jordan vascular plants. *Mitt. Bot. Staatssamml. München* 18: 79-182.
- AL-RAWI, A. (1968). *Wild plants of Iraq*. Division of Botany and Range Management, Ministry of Agriculture, Baghdad.
- AL-RAWI, A. (1983). *Key of the flowering plants of Kuwait*. Kuwait Time Press, Kuwait.
- AUSTIN, D. F. & S. A. GHAZANFAR (1979). No. 126. Convolvulaceae. In: NASIR, E. (Ed.), *Flora of West Pakistan*. University of Pakistan and Kew Gardens, London.
- BATANOUNY, K. H. (1981). *Ecology and Flora of Qatar*. University of Qatar, Alden Press, Oxford.
- BHANDARI, M. M. (1978). *Flora of the Indian Desert*. Scientific Publishers, Jodhpur-India.
- COLLENETTE, S. (1985). *An Illustrated Guide to the Flowers of Saudi Arabia*. Scorpion Publishing Ltd., London.
- CORNES, C. D. & M. D. CORNES (1989). *The Wild Flowering plants of Bahrain*. IMMEL Publishing, London.
- DAOUD, H. S. & A. AL-RAWI (1985). *Flora of Kuwait*, vol. 1. Kuwait University Press.
- FEINBRUN-DOTHAN, N. (1977). *Flora Palaestina*, vol. 3. Jerusalem.
- GHAZANFAR, S. A. (1992). *An Annotated Catalogue of the Vascular Plants of Oman and their Vernacular Names*. Scripta Botanica Belgica 2. National Botanic Garden of Belgium. Meise, Belgium.
- JONGBLOED, M. (1987). *The Living Desert*. Arabian Heritage Series, Motivate Publishing, Dubai, U.A.E.
- KAZMI, S. M. A. (1974). No. 62. Plantaginaceae. In: NASIR, E. (Ed.), *Flora of West Pakistan*. University of Pakistan and Kew Gardens, London.
- MIGAHID, A. M. (1978). *Flora of Saudi Arabia*, ed. 2, vol. 1, 2. Riyadh University Publication.
- MILLER, A. G. & M. MORRIS (1988). *Plants of Dhofar*. Oman, Great Britain.
- NASIR, Y. (1973). No. 41. Aizoaceae. In: NASIR, E. (Ed.), *Flora of West Pakistan*. University of Pakistan and Kew Gardens, London.
- POST, G. E. & J. E. DINSMORE (1932-1933). *Flora of Syria, Palestine and Sinai*, ed. 2, 2 vols. Beirut.
- SCHERING, A. G. (1989). *Pflanzenschutz. Weeds in Sugar beet*. Agrochemical Division. A publication of Schering AG. Berlin, Germany.
- TÄCKHOLM, V. & M. DRAR (1973). *Flora of Egypt*, ed. 2, vol. 2. Otto Koeltz Publishing, Germany.
- TOWNSEND, C. C. (1974). No. 71. Amaranthaceae. In: NASIR, E. (Ed.), *Flora of West Pakistan*. University of Pakistan and Kew Gardens, London.
- TOWNSEND, C. C. & E. GUEST (1974-80). *Flora of Iraq*, vol. 3, 4 (part 1 & 2). Baghdad.
- WESTERN, A. R. (1989). *The Flora of the United Arab Emirates; An Introduction*. Al-Bayan, Dubai, U.A.E.
- ZOHARY, M. (1966). *Flora Palaestina*, vol. 1. Jerusalem.
- ZOHARY, M. (1973). *Geobotanical Foundation of the Middle East*, 2 vols. Amsterdam.

