

Chthonius thaleri, a new endogeic species from Venetia, Italy (Pseudoscorpiones: Chthoniidae) : on Italian pseudoscorpions XXXVI

Autor(en): **Gardini, Giulio**

Objektyp: **Article**

Zeitschrift: **Contributions to Natural History : Scientific Papers from the Natural History Museum Bern**

Band (Jahr): - **(2009)**

Heft 12/2

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-786982>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Chthonius thaleri, a new endogeic species from Venetia, Italy (Pseudoscorpiones: Chthoniidae)*

* On Italian pseudoscorpions XXXVI

Giulio Gardini

ABSTRACT

Contrib. Nat. Hist. 12: 545–550.

The endogeic pseudoscorpion *Chthonius (Ehippochthonius) thaleri* sp. nov. is described (♂ ♀; loc. typ.: Venetia, prov. of Padua, municipality of Arquà Petrarca: Monte Ventolone). The new species is related to the epigeic *C. parmensis* BEIER, 1963.

KURZFASSUNG

Chthonius thaleri, eine neue endogäische Art aus Venetien, Italien (Pseudoscorpiones, Chthoniidae) – Der endogäische Pseudoskorpion *Chthonius (Ehippochthonius) thaleri* sp. nov. wird beschrieben (♂ ♀; loc. typ.: Venetia, Provinz Padua, Gemeinde Arquà Petrarca: Monte Ventolone). Die neue Art ist verwandt mit der epigäischen *C. parmensis* BEIER, 1963.

Introduction

An intensive field study of the subterranean fauna of the karstic limestones of Venetia is being carried out since 1982 by Mr E. Piva and collaborators of the Speleo Club Proteo, Vicenza (Italy). A survey of the population of cavernicolous pseudoscorpions was first published by Gardini (1991), then Gardini (1996, 2000) summarized our knowledge of Venetian hypo- and epigeic pseudoscorpions. In this paper, a new endogeic species of *Chthonius* subgenus *Ehippochthonius* is described from Colli Euganei, Venetia. The species is named for the late Konrad Thaler in recognition of his contributions to the taxonomy, faunistics and ecology of alpine arachnids, chiefly Araneae.


Description

Chthonius (Ephippiochthonius) thaleri sp. nov.


Material studied: 1 ♂ and 4 ♀: Veneto, prov. di Padova, Colli Euganei, Arquà Petrarca, pendici N Monte Ventolone, 180 m, scavo in valle attiva, 8. IV. 1996, E. Piva leg. (1 ♂ holotype); id., 19. IV. 1996, E. Piva leg. (1 ♀ paratype); id., 200 m, 22. X. 1995, G. Peretto & E. Piva leg. (3 ♀ paratypes) (Coll. G. Gardini, c/o Dip. Te. Ris., Genoa; 1 ♀ paratype in coll. E. Piva, Vicenza).

Total length (in mm) 1.35 (♂) 1.45–1.55 (♀). Colour yellowish; cuticular surface of carapace, chelicerae and pedipalps slightly rasp-shaped; setae thick. Carapace 1.0 longer than broad, posteriorly slightly restricted; anterior border (Figs. 1, 2) medially a little prominent and dentate, laterally entirely notched; no eyes nor eye-spots; 16 macrosetae (4:4:4:2:2) and 6 microsetae: 2 (rarely one) "preocular" microsetae on each side and 2 microsetae belonging to the ocular series (Figs. 1, 2); length of central anterior macrosetae 0.10 mm. Tergal setae I–X: 4:4:4:4:6:6:6:6:6:6. Manducatory process with 2 setae; coxal setae: pedipalpal coxa 3, I 3 + 3 marginal microsetae, II 4 + 8–9 coxal spines, III 5 + 4–5 coxal spines, IV 6; intercoxal tubercle bisetose. Genital opening (♂) V-shaped, with 6 and 7 setae; genital operculum (♂ ♀) with 10 setae; setae of sternites III–X: 8 (♂) or 6–7 (♀) (+ 3 microsetae on each stigma): 7 (♂) or 6–8 (♀) (+ 2 m. on each stigma): 8:6:6:6:6:6. Chelicera (Figs. 3, 4) 2.2 (♂) 2.1–2.3 (♀) longer than broad, palm with 5 setae and 5 (rarely 6) microsetae laterally; teeth of fixed and movable fingers and spinneret as in Figs 3, 4; one isolated subdistal tooth on movable finger; ratio $gl = 0.54$ (♂) 0.55–0.57 (♀); flagellum with 11 setae.

Pedipalps (Figs. 5–7): femur 6.8 (♂) 6.1–6.5 (♀), patella 2.0 (♂) 1.8 (♀), chela 6.0 (♂) 5.5 (♀), hand of chela 2.2 (♂) 2.1 (♀) times longer than broad; movable chelal finger 1.7 (♂) 1.6 (♀) times longer than hand, femur 1.2 (♂ ♀) times longer than movable chelal finger and 1.5 (♂) 1.4 (♀) times longer than carapace. Hand of chela with a rounded hump distally of *ib/isb* and a marked hollow just before the base of the movable finger (Figs. 5–7). Fixed chelal finger with 16 triangular teeth (all with dental canal but the distal two) and 2–3 rounded ones (at level of *sb*); the third to last distal tooth is a little wider and with curved sides; latero-distal tooth acuminate. Movable chelal finger with 10 (♂) –9 (♀) triangular teeth (with dental canal) halfway between *sb/st* and 9–10 rounded vestigial teeth (without dental canal) with high base at level of *sb*; sensillum at level of *sb*; trichobothria see Figs. 6, 7.



Figs. 1–5. *Chthonius (E.) thaleri* sp. nov. – 1: Male holotype: antero-lateral portion of carapace; – 2: Female paratype: id.; – 3: Male holotype: chelicera; – 4: Female paratype: id.; – 5: Female paratype: pedipalpal hand, latero-ventral view.



Figs. 6–7. *Chthonius (E.) thaleri* sp. nov. – 6: Male holotype: pedipalpal chela, lateral view; – 7: Female paratype: id.

Measurements (in mm). Carapace: 0.38/0.365 (♂) 0.43–0.44/0.41–0.43 (♀). Chelicera: 0.325/0.145 (♂) 0.365–0.37/0.16–0.17 (♀), movable finger length 0.16 (♂) 0.185–0.19 (♀). Pedipalps: femur 0.575/0.085 (♂) 0.61–0.62/0.095–0.10 (♀), patella 0.215/0.105 (♂) 0.23–0.24/0.13 (♀), chela 0.78/0.13 (♂) 0.86/0.155 (♀), hand length 0.29 (♂) 0.33 (♀), movable finger length 0.485 (♂) 0.52 (♀).

Discussion

Chthonius (E.) thaleri sp. nov. is clearly related to the epigeic *C. (E.) parmensis* BEIER, 1963, the only species in Northern Italy with which it shares the peculiar dentition of the chelal fingers, the presence of a distinct dorsal hump distally of *ib/isb* and the presence of an isolated subdistal tooth on the movable cheliceral finger. *C. thaleri* sp. nov. differs from *C. parmensis* by the following characters (Beier, 1963; Callaini, 1984; De Vore-Scribante, 1999): posterior margin of carapace with 2 setae (4 setae in *C. parmensis*); no eyes nor eye-spots (anterior eyes with lense, posterior ones reduced in *C. parmensis*); length of pedipalpal chela 0.78–0.86 mm (0.48–0.59 mm in *C. parmensis*); pedipalpal chela ratio 5.5–6.0 times, finger/hand ratio 1.6–1.7 times (respectively 3.8–4.8 times and 1.3–1.6 times in *C. parmensis*); movable chelal finger with 9–10 pointed teeth (6–8 in *C. parmensis*).

Acknowledgements

I wish to express my thanks to Mr. Erminio Piva (Speleo Club Proteo, Vicenza) for sending once again the precious fruits of his researches.

References

- Beier, M. (1963): Pseudoscorpione aus dem Museum "Enrico Caffi" in Bergamo. — *Rendiconti dell'Istituto Lombardo di Scienze e Lettere* 97(B): 147–156.
- Callaini, G. (1984): Osservazioni su alcune specie di *Chthonius* del sottogenere *Ephippiochthonius* BEIER (Arachnida, Pseudoscorpionida, Chthoniidae). — *Annali del Museo civico di Storia naturale "G. Doria"* 85: 125–159.
- De Vore-Scribante, A. (1999): Les Pseudoscorpions de la Suisse. Étude systématique, faunistique et biogéographique. — 314 pp., Thèse Faculté des Sciences de l'Université de Genève, Département de Zoologie et Biologie animale.

Gardini, G. (1991): Pseudoscorpioni cavernicoli del Veneto (Arachnida). — Bollettino del Museo civico di Storia naturale di Verona 15(1988): 167–214.

Gardini, G. (1996): Pseudoscorpioni cavernicoli italiani. Seconda nota. — Bollettino del Museo civico di Storia naturale di Verona 20(1993): 157–174.

Gardini, G. (2000): Catalogo degli Pseudoscorpioni d'Italia (Arachnida). — Fragmenta entomologica, Roma 32 (suppl.): 1–181.

Address of the author:

Dr. Giulio Gardini
c/o Dip. Te. Ris., University of Genoa
Corso Europa 26
I-16132 Genoa, Italy

E-mail: giuliogardini@libero.it