

Pimoa thaleri, a new species of the genus Pimoa Chamberlin & Ivie, 1943 from India (Araneae: Pimoidae)

Autor(en): **Trotta, Alessio**

Objekttyp: **Article**

Zeitschrift: **Contributions to Natural History : Scientific Papers from the
Natural History Museum Bern**

Band (Jahr): - **(2009)**

Heft 12/3

PDF erstellt am: **30.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-787032>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Pimoa thaleri, a new species of the genus *Pimoa* CHAMBERLIN & IVIE, 1943 from India (Araneae: Pimoidae)

Alessio Trotta

ABSTRACT

Contrib. Nat. Hist. 12: 1403–1407.

A new species of Pimoidae, *Pimoa thaleri* sp. nov., is described from India (♂♀, loc. typ.: Bakim cave, Bakim, West Sikkim). Diagnostic and descriptive characters are given.

KURZFASSUNG

***Pimoa thaleri*, eine neue Art der Gattung *Pimoa* CHAMBERLIN & IVIE, 1943 von Indien (Araneae: Pimoidae) – *Pimoa thaleri* sp. nov., eine neue Art der Familie Pimoidae von Indien wird beschrieben (♂♀, loc. typ.: Bakim cave, Bakim, West Sikkim). Diagnos-tische und beschreibende Merkmale werden angegeben.**


Keywords: Araneae, Pimoidae, *Pimoa*, new species, cave spider, India, West Sikkim

Introduction

The family Pimoidae is a small relictual group (Hormiga & al., 2005) and contains 33 species separated into four genera (Platnick 2009): *Nanoa* HORMIGA, BUCKLE & SCHARFF, 2005, *Pimoa* CHAMBERLIN & IVIE, 1943, *Putaoa* HORMIGA & Tu, 2008 and *Weintrauboa* HORMIGA, 2003. The genus *Pimoa* includes currently 25 species (from Alaska, Canada, USA, Spain, France, Italy, China, India and Nepal) described by Simon (1884), Keyserling (1886), Fage (1931, 1946), Gertsch & Ivie (1936), Chamberlin & Ivie (1943), Gertsch (1951), Hormiga (1994), Griswold & al. (1999) and Xu & Li (2007). For the diagnosis of the family Pimoidae and the genus *Pimoa* see the detailed descriptions provided by Hormiga (1994). The discovery of a new species brings the total to 26 species.

The new species is named in honour of the great arachnologist Prof. Konrad Thaler, of whom I have been able to appreciate the kindness and seriousness.

Fig. 1. *Pimoa thaleri*
sp. nov. Male pedipalp.
Abbreviations see text.




***Pimoa thaleri* sp. nov. (Figs. 1–2)**

Material studied:

1 ♂ (holotype) and 1 ♀ (paratype) from Bakim cave, Bakim (West Sikkim), 2510 m, 12. IV. 1994, leg. L. De Pasquale, L. Latella & V. Sbordoni. Holotype, in alcohol, deposited in the Museum of Natural History of Verona, paratype in the author's collection.

Figs. 2. *Pimoa thaleri*
sp. nov., Epigynum.



Description and Diagnosis

Bristles are omitted from illustrations for clarity. General appearance as in its congeners. Cephalothorax (male: 2.3 mm long, 2 mm wide; female: 4.1 mm long, 3.4 mm wide) reddish brown, slightly darker at margins. Sternum reddish brown. Abdomen dark greyish with light dorsal pattern composed of four transverse undulating stripes (first and second continuous, third and fourth clearly interrupted). Cheliceral promargin and retromargin with three teeth; cheliceral stridulatory striae present in the male, absent in the female. Pedipalp orange and legs orange-brown with dark annuli.

The male (Fig. 1) can be recognized by its denticulate cymbial process (CDP), well developed with numerous (more than 20) denticles, and by the shape of the pimoid cymbial sclerite (PCS). The following characters are also diagnostic: median apophysis (MA) hook-shaped; pimoid embolic-tegular process (PEP) not bifurcated with apex twisted; embolus (E) and PEP roughly of the same length; paracymbium (P) short and rounded with long bristles.

The female can be recognized by the shape of the distal end of the epigynum (Fig. 2); epigynum is similar to those of *P. gandhii* HORMIGA, 1994 and *P. anatolica* HORMIGA, 1994, but differs from these species in having the apex of the distal end clearly shorter.

Distribution: Known only from the type locality.

Acknowledgements

My sincere thanks to Prof. V. Sbordoni (Rome) and Dr. L. Latella (Verona) for the loan of material and to Dr. G. Gardini (Genoa) for the suggestions and remarks.

References

- Chamberlin, R.V. & Ivie, W. (1943): New genera and species of North American linyphiid spiders. — Bulletin of the University of Utah 33 (10): 1–39.
- Fage, L. (1931): Araneae, 5e série, précédée d'un essai sur l'évolution souterraine et son déterminisme. — In: Biospeologica, LV. Archives de Zoologie expérimentale et générale 71: 91–291.
- Fage, L. (1946): Araignées cavernicoles de l'Inde. — Bulletin du Muséum national d'Histoire naturelle, Paris (2) 18: 382–388.
- Gertsch, W.J. (1951): New American linyphiid spiders. — American Museum Novitates 1514: 1–11.
- Gertsch, W.J. & Ivie, W. (1936): Descriptions of new American spiders. — American Museum Novitates 858: 1–25.
- Griswold, C.E., Long, C.L. & Hormiga, G. (1999): A new spider of the genus *Pimoa* from Gaoligong Mountains, Yunnan, China (Araneae, Araneoidea, Pimoidae). — Acta Botanica Yunnanica, Supplement 11: 91–97.
- Hormiga, G. (1994): A revision and cladistic analysis of the spider family Pimoidae (Araneoidea: Araneae). — Smithsonian Contributions to Zoology 549: 1–104.
- Hormiga, G. (2003): *Weintrauboa*, a new genus of pimoid spiders from Japan and adjacent islands, with comments on the monophyly and diagnosis of the family Pimoidae and the genus *Pimoa* (Araneoidea, Araneae). — Zoological Journal of the Linnean Society 139: 261–281.
- Hormiga, G., Buckle, D.J. & Scharff, N. (2005): *Nanoa*, an enigmatic new genus of pimoid spiders from western North America (Pimoidae, Araneae). — Zoological Journal of the Linnean Society 145: 249–262.
- Hormiga, G. & Tu, L. (2008): On *Putaoa*, a new genus of the spider family Pimoidae (Araneae) from China, with a cladistic test of its monophyly and phylogenetic placement. — Zootaxa 1792: 1–21.
- Keyserling, E. (1886): Die Spinnen Amerikas 2. Theridiidae. — Nürnberg, 1–222.
- Platnick, N.I. (2009): The world spider catalog, version 9.5. — American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html>
- Simon, E. (1884): Les arachnides de France. — Paris 5: 180–885.
- Xu, X. & Li, S.Q. (2007): Taxonomic study of the spider family Pimoidae (Arachnida: Araneae) from China. — Zoological Studies 46 (4): 483–502.

Address of the author:

Dr. Alessio Trotta
via delle Ginestre 1/A2
I-17024 Finale Ligure SV, Italy

E-mail: alessio_trotta@tin.it

