

Some properties of groups without the property P1.

Autor(en): **Losert, V.**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **54 (1979)**

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-41565>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Some properties of groups without the property P_1

V. LOSERT

Let G be a locally compact group with a left Haar measure dx . The group is said to have the property P_1 if for every $\epsilon > 0$ and every compact subset K of G there exists $s \in L^1(G)$ with $\|s\|_1 = 1$ and $\sup_{x \in K} \int_G |s(xy) - s(y)| dy < \epsilon$. (see [1] p. 168).

In [1] A. Derighetti introduced the quantity ρ_1 , which is defined as the infimum of all non-negative real numbers λ such that for every compact subset K of G there exists $s \geq 0$ with $\|s\|_1 = 1$ and $\sup_{x \in K} \int_G |s(xy) - s(y)| dy < \lambda$. He proved that $\rho_1 < 1$ implies property P_1 .

In this paper we are able to show that $\rho_1 < 2$ already implies property P_1 . It follows that ρ_1 can assume only two values, $\rho_1 = 0$ if G has property P_1 and $\rho_1 = 2$ if not. Analogous relations are proved for the constants ρ_p , which are defined in the same manner as ρ_1 , with $L^1(G)$ replaced by $L^p(G)$ ($1 \leq p < \infty$). The same constants are obtained if the system of compact subsets of G is replaced by that of finite subsets.

More generally one can consider the case of a locally compact space X on which G acts continuously and which admits a quasi invariant measure μ (see [5]). This defines a representation π of G on $L^2(X, \mu)$. One can define constants ρ_p as above and it turns out that again only two values for each ρ_p are possible, depending whether π weakly contains the one dimensional identity representation of G or not.

In the last section we show that the Fourier algebra $A(G)$ (or more generally $A_p(G)$) factorizes iff the group has property P_1 . This generalizes a result proved in [7] 2.3 for the free group of two generators.

Acknowledgement I want to thank H. Rindler with whom I had a number of valuable discussions on the subject of this paper.

Notations

Let G be a locally compact group with unit element e , X a locally compact topological space on which G acts continuously and which admits a quasi invariant Radon measure μ with modular function χ (cp. [5]).

If f is a function on X and $a \in G$, we define $af(x) = f(ax)$ for $x \in X$. \mathcal{F} (resp. \mathcal{K}) shall denote the family of all finite (resp. compact) subsets of G .

We have a representation $\pi_p(x)s = \gamma(x^{-1}, .)^{1/p}x^{-1}s$ for $x \in G$, $s \in L^p(X, \mu)$. (We will write π instead of π_2 .)

We define $\rho(\pi_p) = \sup_{K \in \mathcal{K}} \inf \{\sup_{x \in K} \|\pi(x)s - s\|_p : s \in L^p(X, \mu), s \geq 0, \|s\|_p = 1\}$ ($1 \leq p < \infty$)

$\rho^*(\pi_p)$ is defined in the same manner with \mathcal{K} replaced by \mathcal{F} .

$$d(\pi) = \sup_{K \in \mathcal{K}} \inf_{x \in K} \{(\pi(x)s, s) - 1 : s \in L^2(X, \mu)\}$$

$d^*(\pi)$ with \mathcal{F} instead of \mathcal{K} .

$\rho(\pi_2)$ coincides with the quantity $\rho(\pi)$ of [1]. For $X = G$, μ Haar measure $\rho(\pi_1)$ coincides with ρ_1 .

If M is a mean on $L^\infty(X, \mu)$ (a positive linear functional with $M(1) = 1$) we put $\alpha(M) = \sup \{|M(x\phi) - M(\phi)| : x \in G, \phi \in L^\infty(X, \mu), \|\phi\|_\infty \leq 1\}$.

If $x \in G$, ϵ_x shall denote the point measure of mass one concentrated in x . We write $L_R^1(G)$ for the space of *real*-valued integrable functions on G and $L_R^0(G) = \{f \in L_R^1(G) : \int f(x) dx = 0\}$.

LEMMA 1 Put $B = \{\epsilon_x * f - \epsilon_y * f : x, y \in G, f \in L_R^1(G), \|f\|_1 \leq 1\}$. Then the closed absolutely convex hull of B in $L_R^1(G)$ coincides with $\{f \in L_R^0(G) : \|f\|_1 \leq 2\}$.

Proof. We use the bipolar theorem [10] Th. 4 p. 35:

$$\begin{aligned} B^0 &= \{\phi \in L^\infty(G) : |\int g(x)\phi(x) dx| \leq 1 \quad \forall g \in B\} \\ &= \{\phi \in L_R^\infty(G) : \|_x\phi - {}_y\phi\|_\infty \leq 1 \quad \forall x, y \in G\}. \end{aligned}$$

If $\phi \in B^0$ is continuous this means that $|\phi(x) - \phi(y)| \leq 1$ for all $x, y \in G$. It follows that $\sup_{x \in G} \phi(x) - \inf_{x \in G} \phi(x) \leq 1$ (ϕ is real valued) and so there exists some $c \in \mathbf{R}$ such that $\|\phi - c\|_\infty \leq \frac{1}{2}$. B^0 is by definition closed with respect to the topology $\sigma(L^\infty, L^1)$, it is convex and left-translation invariant. It follows that $\phi \in B^0, f \in L^1(G), f \geq 0, \int f dx = 1$ implies $f * \phi \in B^0$.

If $\phi \in B^0$ is arbitrary, we choose an appropriate approximate identity (u_i) in $L^1(G)$ such that $u_i * \phi \in B^0$ and $u_i * \phi$ converges to ϕ in the topology $\sigma(L^\infty, L^1)$. Since $u_i * \phi$ is continuous we can find $c_i \in \mathbf{R}$ such that $\|u_i * \phi - c_i\| \leq \frac{1}{2}$. Let $c \in \mathbf{R}$ be a cluster point of the c_i , then some subset of $u_i * \phi - c_i$ converges to $\phi - c$ (for $\sigma(L^\infty, L^1)$) and it follows that $\|\phi - c\|_\infty \leq \sup_i \|u_i * \phi - c_i\|_\infty \leq \frac{1}{2}$. This means that $B^0 = \{\phi \in L_R^\infty(G) : \exists c \in \mathbf{R} : \|\phi - c\|_\infty \leq \frac{1}{2}\}$ (the converse inclusion is trivial).

If $f \in L_R^1(G), \int f dx = 0, \|f\|_1 \leq 2, \phi \in B^0, \|\phi - c\|_\infty < \frac{1}{2}$ then $|\int f(x)\phi(x) dx| = |\int f(x)\phi(x) - c dx| \leq 1$.

Consequently $f \in B^{00}$ and it is again trivial that any f in the closed absolutely convex hull of B satisfies $\int f dx = 0$ and $\|f\|_1 \leq 2$.

Remark. The conclusion of Lemma 1 is not valid in the case of complex valued functions:

Let $\alpha_1, \alpha_2, \alpha_3 \in \mathbf{C}$ be the roots of $z^3 = 1$ and assume that $\phi \in L^\infty(G)$ takes only the values $3^{-1/2} \alpha_i$ ($i = 1, 2, 3$) each on a set A_i of positive measure. Then ϕ belongs to B^0 but there does not exist a constant $c \in \mathbf{C}$ such that $\|\phi - c\|_\infty \leq \frac{1}{2}$. Now one can easily construct a function $f \in L^0(G)$ such that $\|f\|_1 = 2$ and $\arg f(x) = \bar{\alpha}_i$ for $x \in A_i$. Consequently $\int f(x)\phi(x) dx = 3^{-1/2} \int |f(x)| dx = 2 \cdot 3^{-1/2} > 1$ and so f does not belong to the closed, absolutely convex hull of B .

LEMMA 2. *If M is a mean on $L^\infty(X, \mu)$, $\phi \in L^\infty(X, \mu)$, $f \in L_R^0(G)$, we have $|M(f * \phi)| \leq 2^{-1} \alpha(M) \|f\|_1 \|\phi\|_\infty$.*

Proof. For fixed $\phi \in L^\infty(X, \mu)$ the map $f \rightarrow f * \phi$ is continuous from $L^1(G)$ to $L^\infty(X, \mu)$ (for the norm topology). For $x, y \in G$, $f \in L^1(G)$ we have by the definition of $\alpha(M)$:

$$|M(\epsilon_x * f * \phi - \epsilon_y * f * \phi)| \leq \alpha(M) \|f * \phi\|_\infty \leq \alpha(M) \|f\|_1 \|\phi\|_\infty.$$

The set $B_1 = \{g \in L_R^1(G) : |M(g * \phi)| \leq \alpha(M) \|\phi\|_\infty\}$ is a closed convex subset of $L_R^1(G)$ which contains the set $0.B = \{\epsilon_x * f - \epsilon_y * f : x, y \in G, f \in L_R^1(G), \|f\|_1 \leq 1\}$. By Lemma 1 B_1 contains all functions $g \in L_R^0(G)$ with $\|g\| \leq 2$ and so the result follows.

LEMMA 3. *If M is a mean on $L^\infty(X, \mu)$, C a compact subset of G , $\epsilon > 0$, then there exists $f \in L^1(X, \mu)$, $f \geq 0$ with $\int f(x) d\mu(x) = 1$ such that $\|\pi_1(x)f - f\| \leq \alpha(M) + \epsilon$ for all $x \in C$.*

Proof. Let h be an arbitrary function in $L^1(G)$ with $h \geq 0$, $\int h(x) dx = 1$. Then there exists some neighbourhood U of the unit element e in G such that $\|\epsilon_x * h - h\|_1 \leq \epsilon/2$ for $x \in U$. Let $F = \{x_1, \dots, x_n\}$ be a finite subset of G such that $C \subseteq FU$.

Now we consider the space Y which is defined as a product of $n+1$ -copies of $L_R^1(X)$ with the norm $\|(f_i)\| = \sup \|f_i\|_1$. Put $\alpha = \alpha(M)^{-1}$ (if $\alpha(M) = 0$ the same proof works if α is sufficiently large) and define a map $u : L_R^1(X, \mu) \rightarrow Y$ by

$$u(f) = (\alpha \pi_1(\epsilon_{x_1} * h - h)f, \dots, \alpha \pi_1(\epsilon_{x_n} * h - h)f, f).$$

(We write also π_1 for the extension of the representation to $L^1(G)$.) u is clearly linear and satisfies $\|f\|_1 \leq \|u(f)\| \leq 2\alpha \|f\|_1$.

The dual space Y' of Y can be identified with the sum of $(n+1)$ -copies of $L_R^\infty(X, \mu)$, equipped with the norm $\|(\phi_i)\| = \sum_{i=1}^n \|\phi_i\|_\infty$ and the dual map u'

of u is then given by $u((\phi_i)) = \sum_{i=1}^{n+1} \alpha[(\epsilon_{x_i} * h)^* * \phi_i - h^* * \phi_i] + \phi_{n+1}$. ($h * (x) = h(x^{-1}) \wedge_G (x^{-1})$). Let K be the preimage of the unit ball of Y by u , then the polar K^0 is the image of the unit ball of Y' by u' . If $\phi \in K^0$, it follows that there exist $(\phi_i)_{i=1}^{n+1} \in Y'$ with $\sum_{i=1}^{n+1} \|\phi_i\|_\infty \leq 1$ such that $\phi = \sum_{i=1}^n \alpha[(\epsilon_{x_i} * h)^* * \phi_i - h^* * \phi_i] + \phi_{n+1}$. By Lemma 2 $|M(\phi)| \leq 1$ and this means that M belongs to the bipolar K^{00} of K in $L_R^1(X, \mu)''$. By the bipolar theorem K is dense in K^{00} for the topology $\sigma(L_R^1(X, \mu)'', L_R^\infty(X, \mu))$. Since $M(1) = 1$ it follows that there exists $f_0 \in K$ such that $|\int_X f_0(x) d\mu(x) - 1| < \epsilon/8$. $f_0 \in K$ means that $\|f_0\| \leq 1$ and $\|\pi_1(\epsilon_{x_i} * h - h) f_0\|_1 \leq \alpha(M)$ for $i = 1, \dots, n$.

We decompose f_0 into its positive and negative part: $f_0 = f_0^+ - f_0^-$. From $\|f_0^+\|_1 + \|f_0^-\|_1 \leq 1$, $|1 - \|f_0^+\|_1 + \|f_0^-\|_1| < \epsilon/8$ it follows that $\|f_0 - f_0^+\|_1 = \|f_0^-\|_1 \leq \epsilon/16$ and $\|f_0^+\|_1 \geq 1 - \epsilon/8$. Put $f = f_0^+/\|f_0^+\|$. Then $\|\pi_1(\epsilon_{x_i} * h - h) f\|_1 \leq (\alpha(M) + \epsilon/8) (1 - \epsilon/8)^{-1} \leq \alpha(M) + \epsilon/2$. (We assume $\epsilon < 1$.)

Finally if $x \in U$ then $\|\pi_1(\epsilon_{x,x} * h - h) f\|_1 \leq \epsilon/2 + \|\pi_1(\epsilon_{x_i} * h - h) f\|_1 \leq \alpha(M) + \epsilon$. The following lemma, as well as its proof, is due to H. Rindler:

LEMMA 4. $\rho(\pi_1) = 2$ if and only if $\rho(\pi_2) = 2^{1/2}$ $\rho^*(\pi_1) = 2$ if and only if $\rho^*(\pi_2) = 2^{1/2}$.

Proof. Assume that $x \in G$, $f \in L^1(X, \mu)$, $f \geq 0$, $\|f\|_1 = 1$, $\|\pi_1(x)f - f\|_1 \geq 2 - \epsilon$. Put $g = f^{1/2}$.

Then

$$\begin{aligned} 2 - \epsilon &\leq \int_X |(\pi_2(x)g)^2 - g^2| d\mu \\ &= \int_X |\pi_2(x)g - g| |\pi_2(x)g + g| d\mu \\ &\leq \|\pi_2(x)g - g\|_2 \|\pi_2(x)g + g\|_2 \\ &= (2\langle g, g \rangle - 2\langle \pi_2(x)g, g \rangle)^{1/2} (2\langle g, g \rangle \\ &\quad + 8\langle \pi_2(x)g, g \rangle)^{1/2} \\ &= 2(1 - \langle \pi_2(x)g, g \rangle^2)^{1/2}. \end{aligned}$$

It follows that $\langle \pi_2(x)g, g \rangle \leq (1 - (1 - \epsilon/2)^2)^{1/2} \leq \epsilon^{1/2}$ and

$$\|\pi_2(x)g - g\|_2^2 = 2 - 2\langle \pi_2(x)g, g \rangle \geq 2 - 2\epsilon^{1/2}.$$

Conversely if $\|\pi_2(x)g - g\|_2^2 \geq 2 - \epsilon$ it follows from the equation $|a - b|^2 \leq |a^2 - b^2|$ ($a, b \geq 0$) that $\|\pi_1(x)f - f\|_1 \geq 2 - \epsilon$.

THEOREM 1. *The following statements are equivalent*

- (i) π does not contain weakly i_G
- (ii) $\rho(\pi_p) = 2^{1/p}$ for $1 \leq p < \infty$
- (iii) $\rho^*(\pi_p) = 2^{1/p}$ for $1 \leq p < \infty$
- (iv) $d(\pi) = 1$
- (v) $d^*(\pi) = 1$

Proof. If π contains i_G weakly then all the quantities in (ii)–(v) are zero and so (ii)–(v) implies (i).

If $\rho^*(\pi_2) < 2^{1/2}$ then by Lemma 4 $\rho^*(\pi_1) < 2$. The same argument as in [1] Prop. 1 shows that there exists a mean M on $L^\infty(X, \mu)$ with $\alpha(M) < 2$. By Lemma 3 $\rho(\pi_1) < 2$ and by Lemma 4 $\rho(\pi_2) < 2^{1/2}$. If $\|\pi_2(x)s - s\|_2^2 < \alpha < 2$ then $\langle \pi_2(x)s, s \rangle \geq 1 - \alpha/2$. It follows that $d(\pi) < 1$ and by [1] Cor. 14 π contains i_G weakly. This shows that (i) implies (ii), (iii) for $p = 1, 2$ and (iv).

If $d^*(\pi) = 1$ a simple computation shows that $\rho^*(\pi_2) = 2^{1/2}$. It remains to show that $\rho(\pi_2)^2 = 2$ implies $\rho(\pi_p) = 2^{1/p}$ for $1 \leq p < \infty$ (analogously for $\rho^*(\pi_p)$). By the inequality $|a - b|^p \leq a^p + b^p$ for $a, b \geq 0, p \geq 1$ it follows that $\rho(\pi_p) \leq 2^{1/p}$. Assume that $\|\pi_2(x)s - s\|_2^2 \geq 2 - 2\epsilon^2$ i.e. $\langle \pi_2(x)s, s \rangle \leq \epsilon^2$ and that $s \geq 0, \|s\|_2 = 1$. Put $t = \pi_2(x)s, A_1 = \{x \in X : s(x) \leq \epsilon t(x)\}, A_2 = \{x \in X : t(x) \leq \epsilon s(x)\}$ (we assume $\epsilon < 1$). Then it follows that $\epsilon \int_{X/A_2} s(x)^2 d\mu(x) \leq \int_{X/A_2} s(x)t(x) d\mu(x) \leq \epsilon^2$.

Consequently $\int_{X \setminus A_2} s(x)^2 d\mu(x) \leq \epsilon$ and similarly $\int_{X \setminus A_1} t(x)^2 d\mu(x) \leq \epsilon$. For $1 \leq p < \infty$ we get:

$$\begin{aligned} & \left(\int_{A_1} |s(x)^{2/p} - t(x)^{2/p}|^p d\mu(x) \right)^{1/p} \\ & \geq \left(\int_{A_1} t(x)^2 d\mu(x) \right)^{1/p} - \left(\int_{A_1} s(x)^2 d\mu(x) \right)^{1/p} \geq (1 - \epsilon)^{1/p} - \epsilon^{1/p}. \end{aligned}$$

The same estimate holds for $(\int_{A_2} |s(x)^{1/p} - t(x)^{1/p}|^p d\mu(x))/p$. This gives combined $\int_X |s(x)^{2/p} - t(x)^{2/p}| d\mu(x) \geq 2 - \delta(\epsilon)$ where $\delta(\epsilon) \rightarrow 0$ for $\epsilon \rightarrow 0$. If one puts $s_1 = s^{2/p}$ then $\|\pi_p(x)s_1 - s_1\|_p^p \geq 2 - \delta(\epsilon)$ and it follows that $\rho(\pi_p) \geq 2^{1/p}$.

For $f \in L^1(G)$, $x \in G$ put $A_x f = A_G(x)f_x$. If H is a closed normal subgroup we write T_H for the canonical map $L^1(G) \rightarrow L^1(G/H)$ and \mathcal{A} for the convex hull of $\{A_x : x \in H\}$. It is well known (see [8] p. 174) that if H has the property P_1 then

$$\inf_{A \in \mathcal{A}} \|Af\|_1 = \|T_H f\|_1 \text{ for all } f \in L^1(G).$$

The next proposition shows that a converse similar to Th. 1 holds for this characterization:

PROPOSITION 1. *If H does not have the property P_1 , then for any $\lambda < 1$ there exists $f \in L^1(G)$ such that $\|f\|_1 = 1$, $T_H f = 0$ and $\inf_{A \in \mathcal{A}} \|Af\| > \lambda$.*

Proof. If $\lambda < 1$ and for any $f \in L^1(G)$ with $T_H f = 0$ there exists $A \in \mathcal{A}$ such that $\|Af\| \leq \lambda \|f\|$, one can iterate this procedure to get $A \in \mathcal{A}$ such that $\|Af\| \leq \lambda^n \|f\|$. An analogous argument as in [8] Ch. 8, 4.5 p. 176 shows that for $f_1, \dots, f_m \in L^1(G)$ with $T_H f_i = 0$ and $\epsilon > 0$ there exists $A \in \mathcal{A}$ such that $\|Af_i\|_i < \epsilon$ for $i = 1, \dots, m$. Let $\{x_1, \dots, x_n\}$ be a finite subset of H , $\epsilon > 0$. If $f \in L^1(G)$ with $f \geq 0$, $\|f\|_1 = 1$ is arbitrary put $f_i = \epsilon_{x_i} * f - f$. Then $T_H f_i = 0$ and consequently there exists $A \in \mathcal{A}$ such that $\|Af_i\| < \epsilon$. Then $g = Af$ satisfies $g \geq 0$, $\|g\|_1 = 1$ and $\|\epsilon_{x_i} * g - g\|_1 < \epsilon$ for $i = 1, \dots, n$. This means that (G, H) has the property $P_*(G, H)$ (see [9] §3 p. 12) and by Prop. 1 p. 13 of [9], H has the property P_1 .

2. In the last section we consider some function algebras on a locally compact group, whose factorization properties depend on the property P_1 . The proof of this result was obtained independently by A. Derighetti.

Let $A_p(G)$ be the algebra of all functions $\sum f_n * g_n$, where $f_n \in L^p(G)$, $g_n \in L^q(G)$, $1/p + 1/q = 1$, $\check{g}_n(x) = g(x^{-1})$ (see [6]). – for $p = 2$ one gets the ordinary Fourier algebra of G . Let $B_p(G)$ be the algebra generated by those positive definite functions which are subordinate to the left regular representation of G on $L^2(G)$. [2] An algebra A is said to *factorize weakly*, if it coincides with the linear span of $A \cdot A$.

PROPOSITION 2. *If one of the algebras $A_p(G)$ ($1 \leq p < \infty$) or $B_p(G)$ factorizes weakly, then G has the property P_1 .*

Remark. If G has the property P_1 then $B_p(G)$ has a unit element and each $A_p(G)$ has a bounded approximate identity (see e.g. [6] p. 121).

Proof. If a function f belongs to one of these algebras, then its complex conjugate does also and has the same norm. By Th. 2.3 of [3] there exists a constant $K_1 > 0$ such that for each compact subset C of G there exists $f \in A$ such that $f \geq 1$ on C , $f \geq 0$ on G and $\|f\| \leq K_1$ (A denotes of the algebras cited above.) In the case of $A_p(G)$ one can now use the same argument as given in [6] Th. 6 to see that G has property P_1 . An analogous argument holds for $B_p(G)$ since this is the dual of the C^* -algebra on $L^2(G)$ generated by left convolution operators from $L^1(G)$ ([2] p. 192).

REFERENCES

- [1] A. DERIGHETTI, *On the Property P_1 of Locally Compact Groups*. Comment. Math. Helv. 46 (1971), 226–239.
- [2] P. EYMARD, *L’algèbre de Fourier d’un groupe localement compact*, Bull. Soc. Math. France 92 (1964), 181–236.

- [3] H. G. FEICHTINGER, C. G. GRAHAM, E. H. LAKIEN, *Partial Converses to the Cohen Factorization Theorem and Applications to Segral Algebras*, preprint.
- [4] F. P. GREENLEAF, *Ameanable Actions of Locally Compact Groups*, J. Funct. Analysis 4 (1969), 295–315.
- [5] S. L. GULICK, T. S. LIU, A. C. M. VAN ROOIJ, *Group Algebra Modules II*, Canad. J. Math. 19 (1967), 151–173.
- [6] C. HERZ, *Harmonic Synthesis for Subgroups*, Ann. Inst. Fourier 23 (1973), 91–123.
- [7] M. LEINERT, *Faltungsoperatoren auf gewissen diskreten Gruppen*, Studia Math. 52 (1975), 149–158.
- [8] H. REITER, *Classical Harmonic Analysis and Locally Compact Groups*, Oxford University Press 1968.
- [9] ——, *L^1 -Algebras and Segal Algebras*, Springer Lecture Notes 231.
- [10] A. P. ROBERTSON, W. J. ROBERTSON, *Topological Vector Spaces*, Cambridge Tracts in Math.

Viktor Losert
Mathematisches Institut
der Universität Wien
A-1900 Wien,
Strudlhofgasse 4
AUSTRIA

Received January 9, 1978