

Assembling homology classes in automorphism groups of free groups

Autor(en): **Conant, James / Hatcher, Allen / Kassabov, Martin**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **91 (2016)**

Heft 4

PDF erstellt am: **26.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-630576>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Assembling homology classes in automorphism groups of free groups

James Conant, Allen Hatcher, Martin Kassabov* and Karen Vogtmann**

Abstract. The observation that a graph of rank n can be assembled from graphs of smaller rank k with s leaves by pairing the leaves together leads to a process for assembling homology classes for $\text{Out}(F_n)$ and $\text{Aut}(F_n)$ from classes for groups $\Gamma_{k,s}$, where the $\Gamma_{k,s}$ generalize $\text{Out}(F_k) = \Gamma_{k,0}$ and $\text{Aut}(F_k) = \Gamma_{k,1}$. The symmetric group \mathfrak{S}_s acts on $H_*(\Gamma_{k,s})$ by permuting leaves, and for trivial rational coefficients we compute the \mathfrak{S}_s -module structure on $H_*(\Gamma_{k,s})$ completely for $k \leq 2$. Assembling these classes then produces all the known nontrivial rational homology classes for $\text{Aut}(F_n)$ and $\text{Out}(F_n)$ with the possible exception of classes for $n = 7$ recently discovered by L. Bartholdi. It also produces an enormous number of candidates for other nontrivial classes, some old and some new, but we limit the number of these which can be nontrivial using the representation theory of symmetric groups. We gain new insight into some of the most promising candidates by finding small subgroups of $\text{Aut}(F_n)$ and $\text{Out}(F_n)$ which support them and by finding geometric representations for the candidate classes as maps of closed manifolds into the moduli space of graphs. Finally, our results have implications for the homology of the Lie algebra of symplectic derivations.

Mathematics Subject Classification (2010). 20J06, 20F65, 20F28.

Keywords. Automorphisms of free groups, moduli spaces of graphs, cohomology.

1. Introduction

In this paper we develop a new approach to studying the homology of automorphism groups of free groups which gives fresh group theoretic and geometric insight into known families of homology classes, and also helps direct the search for new classes. We restrict attention to homology and cohomology with untwisted coefficients in a field \mathbb{k} of characteristic zero unless explicitly specified otherwise.

Let us recall briefly what is known about these homology groups. First of all, $H_i(\text{Aut}(F_n))$ and $H_i(\text{Out}(F_n))$ are finite-dimensional over \mathbb{k} for all i , and vanish

*Martin Kassabov was partially supported by Simons Foundation grant 305181 and NSF grants DMS 0900932 and 1303117.

**Karen Vogtmann was partially supported by NSF grant DMS 1011857 and a Royal Society Wolfson Award.

for i greater than the virtual cohomological dimension (vcd), which is $2n - 2$ for $\text{Aut}(F_n)$ and $2n - 3$ for $\text{Out}(F_n)$ [14]. The groups $H_i(\text{Aut}(F_n))$ and $H_i(\text{Out}(F_n))$ are independent of n for $n \geq 5(i + 1)/4$ as shown in [23, 24], and these stable groups are in fact zero as well, as Galatius proved in [16]. Thus in the first quadrant of the (i, n) plane (see Figure 1 below) there is a wedge-shaped region bounded by lines of slope $1/2$ and $5/4$ that contains all the nonzero groups $H_i(\text{Aut}(F_n))$, and similarly for $H_i(\text{Out}(F_n))$. There are only a small number of these groups which are explicitly known to be nonzero. For $\text{Aut}(F_n)$ these occur for $(i, n) = (4, 4), (7, 5), (8, 6), (8, 7), (11, 7)$ and $(12, 8)$; for $\text{Out}(F_n)$ the list is the same except that $(7, 5)$ is omitted. (The natural map $H_i(\text{Aut}(F_n)) \rightarrow H_i(\text{Out}(F_n))$ is known to be surjective for all i and n [27] and we give a different proof of this in Theorem 2.4.) These low-dimensional calculations are done mostly by computer; see [1, 12, 17, 19, 23, 33]. Complete calculations of $H_i(\text{Aut}(F_n))$ have been made only for $n \leq 5$ and for $H_i(\text{Out}(F_n))$ only for $n \leq 7$.




Figure 1. Classes in the homology of $\text{Aut}(F_n)$ for $n \leq 14$. The Morita classes are shown as squares and the Eisenstein classes are shown as circles, filled in if the classes are known to be nontrivial. The nontrivial classes recently found by Bartholdi are shown as hexagons.

There are two potentially infinite sequences which begin with nontrivial classes: these are classes μ_k for $(i, n) = (4k, 2k + 2)$ defined by Morita [33] and classes \mathcal{E}_k for $(i, n) = (4k + 3, 2k + 3)$ constructed in [10]. The latter are known as *Eisenstein classes* because they arise from Eisenstein series via the connection between modular forms and the cohomology of $\text{SL}_2(\mathbb{Z})$ established by the Eichler–

Shimura isomorphism. The Morita classes μ_k are defined for both Aut and Out , while the \mathcal{E}_k 's are defined for Aut and map to zero in Out . Note that these classes are all either one or two dimensions below the vcd .

One of the big open questions is to determine which of the classes μ_k and \mathcal{E}_k are nonzero. However, even if they are nonzero it seems that they account for only a small fraction of the homology. The Euler characteristic for $H_*(\text{Out}(F_n))$ was computed for $n \leq 11$ by Morita, Sakasai, and Suzuki in [35, 36], and after starting with the values 1 and 2 for $n \leq 8$, it becomes $-21, -124, -1202$ for $n = 9, 10, 11$. If this trend continues for larger n , it would say there are many odd-dimensional classes for $\text{Out}(F_n)$, though the only one discovered to date is the 11-dimensional class in $\text{Out}(F_7)$ recently found by Bartholdi [1]. (This class is balanced by a single 8-dimensional class, consistent with the Euler characteristic calculation for $n = 7$.)

| n | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|-------------------------|---|---|---|---|---|---|-----|------|-------|----|
| $\chi(\text{Out}(F_n))$ | 1 | 2 | 1 | 2 | 1 | 1 | -21 | -124 | -1202 | ? |

Figure 2. Euler characteristic of $\text{Out}(F_n)$

The Morita classes μ_k were first constructed using Lie algebra techniques underpinned by Kontsevich's "formal noncommutative symplectic geometry" [28, 29]. In [12] these classes were interpreted explicitly in Lie graph cohomology and generalized; further generalizations including the classes \mathcal{E}_k were obtained using "hairy graph homology" in [10]. In the present paper we show how to construct all of these classes in an elementary fashion which bypasses both graph homology and Kontsevich's work. The idea is to use the fact that $\text{Out}(F_n)$ and $\text{Aut}(F_n)$ are the first two groups in a series $\Gamma_{n,0}, \Gamma_{n,1}, \Gamma_{n,2}, \dots$ where $\Gamma_{n,s}$ is the group of homotopy classes of self-homotopy equivalences of a rank n graph fixing s leaves (vertices of valence one) [2, 22, 24]. These groups are related by natural surjective homomorphisms $\Gamma_{n,s} \rightarrow \Gamma_{n,s-k}$ with kernel $(F_n)^k$. These homomorphisms split for $k < s$ but not for $k = s$.

The groups $\Gamma_{n,s}$ are of interest because by gluing graphs together along a subset of their leaf vertices we obtain many homomorphisms $\Gamma_{n_1,s_1} \times \dots \times \Gamma_{n_k,s_k} \rightarrow \Gamma_{n,s}$. On the level of homology, each such map induces a homomorphism

$$H_*(\Gamma_{n_1,s_1}) \otimes \dots \otimes H_*(\Gamma_{n_k,s_k}) \longrightarrow H_*(\Gamma_{n,s}),$$

which we call an *assembly map* (see Section 5). For example by pairing up all of the leaves of two rank one graphs with s leaves (in any way) we obtain an assembly map

$$H_*(\Gamma_{1,s}) \otimes H_*(\Gamma_{1,s}) \longrightarrow H_*(\Gamma_{s+1,0}) = H_*(\text{Out}(F_{s+1})).$$

Restricting to the case that s is odd, say $s = 2k + 1$, it is easy to calculate that $H_{2k}(\Gamma_{1,2k+1}) \cong \mathbb{k}$ (see Section 3.4), and in Section 5.1 we note that the Morita

class μ_k is the image of $\alpha_k \otimes \alpha_k$ under this assembly map, where α_k is a generator of $H_{2k}(\Gamma_{1,2k+1})$. This graphical interpretation of the original Morita series allows us to give two new proofs that Morita classes vanish after one stabilization, one proof being algebraic (Section 6.2) and the other geometric (Section 7). This result was first proved via a more elaborate combinatorial computation in graph homology in [13].

As a consequence of the elementary construction, we find that all the classes μ_k in Morita's original series, as well as the generalized Morita classes given in [12], are supported on abelian subgroups of $\text{Aut}(F_n)$. This naturally gives rise to the question of whether the standard maximal abelian subgroup can support nontrivial homology classes, and we show in Section 8 that it cannot. For the Eisenstein classes we find slightly more complicated nonabelian subgroups that support them.

Parallel to these group-theoretic descriptions of Morita and Eisenstein classes there are geometric representations as maps of closed orientable manifolds into the classifying spaces for $\text{Aut}(F_n)$ or $\text{Out}(F_n)$ carrying top-dimensional homology classes of the manifolds to the Morita or Eisenstein classes. In the case of Morita classes the manifolds are tori while for the Eisenstein classes they are products of a certain 3-manifold with tori.

The computational heart of the paper is in Section 3 where we use the natural action of the symmetric group \mathfrak{S}_s on $\Gamma_{n,s}$ to study $H_*(\Gamma_{n,s})$. For $n = 1$ and $n = 2$ we determine the \mathfrak{S}_s -module structure of $H_*(\Gamma_{n,s})$ completely. This can be applied in the search for nontrivial classes in $H_*(\Gamma_{n,s})$ that lie in the images of assembly maps. In particular we show in Section 5.5 that many of the generalized Morita classes are in fact zero, though this does not apply to the μ_k 's themselves. In Section 5.7 we show that certain odd-dimensional classes constructed in [35] must vanish, but we also find some new candidates for nontrivial odd-dimensional classes.

The calculation of $H_*(\Gamma_{1,s})$ is an easy consequence of the fact that $\Gamma_{1,s} \cong \mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1}$. To calculate the homology of $\Gamma_{2,s}$ we use the short exact sequence

$$1 \longrightarrow F_2^s \longrightarrow \Gamma_{2,s} \longrightarrow \Gamma_{2,0} = \text{Out}(F_2) \longrightarrow 1.$$

In the Leray–Serre spectral sequence associated to this short exact sequence we note that all differentials are zero, allowing us to completely calculate the homology. (Actually, for convenience we use cohomology rather than homology for spectral sequence arguments and indeed for most algebraic calculations.) The results of our computations for $n = 1$ and $n = 2$ and small values of s are recorded in several tables at the end of the paper.

These computations show that even though the dimension of $H_i(\Gamma_{n,s})$ as a vector space over \mathbb{k} increases with s for fixed $n = 1, 2$, it is nevertheless true that as representations of \mathfrak{S}_s these vector spaces eventually stabilize in the sense of [7]. This representation stability holds for all n in fact, as a corollary of a result of Jiménez Rolland [26]; see Proposition 6.1.

We also apply some elementary representation theory to show that the group $H_i(\Gamma_{n,s})$ is nontrivial whenever i is an even multiple of n and s is sufficiently large

with respect to i and n . This can be contrasted with the situation for stabilization with respect to n , where $H_i(\Gamma_{n,s})$ becomes trivial as n increases, by Galatius' theorem for $s = 1$ and hence for all s since the groups $H_i(\Gamma_{n,s})$ are independent of both n and s when $n \geq 2i + 4$ by [24].

In Section 9 we point out the relationship of the homology of $\Gamma_{n,s}$ with both hairy graph homology [10, 11] and the cohomology of the Lie algebra of symplectic derivations, as studied by Kontsevich, Morita and many others. In particular, we show how our computations for $\Gamma_{n,s}$ imply that the cohomology in each dimension of this Lie algebra contains infinitely many simple Sp-modules.

Section 10 contains some conjectures and open questions. Most nontrivial rational homology classes for any $\Gamma_{n,s}$ which occur below the vcd $2n - 3 + s$ have been shown to be in the image of assembly maps. The only exceptions are new classes for $\text{Out}(F_7)$ and $\text{Aut}(F_7)$ recently found by Bartholdi [1], for which this is still unclear. It is then natural to ask whether assembly maps generate all classes below the vcd. The number of potential homology classes for $H_i(\Gamma_{n,s})$ constructed from assembly maps grows exponentially with n , leading to the expectation that the rank of the homology also grows very fast. For $s = 0$ this expectation coincides nicely with the Euler characteristic calculations of Morita, Sakasai, and Suzuki cited earlier.

Finally, we remark that the rational classifying spaces for the groups $\Gamma_{n,s}$ have natural compactifications, whose homology has recently been studied by Chan, Galatius and Payne [5, 6]. One thing they show is that this homology vanishes in dimensions less than $s - 3$. It is easy to see that all homology classes which are in the image of an assembly map must vanish in this compactification, consistent with their calculations.

2. The groups $\Gamma_{n,s}$

2.1. Definitions. The group $\text{Out}(F_n)$ is the group of homotopy classes of self-homotopy equivalences of a finite connected graph X of rank n , and $\text{Aut}(F_n)$ is the basepointed version of this, the homotopy classes of homotopy equivalences of X fixing a basepoint, where homotopies are also required to fix the basepoint. A natural generalization is to choose s distinct marked points x_1, \dots, x_s in X and then define $\Gamma_{n,s}$ to be the group of homotopy classes of self-homotopy equivalences of X fixing each x_i , with homotopies also required to fix these points. The group operation in $\Gamma_{n,s}$ is induced by composition of homotopy equivalences, which is obviously associative with an identity element. To check that inverses exist one uses the following elementary fact:

Lemma 2.1. *If $f: X \rightarrow Y$ is a homotopy equivalence of finite connected graphs taking a set of s marked points $\mathbf{x} = \{x_1, \dots, x_s\}$ bijectively to another such set $\mathbf{y} = \{y_1, \dots, y_s\}$, then f is a homotopy equivalence of pairs $(X, \mathbf{x}) \rightarrow (Y, \mathbf{y})$, so*

there is a map $g : Y \rightarrow X$ restricting to f^{-1} on \mathbf{y} with the compositions gf and fg homotopic to the identity fixing \mathbf{x} and \mathbf{y} respectively.

Proof. Let Z be the quotient of the mapping cylinder of f obtained by collapsing $\mathbf{x} \times I$ to $\mathbf{x} = \mathbf{y}$. The quotient map collapses a finite number of intervals to a point so it is a homotopy equivalence. If f is a homotopy equivalence, then the inclusions of X and Y into the mapping cylinder are homotopy equivalences, hence the same is true for the inclusions into Z . It follows that Z deformation retracts onto the copies of X and Y at either end. The deformation retraction to X gives the map g . \square

This lemma also shows that $\Gamma_{n,s}$ does not depend on the choice of (X, \mathbf{x}) , up to isomorphism. Throughout most of the paper we will take X to be a rank n graph with exactly s leaves, with the leaf vertices as the marked points. Here a *leaf* means a vertex of valence one together with the adjoining edge. Our generic notation for a graph of rank n with s leaves will be $X_{n,s}$. Two examples of rank 3 graphs with 4 leaves are shown in Figure 3.




Figure 3. Two possibilities for $X_{3,4}$

A homotopy equivalence $f : X_{n,s} \rightarrow X_{n,s}$ that permutes the leaf vertices induces an automorphism of $\Gamma_{n,s}$ via conjugation by f . If f fixes the leaf vertices this is an inner automorphism, hence induces the identity on the homology of $\Gamma_{n,s}$, so there is an induced action of the symmetric group \mathfrak{S}_s on the homology. If we choose $X_{n,s}$ to have a single nonleaf vertex as in the left half of Figure 3 then this action of \mathfrak{S}_s on homology comes from the action on $X_{n,s}$ permuting the leaves. The \mathfrak{S}_s -action on $H_*(\Gamma_{n,s})$ will play a major role in later sections of the paper.

The groups $\Gamma_{0,s}$ are trivial since any $X_{0,s}$ is a tree and any homotopy equivalence of a tree which fixes all of its leaf vertices is homotopic to the identity by a homotopy fixing the leaf vertices.

As shown in [2], the group $\Gamma_{1,s}$ is the semidirect product $\mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1}$. The free abelian subgroup \mathbb{Z}^{s-1} is generated by homotopy equivalences which wrap one leaf edge around the (unique) loop while fixing the leaf vertex and the rest of the graph. These generators commute since they have disjoint supports. Note that wrapping all of the leaf edges around the loop in the same direction results in a map which is homotopic to the identity fixing the leaf vertices, so there are only $s - 1$ independent generators. The generator of \mathbb{Z}_2 flips the loop, so acts on \mathbb{Z}^{s-1} by $x \mapsto -x$.

We remark that $\Gamma_{n,s}$ could also be defined as the mapping class group of the 3-manifold $M_{n,s}$ formed by removing s disjoint balls from the connected sum of n copies of $S^1 \times S^2$, modulo the subgroup generated by Dehn twists along embedded 2-spheres. This follows from results of Laudenbach and is made explicit in Proposition 1 of [24].

The groups $\Gamma_{n,s}$ for $s > 1$ were first considered in [22] in work on homological stability and also appeared in Bestvina and Feighn's proof that $\text{Out}(F_n)$ is a virtual duality group [2]. It was observed in [10] that their homology is very closely related to hairy graph homology groups for the Lie operad.

2.2. Short exact sequences. In this section we observe that there are natural short exact sequences relating the groups $\Gamma_{n,s}$.

Proposition 2.2. *If $n > 1$ and $k \leq s$ there is a short exact sequence*

$$1 \longrightarrow F_n^k \longrightarrow \Gamma_{n,s} \longrightarrow \Gamma_{n,s-k} \longrightarrow 1$$

which splits if $k < s$. This holds also when $n = 1$ and $k < s$, but in the exceptional case $(n, k) = (1, s)$ there is a split short exact sequence

$$1 \longrightarrow \mathbb{Z}^{s-1} \longrightarrow \Gamma_{1,s} \longrightarrow \Gamma_{1,0} \longrightarrow 1$$

expressing $\Gamma_{1,s}$ as the semidirect product $\mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1}$.

For $k = s - 1$ the proposition follows from [2, Section 2.5], where it is shown that $\Gamma_{n,s} \cong \text{Aut}(F_n) \ltimes F_n^{s-1}$.

Proof. Let X be a rank n graph containing a set $\mathbf{x} = \{x_1, \dots, x_s\}$ of s distinct marked points. Let $E_{n,s}$ be the space of homotopy equivalences $X \rightarrow X$ fixed on \mathbf{x} , so $\Gamma_{n,s} = \pi_0(E_{n,s})$. For $k \leq s$ there is an inclusion $E_{n,s} \subset E_{n,s-k}$ obtained by no longer requiring homotopy equivalences to fix x_1, \dots, x_k . Evaluating homotopy equivalences $X \rightarrow X$ on x_1, \dots, x_k gives a map $E_{n,s-k} \rightarrow X^k$ which is a fibration with fiber $E_{n,s}$ over the point (x_1, \dots, x_k) . The long exact sequence of homotopy groups for this fibration ends with the terms

$$\pi_1(E_{n,s-k}) \longrightarrow \pi_1(X^k) \longrightarrow \Gamma_{n,s} \longrightarrow \Gamma_{n,s-k} \longrightarrow 1.$$

When $k < s$ the first term $\pi_1(E_{n,s-k})$ is trivial by obstruction theory. Namely, we can assume X is obtained by attaching 1-cells to a set of $s - k$ 0-cells, and then any loop of homotopy equivalences $f_t : X \rightarrow X$ fixing the 0-cells can be deformed to the trivial loop since $\pi_2(X) = 0$. Thus we obtain the first short exact sequence in statement of the proposition when $k < s$, for arbitrary n .

To split this short exact sequence when $k < s$ it suffices to find a map $E_{n,s-k} \rightarrow E_{n,s}$ such that the composition $E_{n,s-k} \rightarrow E_{n,s} \rightarrow E_{n,s-k}$ is homotopic to the identity. We are free to choose the marked points x_1, \dots, x_k anywhere

in the complement of the remaining $s - k$ points, so we choose them in a small contractible neighborhood N of the point x_{k+1} . We can then deformation retract $E_{n,s-k}$ onto the subspace $E'_{n,s-k}$ of homotopy equivalences that are fixed on N . (This is particularly easy if we choose X to have a valence one vertex with x_{k+1} as this vertex.) The subspace $E'_{n,s-k}$ includes naturally into $E_{n,s}$, and this inclusion gives the desired map $E_{n,s-k} \rightarrow E_{n,s}$ as the composition of the first two maps $E_{n,s-k} \rightarrow E'_{n,s-k} \hookrightarrow E_{n,s} \hookrightarrow E_{n,s-k}$, the first map being the retraction produced by the deformation retraction. The composition of the three maps is homotopic to the identity by the deformation retraction itself.

There remain the cases $k = s$. The issue is whether $\pi_1(E_{n,0})$ is trivial, so that the long exact sequence becomes a short exact sequence. To settle this, consider the fibration $E_{n,1} \rightarrow E_{n,0} \rightarrow X$ which gives a long exact sequence

$$1 \longrightarrow \pi_1(E_{n,0}) \longrightarrow \pi_1(X) \longrightarrow \Gamma_{n,1} \longrightarrow \Gamma_{n,0} \longrightarrow 1$$

where the initial 1 is $\pi_1(E_{n,1})$. The middle map in this sequence is the map from π_1 of the base of the fibration to π_0 of the fiber, and it is easy to check the definitions to see that this is the map $F_n \rightarrow \text{Aut}(F_n)$ sending an element of F_n to the inner automorphism it determines. The kernel of this map is the center of F_n so it is trivial when $n > 1$ and we deduce that $\pi_1(E_{n,0}) = 1$ in these cases, so we again have the short exact sequence claimed in the proposition.

When $n = 1$ and $k = s$ the space $E_{1,0}$ is homotopy equivalent to S^1 and the exact sequence of the fibration $E_{1,s} \rightarrow E_{1,0} \rightarrow X^s$ becomes

$$1 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}^s \longrightarrow \Gamma_{1,s} \longrightarrow \Gamma_{1,0} \longrightarrow 1,$$

with the map $\mathbb{Z} \rightarrow \mathbb{Z}^s$ the diagonal inclusion. This yields the short exact sequence displaying $\Gamma_{1,s}$ as the semidirect product $\mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1}$. \square

Remark 2.3. If we use Laudenbach's theorem to express $\Gamma_{n,s}$ in terms of the mapping class group of $M_{n,s}$ then the short exact sequence of Proposition 2.2 can be derived from a 3-dimensional analog of the Birman exact sequence for mapping class groups of surfaces [2]. From this viewpoint the space $E_{n,s}$ is replaced by the diffeomorphism group of $M_{n,s}$, and the resulting fibration is a very simple special case of much more general fibrations due to Cerf, Palais, and Lima.

2.3. Homology splitting. If $s = k$ and $n \geq 2$ the map $\Gamma_{n,s} \rightarrow \Gamma_{n,s-k} = \Gamma_{n,0}$ does not split. The reason is that $\Gamma_{n,0} = \text{Out}(F_n)$ contains finite subgroups which do not lift. For example, consider the symmetry group of the graph consisting of two vertices joined by $n + 1$ edges. This is a subgroup of $\Gamma_{n,0}$ which cannot be realized on any graph of rank n by graph symmetries which fix a basepoint, so the subgroup does not lift to any $\Gamma_{n,s}$ with $s \geq 1$.

Homology with coefficients in \mathbb{k} does not see finite subgroups, and in fact when we pass to homology we do obtain a splitting. Note that it suffices to prove this for

$s = 1$, i.e. the map from $\text{Aut}(F_n)$ to $\text{Out}(F_n)$. Homology splitting of this map was first proved by Kawazumi in [27]. There is a simple proof of this using the fact that the moduli space of graphs (respectively basepointed graphs) is a rational $K(\pi, 1)$ for $\text{Out}(F_n)$ (respectively $\text{Aut}(F_n)$). The idea is that although there is no natural way to choose a basepoint in a graph one can compensate by taking a suitably weighted sum of all possible basepoints.

Theorem 2.4. *The natural map $\text{Aut}(F_n) \rightarrow \text{Out}(F_n)$ splits on the level of rational homology, so $H_k(\text{Out}(F_n))$ embeds into $H_k(\text{Aut}(F_n))$.*

Proof. We define a backwards map on the chain level. We take $C_*(\text{Aut}(F_n))$ and $C_*(\text{Out}(F_n))$ to be defined in terms of the spine of the moduli space of (basepointed) graphs. We refer to [13, section 2] for complete details. The chain complex $C_*(\text{Aut}(F_n))$ is spanned by graphs with specified subforests and a chosen basepoint, while $C_*(\text{Out}(F_n))$ is defined in the same way except the graphs do not have basepoints. In both cases, the edges in the subforests are ordered, and changing the order incurs the sign of the permutation. There are two boundary operators ∂_C and ∂_R which sum over contracting and removing forest edges respectively. In both the basepointed and unbasepointed cases, contracting the i th edge of a forest comes with the sign $(-1)^{i+1}$, while removing that edge comes with the sign $(-1)^i$.

The natural projection $\pi: \text{Aut}(F_n) \rightarrow \text{Out}(F_n)$ corresponds to the map

$$\pi_*: C_*(\text{Aut}(F_n)) \rightarrow C_*(\text{Out}(F_n))$$

which forgets the basepoint.

We now define a map $r: C_*(\text{Out}(F_n)) \rightarrow C_*(\text{Aut}(F_n))$, by

$$r(G) = \sum_{v \in V(G)} (|v| - 2)r_v(G).$$

Here $V(G)$ is the vertex set of G , $|v|$ is the valence of v and $r_v(G)$ is the forested graph G with v specified as the basepoint. We need to check that r is a chain map. Clearly $\partial_R r = r \partial_R$ since the definition of r makes no reference to the forest, and the signs in ∂_R make no reference to the basepoint. For ∂_C we must check whether the order of performing the two operations of adding a basepoint and contracting an edge matters, the signs in ∂_C being the same in both cases. If e is an edge with vertices v and w , then adding a basepoint distinct from v and w clearly commutes with contracting e . Adding basepoints at v and at w followed by contracting e results in the same basepointed graph with multiplicity $|v| + |w| - 4$, whereas contracting e first results in a vertex vw of valence $|v| + |w| - 2$, so adding a basepoint there also gives multiplicity $|v| + |w| - 4$ (see Figure 4).

Now observe that $\pi_* \circ r(G) = k_G G$, where $k_G = \sum_{v \in V(G)} (|v| - 2) = 2n - 2$. Thus if we are not in the trivial case $n = 1$ the composition $\pi_* \circ r$ is represented by a diagonal matrix with nonzero diagonal entries and is therefore invertible. So r_* is injective on homology. \square




Figure 4. Diagram commutes because $|vw| = |v| + |w| - 2$

3. Cohomology of $\Gamma_{n,s}$

We are interested in studying the homology of $\text{Out}(F_n)$ and $\text{Aut}(F_n)$, with trivial coefficients in a field \mathbb{k} of characteristic 0. The idea is to glue together homology classes of the $\Gamma_{n,s}$ using the assembly maps described briefly in the Introduction and defined more precisely in Section 5. To find nontrivial classes which can be fed to the assembly maps we use some elementary representation theory of symmetric groups and $\text{GL}_n(\mathbb{Z})$ together with the Leray–Serre spectral sequence applied to the group extensions

$$1 \longrightarrow F_n^s \longrightarrow \Gamma_{n,s} \longrightarrow \Gamma_{n,0} = \text{Out}(F_n) \longrightarrow 1 \quad (3.1)$$

from Section 2.2.

For the calculations it will be convenient to switch from homology to cohomology, which is isomorphic by the universal coefficient theorem since we are taking coefficients in \mathbb{k} and all homology is finite-dimensional over \mathbb{k} . In the course of our study we will exploit the structure of $H_i(\Gamma_{n,s})$ as an \mathfrak{S}_s -module. Since all the modules we consider are finite-dimensional and all \mathfrak{S}_s -modules are self-dual, the cohomology is isomorphic to the homology also as an \mathfrak{S}_s -module, though the isomorphism is not canonical.

3.1. A little representation theory. In this section we establish some notation and collect some results from representation theory which we will use. All the contents of this section are well known and can be found, for example, in [15].

Recall that the irreducible representations of \mathfrak{S}_s correspond to partitions of s and are often represented by drawing Young diagrams with s boxes arranged in rows of non-increasing size. We use P_λ to denote the representation corresponding to the partition $\lambda = (\lambda_1, \dots, \lambda_k)$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$ and $\sum_i \lambda_i = s$. Exponential notation denotes equal values of λ_i , e.g., $P_{(2,1,1,1,1)}$ is written as $P_{(2,1^4)}$.

Example 3.1. The module $P_{(s)}$ is the 1-dimensional trivial module. The module $P_{(1^s)} = \text{alt}$ is the 1-dimensional *alternating* representation of \mathfrak{S}_s , where a permutation σ acts as multiplication by $\text{sign}(\sigma) = \pm 1$. The module $P_{(s-1,1)}$ is the $(s-1)$ -dimensional *standard* representation $\mathbb{k}^s / \mathbb{k}$ of \mathfrak{S}_s . It contains distinguished elements v_i , $1 \leq i \leq s$, which satisfy $\sum v_i = 0$.

The tensor product of two \mathfrak{S}_s -representations is also an \mathfrak{S}_s -representation with the diagonal action. In general the multiplicity of an irreducible representation P_ν in the decomposition of $P_\lambda \otimes P_\mu$ is difficult to compute, but for $\nu = (s)$ it is known that $P_{(s)}$ occurs with multiplicity 1 if $\lambda = \mu$ and with multiplicity 0 otherwise. One tensor product we will encounter is $P_\lambda \otimes \text{alt}$. This is equal to $P_{\lambda'}$ where λ' denotes the *transpose partition*, obtained by switching the rows and columns of the Young diagram.

If P is a representation of \mathfrak{S}_{s-k} and Q is a representation of \mathfrak{S}_k , then $P \otimes Q$ is a representation of $\mathfrak{S}_{s-k} \times \mathfrak{S}_k$. If we consider $\mathfrak{S}_{s-k} \times \mathfrak{S}_k$ as a subgroup of \mathfrak{S}_s we can form the induced representation. Following Fulton and Harris [15], we denote this induced representation by $P \circ Q$, i.e.,

$$P \circ Q = \text{Ind}_{\mathfrak{S}_{s-k} \times \mathfrak{S}_k}^{\mathfrak{S}_s} P \otimes Q.$$

The Littlewood–Richardson rule can be used to compute the decomposition of $P_\lambda \circ P_\mu$ into irreducible modules. When $\mu = (k)$ this specializes to the *Pieri rule*, see [15, Appendix A]. This says that the terms of $P_\lambda \circ P_{(k)}$ correspond to all Young diagrams which can be obtained by adding k boxes to the diagram for λ , each in a different column. An example is illustrated in Figure 5.




Figure 5. Pieri rule for decomposing $P_{(2,2)} \circ P_{(3)}$

Now let V be an n -dimensional vector space. The irreducible representations of $\text{GL}(V) = \text{GL}_n(\mathbb{k})$ also correspond to partitions, and we let $\mathbb{S}_\lambda V$ denote the GL_n -representation associated to the partition λ . Since $\dim(V) = n$ only partitions into at most n pieces occur. *Schur–Weyl duality* gives the irreducible decomposition

of the representation $V^{\otimes q}$ as a module over $\mathrm{GL}(V) \times \mathfrak{S}_q$, namely

$$V^{\otimes q} \cong \bigoplus_{\lambda} \mathbb{S}_{\lambda} V \otimes P_{\lambda},$$

where the sum is over all partitions of q into at most n pieces (if λ has more than n rows the module $\mathbb{S}_{\lambda} V$ is zero) (see, e.g., [15, Cor. 6.6]). We emphasize that $\mathrm{GL}(V)$ acts trivially on P_{λ} and \mathfrak{S}_q acts trivially on $\mathbb{S}_{\lambda} V$.

Example 3.2. For $q = 2$ the Schur–Weyl formula gives

$$V \otimes V = (\mathbb{S}_{(2)} V \otimes P_{(2)}) \oplus (\mathbb{S}_{(1^2)} V \otimes P_{(1^2)}) = \mathrm{Sym}^2 V \oplus \bigwedge^2 V.$$

where Sym^k denotes the k -th symmetric power functor on vector spaces and \bigwedge^k is the k -th exterior power.

Notation. We denote by $V^{\wedge q}$ the \mathfrak{S}_q -module which is isomorphic as a vector space to $V^{\otimes q}$, with \mathfrak{S}_q acting by permuting the factors and multiplying by the sign of the permutation, i.e.,

$$V^{\wedge q} = V^{\otimes q} \otimes \mathrm{alt}.$$

The Schur–Weyl formula translates to a similar formula for $V^{\wedge q}$:

$$V^{\wedge q} = V^{\otimes q} \otimes \mathrm{alt} \cong \bigoplus_{\lambda} \mathbb{S}_{\lambda} V \otimes P_{\lambda} \otimes \mathrm{alt} = \bigoplus_{\lambda} \mathbb{S}_{\lambda} V \otimes P_{\lambda'},$$

where the sum is over all partitions of q into at most n pieces.

Finally, we record a computation we will use later.

Lemma 3.3. Suppose $\dim(V) = 2$. Then $\mathbb{S}_{(q-k,k)} V \cong \mathrm{Sym}^{q-2k} V \otimes \det^k$ as $\mathrm{GL}(V)$ -modules, where $\det^k = (\bigwedge^2 V)^{\otimes k}$ is the 1-dimensional representation given by the k -th power of the determinant.

Proof. This can be seen by calculating the Schur polynomials S_{λ} for the two sides, which determine the representations uniquely. Using the formula A.4 of [15, Appendix A] one obtains

$$S_{(a,b)} = (x_1 x_2)^b \left[\frac{x_1^{a-b+1} - x_2^{a-b+1}}{x_1 - x_2} \right] = (S_{(1,1)})^b S_{(a-b)}.$$

The lemma now follows because Schur polynomials of tensor products multiply, $\mathbb{S}_{(1,1)} H = \det$ and $\mathbb{S}_{(c)} V = \mathrm{Sym}^c V$. \square

3.2. The Leray–Serre spectral sequence. Shifting from homology to cohomology now, the Leray–Serre spectral sequence of a group extension $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ is a first-quadrant spectral sequence with $E_2^{p,q} = H^p(Q; H^q(N))$, which converges to $H^{p+q}(G)$. Applied to the short exact sequence (3.1) it reads

$$E_2^{p,q} = H^p(\mathrm{Out}(F_n); H^q(F_n^s)) \implies H^{p+q}(\Gamma_{n,s}). \quad (3.2)$$

The symmetric group \mathfrak{S}_s which permutes the factors of F_n^s induces an action on each of the E_2 terms which commutes with all differentials. We begin by identifying the coefficients $H^q(F_n^s)$ as \mathfrak{S}_s -modules.

Throughout this section we set $H = H^1(F_n) \cong \mathbb{k}^n$. Note that the action of $\text{Out}(F_n)$ on H factors through the natural $\text{GL}_n(\mathbb{Z})$ action on H .

Lemma 3.4. *The cohomology of F_n^s is given as an \mathfrak{S}_s -module by the formula*

$$H^q(F_n^s) = H^{\wedge q} \circ P_{(s-q)}.$$

Proof. The Künneth formula gives an isomorphism

$$H^*(F_n) \otimes \cdots \otimes H^*(F_n) \cong H^*(F_n \times \cdots \times F_n)$$

via the cohomology cross product. The group \mathfrak{S}_s acts by permuting the factors, with signs determined by the permutation and the dimension of the cohomology groups on the left-hand side (see, e.g., [21, Chapter 3B]). The cohomology of F_n is \mathbb{k} in dimension 0, H in dimension 1 and zero in higher dimensions, so in dimension q the cohomology of F_n^s is the direct sum of $\binom{s}{q}$ copies of $H^{\otimes q}$. These copies are permuted by the action of \mathfrak{S}_s . The stabilizer of each copy is isomorphic to $\mathfrak{S}_q \times \mathfrak{S}_{s-q}$, where the action of \mathfrak{S}_q on $H^{\otimes q}$ is modified by the sign of the permutation since all classes are in dimension 1.

In other words, $H^q(F_n^s)$ is obtained by inducing up to \mathfrak{S}_s the $\mathfrak{S}_q \times \mathfrak{S}_{s-q}$ -module $H^{\wedge q} \otimes P_{(s-q)}$. \square

We now read off information which we obtain immediately from the spectral sequence (3.2). The first observation applies to the case $s = 1$. The same result was obtained earlier by Kawazumi [27] using a different method.

Proposition 3.5. *There is an isomorphism*

$$H^k(\text{Aut}(F_n); \mathbb{k}) \cong H^k(\text{Out}(F_n); \mathbb{k}) \oplus H^{k-1}(\text{Out}(F_n); H).$$

Proof. In the spectral sequence associated to $1 \rightarrow F_n \rightarrow \text{Aut}(F_n) \rightarrow \text{Out}(F_n) \rightarrow 1$ we have $E_2^{p,q} \cong H^p(\text{Out}(F_n); H^q(F_n))$ with differentials of bidegree $(2, -1)$. Since F_n has cohomological dimension one there are only two nontrivial rows, namely $q = 0$ and $q = 1$, so the only possible nonzero differentials in the entire spectral sequence are on the E_2 page; they start in the top row $q = 1$ with target in the bottom row $q = 0$.

The map on cohomology induced by $p: \text{Aut}(F_n) \rightarrow \text{Out}(F_n)$ factors through the edge homomorphism $e: E_\infty^{p,0} \rightarrow H^p(\text{Aut}(F_n))$

$$\begin{array}{ccc} H^p(\text{Out}(F_n)) & \xrightarrow{p^*} & H^p(\text{Aut}(F_n)) \\ & \searrow & \nearrow e \\ & H^p(\text{Out}(F_n))/\text{Im}(d^2) & \end{array}$$

The top arrow is injective by Theorem 2.4, so the left arrow is as well, i.e., $\text{Im}(d^2) = 0$ and the differentials on the E_2 page are also trivial. Thus

$$\begin{aligned} H^k(\text{Aut}(F_n)) &= E_2^{k,0} \oplus E_2^{k-1,1} \\ &= H^k(\text{Out}(F_n)) \oplus H^{k-1}(\text{Out}(F_n); H^1(F_n)) \\ &= H^k(\text{Out}(F_n)) \oplus H^{k-1}(\text{Out}(F_n); H). \end{aligned} \quad \square$$

The next observation has to do with the top-dimensional cohomology of $\Gamma_{n,s}$.

Proposition 3.6. $H^k(\Gamma_{n,s})$ vanishes for $k > 2n - 3 + s$ and $H^{2n-3+s}(\Gamma_{n,s})$ is given as an \mathfrak{S}_s -module by

$$H^{2n-3+s}(\Gamma_{n,s}) \cong H^{2n-3}(\text{Out}(F_n); H^{\wedge s}).$$

Proof. The cohomology group $H^p(\text{Out}(F_n); H^q(F_n^s))$ vanishes if either $p > 2n - 3$ or $q > s$ since the virtual cohomological dimension of $\text{Out}(F_n)$ is $2n - 3$ and the virtual cohomological dimension of F_n^s is equal to s . Thus the only possible nonzero terms in the spectral sequence (3.2) lie in a rectangle with $E_2^{2n-3,s}$ at its upper right-hand corner, so all differentials into or out of $E_2^{2n-3,s}$ are zero and

$$H^{2n-3+s}(\Gamma_{n,s}) \cong H^{2n-3}(\text{Out}(F_n); H^s(F_n^s)) \cong H^{2n-3}(\text{Out}(F_n); H^{\wedge s}). \quad \square$$

3.3. Rank zero. Since $\Gamma_{0,s}$ is trivial, we just have

$$H^i(\Gamma_{0,s}) = \begin{cases} P_{(s)} = \mathbb{k} & \text{if } i = 0 \\ 0 & \text{if } i \neq 0. \end{cases}$$

3.4. Rank one. For $n = 1$ the short exact sequence (3.1) is a restatement of the fact that

$$\Gamma_{1,s} = \mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1},$$

where the \mathbb{Z}_2 acts via $x \mapsto -x$. We can use this to compute the cohomology of $\Gamma_{1,s}$ as an \mathfrak{S}_s -module without appealing to the Leray–Serre spectral sequence, as follows.

Proposition 3.7. As a representation of \mathfrak{S}_s

$$H^i(\Gamma_{1,s}) \cong \begin{cases} P_{(s-i,1^i)} & \text{if } i \text{ is even} \\ 0 & \text{if } i \text{ is odd.} \end{cases}$$

In particular, $H^{2k}(\Gamma_{1,2k+1}) \cong P_{(1^{2k+1})} = \mathbb{k}$ with the alternating action.

Proof. The rational cohomology of $\Gamma_{1,s} = \mathbb{Z}_2 \ltimes \mathbb{Z}^{s-1}$ is the invariants of the \mathbb{Z}_2 -action on the cohomology of \mathbb{Z}^{s-1} induced from the action on \mathbb{Z}^{s-1} . The

cohomology of \mathbb{Z}^{s-1} is the exterior algebra on $s - 1$ generators. Thus we get the even degree part of this exterior algebra:

$$H^i(\Gamma_{1,s}) \cong \begin{cases} \bigwedge^i \mathbb{k}^{s-1} & \text{if } i \text{ is even} \\ 0 & \text{if } i \text{ is odd.} \end{cases}$$

To see the \mathfrak{S}_s action, write $\Gamma_{1,s} = \mathbb{Z}_2 \ltimes (\mathbb{Z}^s/\mathbb{Z})$. The representation \mathbb{k}^s/\mathbb{k} is the standard representation $P_{(s-1,1)}$ of \mathfrak{S}_s , and by [15, Ex. 4.6] we have $\bigwedge^i (\mathbb{k}^s/\mathbb{k}) = P_{(s-i,1^i)}$ as an \mathfrak{S}_s -module. \square

We record this calculation for small values of s in table form at the end of the paper (Section 11). We note that the results agree with the calculations via dihedral homology in [11].

3.5. Rank two. Recall that $\Gamma_{2,0} = \text{Out}(F_2) \cong \text{GL}_2(\mathbb{Z})$, so that for $n = 2$ the E_2 term of (3.2) is

$$E_2^{p,q} = H^p(\text{GL}_2(\mathbb{Z}); H^q(F_2^s)).$$

Since $\text{GL}_2(\mathbb{Z})$ has virtual cohomological dimension 1, the only potentially nonzero terms on the E_2 -page of this spectral sequence lie in the first two columns $p = 0$ and $p = 1$. For $p = 1$ the cohomology of $\text{GL}_2(\mathbb{Z})$ is closely related to modular forms; we review this relation in the next subsection.

3.5.1. Modular forms. Let \mathcal{M}_s be the vector space of classical modular forms for $\text{SL}_2(\mathbb{Z})$ of weight s , and let $\mathcal{S}_s \subset \mathcal{M}_s$ be the subspace of cusp forms. See [31] for an elementary introduction to these spaces. They satisfy

$$\bigoplus_{s \geq 0} \mathcal{M}_s \cong \mathbb{k}[E_4, E_6],$$

where E_4 and E_6 are generators of weight 4 and 6 respectively. In particular, \mathcal{M}_s is nonzero only for $s > 2$ even. In these cases the subspace \mathcal{S}_s has codimension 1. The classical Eichler–Shimura isomorphism (see, e.g., [20]) relates modular forms to the cohomology of $\text{SL}_2(\mathbb{Z})$:

$$H^1(\text{SL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2)) \cong \mathcal{M}_{s+2} \oplus \mathcal{S}_{s+2}.$$

We next review the relation between cusp forms and the stabilizer of the cusp at infinity. Let $P \leq \text{SL}_2(\mathbb{Z})$ be the (parabolic) subgroup generated by the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and consider the map

$$\rho: H^1(\text{SL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2)) \longrightarrow H^1(P; \text{Sym}^s(\mathbb{k}^2))$$

induced by inclusion. Since $P \cong \mathbb{Z}$, its first cohomology with any coefficients is simply the coinvariants of the action, which is isomorphic to the space of

invariants. If x and y are a basis for \mathbb{k}^2 , the generator of P acts on $\text{Sym}^s(\mathbb{k}^2) = \mathbb{k}[x^s, x^{s-1}y, \dots, xy^{s-1}, y^s]$ by sending $x \mapsto x$ and $y \mapsto x + y$, so the space of invariants is 1-dimensional, spanned by x^s . The map ρ can be identified with the map $\mathcal{M}_{s+2} \oplus \mathcal{S}_{s+2} \rightarrow \mathcal{M}_{s+2}/\mathcal{S}_{s+2}$ (projection on the first factor, zero on the second factor) given by the normalized value of the modular form at infinity (see [20]).

We can reinterpret ρ in terms of the cohomology of $\text{GL}_2(\mathbb{Z})$ using the short exact sequence $1 \rightarrow \text{SL}_2(\mathbb{Z}) \rightarrow \text{GL}_2(\mathbb{Z}) \rightarrow \mathbb{Z}_2 \rightarrow 1$ (see, e.g., [10]); this gives

$$H^1(\text{GL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2) \otimes \det) \cong \mathcal{M}_{s+2} \quad \text{and} \quad H^1(\text{GL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2)) \cong \mathcal{S}_{s+2}.$$

Since $H^1(\text{SL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2)) = \mathcal{M}_{s+2} \oplus \mathcal{S}_{s+2}$ we see that the restriction of ρ to the second factor

$$\rho: H^1(\text{GL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2)) \longrightarrow H^1(P; \text{Sym}^s(\mathbb{k}^2))$$

is zero, but on the first factor

$$\rho: H^1(\text{GL}_2(\mathbb{Z}); \text{Sym}^s(\mathbb{k}^2) \otimes \det) \longrightarrow H^1(P; \text{Sym}^s(\mathbb{k}^2) \otimes \det)$$

has 1-dimensional image when $s > 0$ is even.

3.5.2. Cohomology calculations.

Lemma 3.8. *Let $H = H^1(F_2) \cong \mathbb{k}^2$. Then as an \mathfrak{S}_q -module,*

$$H^0(\text{GL}_2(\mathbb{Z}); H^{\wedge q}) = \begin{cases} P_{(2^{2m})} & \text{if } q = 4m \\ 0 & \text{otherwise.} \end{cases}$$

$$H^1(\text{GL}_2(\mathbb{Z}); H^{\wedge q}) = \begin{cases} 0 & \text{if } q \text{ is odd} \\ W_q := \bigoplus_{0 \leq i < \frac{q}{2}} \mathcal{X}_{q,i} \otimes P_{(2^i, 1^{q-2i})} & \text{if } q \text{ is even} \end{cases}$$

where $\mathcal{X}_{q,i} = \mathcal{S}_{q+2-2i}$ if i is even and \mathcal{M}_{q+2-2i} if i is odd. In either case $\mathcal{X}_{q,i}$ is trivial as an \mathfrak{S}_q -module.

Remark 3.9. The formula in the statement above gives the following pattern for the first few W_q :

$$W_0 = 0$$

$$W_2 = (\mathcal{S}_4 \otimes P_{(1^2)})$$

$$W_4 = (\mathcal{S}_6 \otimes P_{(1^4)}) \oplus (\mathcal{M}_4 \otimes P_{(2,1^2)})$$

$$W_6 = (\mathcal{S}_8 \otimes P_{(1^6)}) \oplus (\mathcal{M}_6 \otimes P_{(2,1^4)}) \oplus (\mathcal{S}_4 \otimes P_{(2^2,1^2)})$$

$$W_8 = (\mathcal{S}_{10} \otimes P_{(1^8)}) \oplus (\mathcal{M}_8 \otimes P_{(2,1^6)}) \oplus (\mathcal{S}_6 \otimes P_{(2^2,1^4)}) \oplus (\mathcal{M}_4 \otimes P_{(2^3,1^2)})$$

$$W_{10} = (\mathcal{S}_{12} \otimes P_{(1^{10})}) \oplus (\mathcal{M}_{10} \otimes P_{(2,1^8)}) \oplus (\mathcal{S}_8 \otimes P_{(2^2,1^6)}) \oplus (\mathcal{M}_6 \otimes P_{(2^3,1^4)}) \\ \oplus (\mathcal{S}_4 \otimes P_{(2^4,1^2)}).$$

However, the dimension of \mathcal{M}_k is 1 for $k = 4, 6, 8, 10$, and hence the dimension of \mathcal{S}_k is trivial in those degrees. The module \mathcal{S}_{12} is 1-dimensional, so we see interesting modular forms entering the picture starting with W_{10} . Using this information, the above list simplifies to

$$\begin{aligned} W_0 &= 0 \\ W_2 &= 0 \\ W_4 &= P_{(2,1^2)} \\ W_6 &= P_{(2,1^4)} \\ W_8 &= P_{(2,1^6)} \oplus P_{(2^3,1^2)} \\ W_{10} &= P_{(1^{10})} \oplus P_{(2,1^8)} \oplus P_{(2^3,1^4)}. \end{aligned}$$

Proof. We first decompose the coefficients $H^{\wedge q}$ into irreducible components using Schur–Weyl duality. Since H has dimension 2, this gives $H^{\wedge q} \cong \bigoplus_{\lambda} \mathbb{S}_{\lambda} H \otimes P_{\lambda'}$, where the sum is over all partitions of q into at most 2 pieces, i.e., $\lambda = (q-k, k)$.

Now $H^0(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge q})$ is equal to the $\mathrm{GL}_2(\mathbb{Z})$ -invariants of $H^{\wedge q}$, so we are looking for the trivial representations $\mathbb{S}_{\lambda} H$ appearing in the Schur–Weyl formula. By Lemma 3.3 we have $\mathbb{S}_{(q-k,k)} H \cong \mathrm{Sym}^{q-2k} H \otimes \det^k$, which is clearly trivial only if $q = 2k$ and k is even. Therefore as an \mathfrak{S}_q -module we have

$$H^0(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge q}) = \begin{cases} P_{(2m,2m)'} = P_{(2^{2m})} & \text{if } q = 4m, \\ 0 & \text{otherwise.} \end{cases}$$

For the first cohomology we have

$$\begin{aligned} H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge q}) &= \bigoplus_{0 \leq k \leq \frac{q}{2}} H^1(\mathrm{GL}_2(\mathbb{Z}); \mathbb{S}_{(q-k,k)} H \otimes P_{(q-k,k)'}) \\ &= \bigoplus_{0 \leq k \leq \frac{q}{2}} H^1(\mathrm{GL}_2(\mathbb{Z}); \mathbb{S}_{(q-k,k)} H) \otimes P_{(q-k,k)'} \\ &= \bigoplus_{0 \leq k \leq \frac{q}{2}} H^1(\mathrm{GL}_2(\mathbb{Z}); \mathrm{Sym}^{q-2k} H \otimes \det^k) \otimes P_{(q-k,k)'} \\ &= \bigoplus_{0 \leq k \leq \frac{q}{2}} H^1(\mathrm{GL}_2(\mathbb{Z}); \mathrm{Sym}^{q-2k} H \otimes \det^k) \otimes P_{(2^k, 1^{q-2k})}. \end{aligned}$$

The computations in Section 3.5.1 now give

$$H^1(\mathrm{GL}_2(\mathbb{Z}); \mathrm{Sym}^r H \otimes \det^{\ell}) = \begin{cases} 0 & \text{if } r \text{ is odd,} \\ \mathcal{S}_{r+2} & \text{if } r \text{ and } \ell \text{ are even,} \\ \mathcal{M}_{r+2} & \text{if } r \text{ is even and } \ell \text{ is odd.} \end{cases}$$

Substituting $r = q - 2k$, $\ell = k$ into these formulas and using the fact that $\mathcal{M}_2 = 0$ completes the calculation. \square

We now have the tools we need to completely compute the cohomology of $\Gamma_{2,s}$ as an \mathfrak{S}_s -module.

Theorem 3.10. *The cohomology of $\Gamma_{2,s}$ is*

$$H^i(\Gamma_{2,s}) = \begin{cases} P_{(2^{2m})} \circ P_{(s-4m)} & i = 4m \leq s, \\ 0 & i = 4m + 2, \\ W_{2m} \circ P_{(s-2m)} & i = 2m + 1 \leq s + 1, \\ 0 & \text{otherwise,} \end{cases}$$

where W_{2m} is the module defined in the statement of Lemma 3.8.

Proof. For all $r \geq 2$ the differential on the r -th page of the Leray–Serre spectral sequence has bidegree $(r, -r + 1)$. Since only the first two columns are nonzero all differentials are too wide to be nonzero, so $E_2 = E_\infty$ and

$$H^k(\Gamma_{2,s}) \cong H^0(\mathrm{GL}_2(\mathbb{Z}); H^k(F_2^s)) \oplus H^1(\mathrm{GL}_2(\mathbb{Z}); H^{k-1}(F_2^s)).$$

By Lemmas 3.4 and 3.8 we have

$$\begin{aligned} H^0(\mathrm{GL}_2(\mathbb{Z}); H^k(F_2^s)) &\cong H^0(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge k} \circ P_{(s-k)}) \\ &\cong H^0(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge k}) \circ P_{(s-k)} \\ &\cong \begin{cases} P_{(2^{2m})} \circ P_{(s-4m)} & \text{if } k = 4m \leq s, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where the second isomorphism holds because the $\mathrm{GL}_2(\mathbb{Z})$ action on $H^{\wedge k}$ commutes with the $\mathfrak{S}_k \times \mathfrak{S}_{s-k}$ action and the \mathfrak{S}_{s-k} action is trivial.

We calculate the irreducible decomposition of $P_{(2^{2m})} \circ P_{(s-4m)}$ using the Pieri rule, which says that the components are obtained by adding $s - 4m$ boxes in different columns to the Young diagram for $\lambda = (2^{2m})$. The only legal way to do this is to put 0, 1, or 2 boxes in a new bottom row and add the rest to the first row. The resulting partitions are $(s - 4m - j, 2^{2m-1}, j)$ for $j = 0, 1, 2$.

The second summand is

$$H^1(\mathrm{GL}_2(\mathbb{Z}); H^{k-1}(F_2^s)) \cong H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge k-1}) \circ P_{(s-k+1)}.$$

By Lemma 3.8, $H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge k-1})$ is nonzero only when k is odd, in which case we have identified it as an \mathfrak{S}_{k-1} -module which we named W_{k-1} . Inducing this up to \mathfrak{S}_s produces $\binom{s}{k-1}$ copies of W_{k-1} , permuted by the action of \mathfrak{S}_s .

The first few rows and columns of the spectral sequence look like this:

| | | | |
|-----------------------------|-----------------------|---|---|
| | 0 | 0 | 0 |
| $P_{(2^4)} \circ P_{(s-8)}$ | $W_8 \circ P_{(s-8)}$ | | 0 |
| 0 | 0 | | 0 |
| 0 | $W_6 \circ P_{(s-6)}$ | | 0 |
| 0 | 0 | | 0 |
| $P_{(2^2)} \circ P_{(s-4)}$ | $W_4 \circ P_{(s-4)}$ | | 0 |
| 0 | 0 | | 0 |
| 0 | $W_2 \circ P_{(s-2)}$ | | 0 |
| 0 | 0 | | 0 |
| $P_{(s)}$ | 0 | | 0 |

Since $E_2 = E_\infty$, the result follows. \square

Remark 3.11. The dimension of $P_{(2^{2m})}$ can be computed by the hook-length formula (see, e.g., [15]); it is the $2m$ -th Catalan number $C_{2m} = \frac{1}{2m+1} \binom{4m}{2m}$. The induced representation $P_{(2^{2m})} \circ P_{(s-4m)} = H^{4m}(\Gamma_{2,s})$ consists of $\binom{s}{4m}$ copies of this, so has dimension equal to

$$\frac{1}{2m+1} \binom{4m}{2m} \binom{s}{4m} = \frac{s!}{(s-4m)!(2m+1)!(2m)!}.$$

If $s \geq 4m+2$ then the irreducible decomposition of $P_{(2^{2m})} \circ P_{(s-4m)}$ obtained by the Pieri rule is

$$P_{(2^{2m})} \circ P_{(s-4m)} = P_{(s-4m+2, 2^{2m-1})} \oplus P_{(s-4m+1, 2^{2m-1}, 1)} \oplus P_{(s-4m, 2^{2m})}$$

Remark 3.12. Using the decomposition of W_q into irreducible \mathfrak{S}_q -modules in Remark 3.9 one can use the Pieri rule to obtain the decomposition of $H^i(\Gamma_{2,s})$ into irreducible \mathfrak{S}_s -modules for odd i . For example

$$H^7(\Gamma_{2,10}) = W_6 \circ P_{(4)} = P_{(2,1^4)} \circ P_{(4)} = P_{(6,1^4)} \oplus P_{(5,2,1^3)} \oplus P_{(5,1^5)} \oplus P_{(4,2,1^4)}.$$

The dimension and module structure of the cohomology of $\Gamma_{1,s}$ and $\Gamma_{2,s}$ for $s \leq 10$ are summarized in the tables at the end of the paper.

Remark 3.13. The calculation of the map on cohomology induced by inclusion $P \rightarrow \mathrm{GL}_2(\mathbb{Z})$ in Section 3.5.1 together with the decomposition

$$H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge q}) = \bigoplus_{0 \leq k \leq \frac{q}{2}} H^1(\mathrm{GL}_2(\mathbb{Z}); \mathrm{Sym}^{q-2k} H \otimes \det^k) \otimes P_{(2^k, 1^{q-2k})}.$$

given in the proof of Lemma 3.8 shows that the image of the map

$$H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge q}) \longrightarrow H^1(P; H^{\wedge q})$$

is isomorphic to $\bigoplus_{2k < q, k \text{ odd}} P_{(2^k, 1^{q-2k})}$ for q even. Combining with this with Theorem 3.10 gives a projection

$$H^i(\Gamma_{2,s}) \longrightarrow \bigoplus_{2k < i-1, k \text{ odd}} P_{(2^k, 1^{i-2k-1})} \circ P_{(s-i+1)}$$

for i odd. This projection will be useful for constructing nice homology classes from cohomology classes, which we do in Section 4.3.

3.6. Arbitrary rank. The representation theory we used to compute the cohomology of $\Gamma_{2,s}$ gives information about the cohomology of $\Gamma_{n,s}$ for all values of n . In this section we show how this works.

Theorem 3.14. *If $s \geq n(2m+1)$ then $H^{2mn}(\Gamma_{n,s})$ contains the \mathfrak{S}_s -module $P_{(s-2mn, n^{2m})}$ as a direct summand with multiplicity 1. In particular, $H^{2mn}(\Gamma_{n,s}) \neq 0$ for all $s \geq 2mn + n$.*

Proof. The E_2 term of the spectral sequence (3.2) is $H^p(\mathrm{Out}(F_n); H^q(F_n^s))$. The $p = 0$ column is straightforward to calculate because it is simply a calculation of $\mathrm{GL}_n(\mathbb{Z})$ invariants of a well-understood module. The other columns consist of groups that are not known, so our strategy will be to look for \mathfrak{S}_s -representations in the $p = 0$ column that cannot appear in the other columns. Such a representation cannot support a nontrivial differential, as all differentials are \mathfrak{S}_s -equivariant, so survives to E_∞ and hence to $H^*(\Gamma_{n,s})$.

The action of $\mathrm{Out}(F_n)$ on $H = H^1(F_n) \cong \mathbb{k}^n$ factors through the usual action of $\mathrm{GL}_n(\mathbb{Z})$ on H , and as before, using Lemma 3.4 we have

$$\begin{aligned} H^0(\mathrm{Out}(F_n); H^q(F_n^s)) &= H^0(\mathrm{Out}(F_n); H^{\wedge q} \circ P_{(s-q)}) \\ &= H^0(\mathrm{Out}(F_n); H^{\wedge q}) \circ P_{(s-q)} \\ &= H^0(\mathrm{GL}_n(\mathbb{Z}); H^{\wedge q}) \circ P_{(s-q)}. \end{aligned}$$

By Schur–Weyl duality, $H^{\wedge q} \cong \bigoplus_{|\lambda|=q} \mathbb{S}_\lambda H \otimes P_{\lambda'}$, where $\mathbb{S}_\lambda H$ is the irreducible GL_n -representation corresponding to λ . It follows from the character formula [15, Theorem 6.3], that $\mathbb{S}_\lambda H$ is 1-dimensional if and only if q is a multiple of n , say $q = kn$ and $\lambda = (k^n)$. In this case $\mathbb{S}_{(k^n)} H$ is the 1-dimensional $\mathrm{GL}_n(\mathbb{k})$ -representation which is the k th power of the determinant (the Schur polynomial is $S_{k^n} = (x_1 x_2 \dots x_n)^k$). Thus $\mathbb{S}_\lambda H$ is a trivial $\mathrm{GL}_n(\mathbb{Z})$ -module only when $q = 2mn$ is an even multiple of n and $\lambda = ((2m)^n)$. We conclude that $H^0(\mathrm{Out}(F_n); H^q(F_n^s)) = 0$ unless $q = 2mn$ in which case we have

$$H^0(\mathrm{Out}(F_n); H^{2mn}(F_n^s)) = P_{((2m)^n)'} \circ P_{(s-2mn)} = P_{(n^{2m})} \circ P_{(s-2mn)}.$$

Using the Pieri rule to decompose this representation, we see that as long as $s - 2mn \geq n$ one of the terms we get is $P_{(s-2mn, n^{2m})}$, obtained by adding one box below each existing column and the rest to the right of the first row; this is illustrated in Figure 6.




Figure 6. Adding boxes to (n^{2m}) to obtain the Young diagram for one term of the induced module $P_{(n^{2m})} \circ P_{(s-2mn)}$

We have shown $P_{(s-2mn, n^{2m})}$ occurs in $E_2^{0, 2mn}$. We now claim it does not appear in any row below the $2mn$ -th row, so that all differentials from $P_{(s-2mn, n^{2m})}$ must vanish, and $P_{(s-2mn, n^{2m})}$ survives in $H^*(\Gamma_{n, s})$. Since

$$\begin{aligned} E_2^{p, q} &= H^p(\text{Out}(F_n), H^q(F_n^s)) \\ &= H^p(\text{Out}(F_n), H^{\wedge q} \circ P_{(s-q)}) \\ &= \bigoplus_{|\lambda|=q} H^p(\text{Out}(F_n), \mathbb{S}_\lambda H \otimes P_{\lambda'} \circ P_{(s-q)}) \\ &= \bigoplus_{|\lambda|=q} H^p(\text{Out}(F_n), \mathbb{S}_\lambda H) \otimes P_{\lambda'} \circ P_{(s-q)}, \end{aligned}$$

it suffices to show that $P_{(s-2mn, n^{2m})}$ cannot occur as a term in any of the induced modules $P_{\lambda'} \circ P_{(s-q)}$ with $|\lambda| = q < 2mn$. This is the case because the first row of any diagram appearing in $P_{\lambda'} \circ P_{(s-q)}$ has length at least $s - q > s - 2mn$. \square

Remark 3.15. The module $P_{(s-2mn, n^{2m})}$ used in the above proof is only a tiny piece of $E_2^{0, 2mn} = H^0(\text{Out}(F_n); H^q(F_n^s))$. It seems likely that a much larger part survives to infinity in the spectral sequence and thus contributes to the cohomology of $\Gamma_{n, s}$.

4. Subgroups supporting homology classes in $\Gamma_{1, s}$ and $\Gamma_{2, s}$

In later sections of the paper it will be more natural to work with homology than cohomology. The universal coefficient theorem formally allows us to do this, since we have finite-dimensional homology groups and coefficients in a field. In this section

we show that in rank 1 and 2 we can also describe some homology classes more directly as classes supported on certain easily understood subgroups.

4.1. Rank one. The situation for rank 1 is quite simple so we describe this first. By Proposition 3.7 the odd-dimensional cohomology of $\Gamma_{1,s}$ vanishes and the inclusion of \mathbb{Z}^{s-1} into $\Gamma_{1,s}$ induces an isomorphism on cohomology in even degrees, hence this holds also for homology. This implies that the top homology class of any subgroup of even rank in \mathbb{Z}^{s-1} maps to a nontrivial class in $H_*(\Gamma_{1,s})$, and $H_*(\Gamma_{1,s})$ has a basis of such classes. If $s = 2k + 1$ then the entire subgroup $\mathbb{Z}^{s-1} = \mathbb{Z}^{2k}$ has even rank and its top homology class maps to a nontrivial class $\alpha_k \in H_{2k}(\Gamma_{1,2k+1})$, which is well defined up to sign. The class α_k will be used to construct the Morita class μ_k in Section 5.1.

4.2. Rank two, even homology degree. Now we turn to rank 2, where $H_*(\Gamma_{2,s})$ is considerably more complicated. This extra complication is relatively mild in even degrees, so we examine those first. It suffices to consider $H_{4k}(\Gamma_{2,s})$ since $H_{4k+2}(\Gamma_{2,s}) = 0$ by Theorem 3.10.

Notation. Throughout this section and the next we fix generators x and y for F_2 and we let \mathbf{x} and \mathbf{y} denote their images in $H_1(F_2)$, with \mathbf{x}^* and \mathbf{y}^* the dual basis of $H^1(F_2)$. We also set $\hat{H} = H_1(F_2)$; the notation is meant to distinguish it from $H = H^1(F_2)$.

For disjoint subsets I and J of $\{1, 2, \dots, s\}$ let $A_{I,J}$ be the abelian subgroup of F_2^s consisting of s -tuples with powers of x in the I coordinates, powers of y in the J coordinates, and the identity in the other coordinates. We have inclusions $A_{I,J} \subset F_2^s \subset \Gamma_{2,s}$, and we let $\alpha_{I,J} \in H_*(\Gamma_{2,s})$ be the image of a generator of the top-dimensional homology of $A_{I,J}$.

Proposition 4.1. *If $|I| = |J| = 2k$ for some k then the class $\alpha_{I,J} \in H_{4k}(\Gamma_{2,s})$ is nonzero and these classes $\alpha_{I,J}$ generate $H_{4k}(\Gamma_{2,s})$.*

Proof. First we show that $\alpha_{I,J}$ is nonzero when $|I| = |J| = 2k$ for some $k \geq 1$. (Here $s \geq 4k$ since $A_{I,J} \subset F_2^s$.) We do this by finding a cohomology class in $H^{4k}(\Gamma_{2,s})$ that pairs nontrivially with $\alpha_{I,J}$.

By Lemma 3.8 we have $H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge 4k-1}) = 0$. Therefore

$$H^1(\mathrm{GL}_2(\mathbb{Z}); H^{4k-1}(F_2^s)) = H^1(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge 4k-1}) \circ P_{(s-4k)} = 0,$$

and

$$H^{4k}(\Gamma_{2,s}) \cong H^0(\mathrm{GL}_2(\mathbb{Z}); H^{4k}(F_2^s)) \cong H^0(\mathrm{GL}_2(\mathbb{Z}); H^{\wedge 4k}) \circ P_{(s-4k)}.$$

Thus to compute $H^{4k}(\Gamma_{2,s})$ as an \mathfrak{S}_s -module it suffices to understand the invariants of the diagonal action of $\mathrm{GL}_2(\mathbb{Z})$ on $H^{\wedge 4k}$. As a $\mathrm{GL}_2(\mathbb{Z})$ -module, $H^{\wedge 4k}$ is the same as $H^{\otimes 4k}$, and we describe the (classical) answer below.

A straightforward calculation shows that the diagonal action of an element $T \in \mathrm{GL}_2(\mathbb{Z})$ on $H \otimes H$ sends $\omega^* = \mathbf{x}^* \otimes \mathbf{y}^* - \mathbf{y}^* \otimes \mathbf{x}^*$ to $(\det T) \cdot \omega^*$. The diagonal action on $H^{\otimes 4k}$ sends $(\omega^*)^{\otimes 2k}$ to $(\det T)^{2k} (\omega^*)^{\otimes 2k}$, so since $\det T = \pm 1$ and $2k$ is even this is an invariant. Any permutation of the indices $\{1, \dots, 4k\}$ produces another invariant, and the invariants defined in this way span the entire space of invariants (see, e.g., [15] for details). Note that each term in each of these invariants has an equal number of \mathbf{x}^* 's and \mathbf{y}^* 's, so this is true of any invariant.

Suppose first that $s = 4k$ and let $I = \{1, 3, \dots, 4k - 1\}$, the odd indices, and $J = \{2, 4, \dots, 4k\}$, the even indices. Then the image of $H_{4k}(A_{I,J}) \cong H_1(\mathbb{Z}) \otimes \dots \otimes H_1(\mathbb{Z})$ in $H_1(F_2^{4k}) \cong \hat{H} \otimes \dots \otimes \hat{H}$ is generated by $\mathbf{z} = \mathbf{x} \otimes \mathbf{y} \otimes \dots \otimes \mathbf{x} \otimes \mathbf{y}$. Since this matches the first term of $(\omega^*)^{\otimes 2k}$, the cohomology class $(\omega^*)^{\otimes 2k} \in H^{4k}(\Gamma_{2,4k})$ pairs nontrivially with $\alpha_{I,J}$, which is the image of \mathbf{z} in $H_{4k}(\Gamma_{2,4k})$. This shows that $\alpha_{I,J}$ is nonzero. Permuting the indices produces other nonzero classes $\alpha_{I,J}$ that span $H_{4k}(\Gamma_{2,4k})$ since the corresponding cohomology classes span $H^{4k}(\Gamma_{2,4k})$.

If $s > 4k$, any of the natural inclusions $\Gamma_{2,4k} \rightarrow \Gamma_{2,s}$ (given by gluing extra leaves to the leaf vertices of $X_{2,4k}$ and extending maps by the identity) induces an injection $H_{4k}(\Gamma_{2,4k}) \rightarrow H_{4k}(\Gamma_{2,s})$ mapping each $\alpha_{I,J}$ nontrivially. On homology, the image of this map depends only on the inclusion $\{1, \dots, 2k\} \rightarrow \{1, \dots, s\}$ of leaf vertices. Since $H_{4k}(\Gamma_{2,s}) = H_{4k}(\Gamma_{2,4k}) \circ P_{(s-4k)}$, these classes span all of $H_{4k}(\Gamma_{2,s})$. \square

Remark 4.2. If I and J are disjoint subsets of $\{1, \dots, s\}$ of different size, then $\alpha_{I,J}$ is trivial because the top-dimensional homology class of $A_{I,J}$ is a simple tensor with an unequal number of \mathbf{x} 's and \mathbf{y} 's, so every invariant evaluates trivially on it.

Remark 4.3. The classes $\alpha_{I,J} \in H_{4k}(\Gamma_{2,s})$ are not linearly independent. There are several possible ways to obtain a subset of these classes which form a basis of $H_{4k}(\Gamma_{2,s})$. Since the dimension of $H_{4k}(\Gamma_{2,s})$ is closely related to the dimension of $P_{(2k,2k)}$, which is equal to the Catalan number C_{2k} , one can use combinatorial objects such as non-crossing partitions or Young tableaux to describe such a basis. Here is one possible description of a basis.

Claim. *The space $H_{4k}(\Gamma_{2,s})$ has a basis consisting of those $\alpha_{I,J}$ for which $I = \{i_1 < i_2 < \dots < i_{2k}\}$ and $J = \{j_1 < j_2 < \dots < j_{2k}\}$ are disjoint subsets of $\{1, \dots, s\}$ such that $i_t < j_t$ for each $t = 1, \dots, 2k$.*

The proof of this involves a deeper use of representation theory so we will not give it here.

4.3. Rank two, odd homology degree. Constructing classes of odd homology degree is more difficult since no subgroups of F_2^s support such classes. As a result we must use slightly more complicated subgroups of $\Gamma_{2,s}$. We use the same notation as in the previous section for generators of F_2 and its homology and cohomology.

Fix disjoint subsets $I, J \subset \{1, \dots, s\}$. Let $B_{I,J} \cong F_2^{|I|} \times \mathbb{Z}^{|J|}$ be the subgroup of F_2^s consisting of s -tuples with arbitrary elements in the I coordinates, powers

of x in the J -coordinates and the identity in coordinates not indexed by I or J . Recall that $\Gamma_{2,s}$ maps onto $\Gamma_{2,0} \cong \mathrm{GL}(2, \mathbb{Z})$ with kernel F_2^s . Let $P \cong \mathbb{Z}$ denote the unipotent subgroup of $\mathrm{GL}_2(\mathbb{Z})$ generated by $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, corresponding to the (outer) automorphism of F_2 fixing x and sending y to xy . Lift the generator of P to an element $\varphi \in \Gamma_{2,s}$ that wraps the y -loop of $X_{2,s}$ around both itself and the x -loop, and fixes the x -loop and all leaves. This normalizes $B_{I,J}$, and we define $M_{I,J}$ to be the subgroup of $\Gamma_{2,s}$ generated by φ and $B_{I,J}$. We now have a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & F_2^s & \longrightarrow & \Gamma_{2,s} & \longrightarrow & \mathrm{GL}_2(\mathbb{Z}) \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & B_{I,J} & \longrightarrow & M_{I,J} & \longrightarrow & P \longrightarrow 1. \end{array}$$

Note that $M_{I,J}$ splits as the product $M_I \times \mathbb{Z}^{|J|}$ where $M_I = M_{I,\emptyset}$ and $\mathbb{Z}^{|J|} = B_{\emptyset,J}$.

We will be interested in the cases when $|I|$ and $|J|$ are even, and we let $|I| + |J| = 2k$. The top-dimensional homology of $B_{I,J}$ is $H_{2k}(B_{I,J}) \cong \hat{H}^{\otimes |I|} \otimes \hat{X}^{\otimes |J|}$, where $\hat{X} \cong \mathbb{k}$ is the subspace of $\hat{H} = H_1(F_2)$ spanned by \mathbf{x} . From the Leray–Serre spectral sequence it follows that $H_*(M_{I,J})$ vanishes above dimension $2k + 1$ and $H_{2k+1}(M_{I,J}) = H_1(P; H_{2k}(B_{I,J}))$.

4.3.1. The case $|I|=2$. The analysis of the case $|I| = 2$ is easier than the general case and will suffice for our construction of the Eisenstein classes in Section 5.6, so we begin with this case.

We first compute $H_{2k+1}(M_{I,J})$. From the splitting $M_{I,J} = M_I \times \mathbb{Z}^{|J|}$ we have

$$H_{2k+1}(M_{I,J}) \cong H_3(M_I) \otimes H_{2k-2}(\mathbb{Z}^{2k-2}) \cong H_3(M_I) \cong H_1(P; H_2(B_{I,\emptyset})).$$

Since $P \cong \mathbb{Z}$ the first homology $H_1(P; H_2(B_{I,\emptyset}))$ is just the invariants of the action of P on $H_2(B_{I,\emptyset}) = \hat{H} \otimes \hat{H}$. This is the diagonal action, where P acts on \hat{H} by sending $\mathbf{x} \rightarrow \mathbf{x}$ and $\mathbf{y} \rightarrow \mathbf{x} + \mathbf{y}$. It is easy to compute that the space of invariants is 2-dimensional, spanned by $\mathbf{x} \otimes \mathbf{x}$ and $\omega = \mathbf{x} \otimes \mathbf{y} - \mathbf{y} \otimes \mathbf{x}$. Thus

$$H_{2k+1}(M_{I,J}) \cong H_1(P; H_{2k}(B_{I,J})) \cong [H_{2k}(B_{I,J})]^P \cong [\hat{H}^{\otimes 2} \otimes \hat{X}^{\otimes 2k-2}]^P \cong \mathbb{k}^2$$

with basis \mathbf{x}^{2k} and $\omega \otimes \mathbf{x}^{2k-2}$.

The class in $H_{2k+1}(M_{I,J})$ corresponding to $\omega \otimes \mathbf{x}^{2k-2}$ is the one whose image $m_{I,J}$ in $H_{2k+1}(\Gamma_{2,s})$ will be used as a building block for Eisenstein classes. In Section 7.2 we give a different, more geometric construction of this class as the image of the fundamental class of a manifold mapped into a moduli space of graphs.

The natural actions of $\mathfrak{S}_I = \mathfrak{S}_2$ and $\mathfrak{S}_J = \mathfrak{S}_{2k-2}$ on $H_{2k+1}(M_{I,J})$ are easy to describe since these two symmetric groups act separately on the factors of the splitting $M_{I,J} = M_I \times \mathbb{Z}^{|J|}$. For \mathfrak{S}_I the transposition σ interchanges the F_2 factors of F_2^2 , so $\sigma(\mathbf{x} \otimes \mathbf{x}) = -\mathbf{x} \otimes \mathbf{x}$ since the cross product in the Künneth formula is

anti-symmetric. For the class $\omega = \mathbf{x} \otimes \mathbf{y} - \mathbf{y} \otimes \mathbf{x}$ we have $\sigma(\omega) = \omega$ since we get one minus sign from the minus sign in ω and another from anti-symmetry in the Künneth formula. For an element $\sigma \in \mathfrak{S}_J$ the action on \mathbf{x}^{2k-2} is just by the sign of σ .

Because of the anti-symmetric action of \mathfrak{S}_J , the classes $m_{I,J} \in H_{2k+1}(\Gamma_{2,s})$ are well defined only up to sign. We now show they are nontrivial and describe how much of $H_{2k+1}(\Gamma_{2,s})$ they account for.

Proposition 4.4. *For $I, J \subset \{1, \dots, s\}$ with $|I| = 2$ and $|J| = 2k - 2 \geq 2$ the map $H_{2k+1}(M_{I,J}) \rightarrow H_{2k+1}(\Gamma_{2,s})$ induced by inclusion has 1-dimensional image spanned by $m_{I,J}$, and the \mathfrak{S}_s -module generated by $m_{I,J}$ is isomorphic to $P_{(2,1^{2k-2})} \circ P_{(s-2k)}$.*

In particular, when $s = 2k$ the classes $m_{I,J}$ generate $H_{2k+1}(\Gamma_{2,2k})$ only when $k \leq 3$; this follows from Theorem 3.10.

Proof. To prove that $m_{I,J}$ is nonzero we find a cohomology class that pairs nontrivially with it.

Assume first that $s = 2k$. To simplify notation we also assume $I = \{1, 2\}$ and $J = \{3, \dots, 2k\}$ and set $B = B_{I,J}$, $M = M_{I,J}$ and $m = m_{I,J}$. The map from $H^{2k+1}(\Gamma_{2,2k}) \cong H^1(\mathrm{GL}_2(\mathbb{Z}); H^{2k}(F_2^{2k}))$ to $H^{2k+1}(M)$ induced by the inclusion $M \hookrightarrow \Gamma_{2,2k}$ factors as

$$H^1(\mathrm{GL}_2(\mathbb{Z}); H^{2k}(F_2^{2k})) \longrightarrow H^1(P; H^{2k}(F_2^{2k})) \longrightarrow H^1(P; H^{2k}(B))$$

where the first map is induced by the inclusion $P \hookrightarrow \mathrm{GL}_2(\mathbb{Z})$ and the second by the map of coefficients induced by $B \hookrightarrow F_2^{2k}$.

In Remark 3.13 we pointed out that the first map is a surjection onto the odd terms of the decomposition of $H^1(P; H^{2k}(F_2^{2k}))$ into irreducible \mathfrak{S}_{2k} -modules. Here is a more explicit description of this map. Since $P \cong \mathbb{Z}$, for any P -module V we have $H^1(P; V) \cong V_P$. If V is a vector space there is a canonical isomorphism $(V^*)_P \cong (V^P)^*$ (sending f to its restriction to V^P). In particular, using the universal coefficient theorem we get natural isomorphisms

$$\begin{aligned} H^1(P; H^{2k}(F_2^{2k})) &= [H^{2k}(F_2^{2k})]_P \cong ([H_{2k}(F_2^{2k})]^P)^* \\ &= ([H_1(F_2)^{\otimes 2k} \otimes \mathrm{alt}]^P)^* \\ &= ([\hat{H}^{\otimes 2k}]^P)^* \otimes \mathrm{alt}, \end{aligned}$$

where alt refers to the \mathfrak{S}_{2k} -action. Now recall that the space of P -invariants in $\hat{H} \otimes \hat{H}$ is spanned by ω and \mathbf{x}^2 . Since $\hat{H} \otimes \hat{H} = \wedge^2 \hat{H} \oplus \mathrm{Sym}^2 \hat{H}$, this shows that the subspace of P -invariants in each summand is 1-dimensional. This is a special instance of the general fact that the space of P -invariants in $(\wedge^2 \hat{H})^{\otimes \ell} \otimes \mathrm{Sym}^{2k-2\ell} \hat{H}$ is 1-dimensional, spanned by $\omega^\ell \otimes \mathbf{x}^{2k-2\ell}$. The Schur–Weyl decomposition of $\hat{H}^{\otimes 2k}$ then shows that the image of the first map can be identified with $\bigoplus_{\ell < k \text{ odd}} P_{(2^\ell, 1^{2k-2\ell})}$, as in Remark 3.13, where each term is generated by $\omega^\ell \otimes \mathbf{x}^{2k-2\ell}$ as an \mathfrak{S}_{2k} -module.

For the second map, note that

$$\begin{aligned} H^1(P; H^{2k}(B)) &= [H \otimes H \otimes X^{\otimes 2k-2}]_P \otimes \text{alt} \\ &\cong ([\hat{H} \otimes \hat{H} \otimes \hat{X}^{\otimes 2k-2}]^P)^* \otimes \text{alt}, \end{aligned}$$

where this alt refers to the action of $\mathfrak{S}_2 \times \mathfrak{S}_{2k-2}$ which permutes the factors of $B = F_2^2 \times \mathbb{Z}^{2k-2}$ independently. Thus the map

$$H^1(P; H^{2k}(F_2^{2k})) \rightarrow H^1(P; H^{2k}(B))$$

becomes

$$([H_{2k}(F_2^{2k})]^P)^* \rightarrow ([H_{2k}(B)]^P)^*$$

i.e.,

$$([\hat{H}^{\otimes 2k}]^P)^* \otimes \text{alt} \rightarrow ([\hat{H}^{\otimes 2} \otimes \hat{X}^{\otimes 2}]^P)^* \otimes \text{alt}.$$

The map on the first factor is just the transpose of the inclusion map $[\hat{H}^{\otimes 2} \otimes \hat{X}^{\otimes 2}]^P \hookrightarrow [\hat{H}^{\otimes 2k}]^P$ and in particular sends $(\omega \otimes \mathbf{x}^{2k-2})^*$ to itself.

Since $(\omega \otimes \mathbf{x}^{2k-2})^*$ is in the image of the first map $H^1(\text{GL}_2(\mathbb{Z}); H^{2k}(F_2^{2k})) \rightarrow H^1(P; H^{2k}(F_2^{2k}))$, there is a cohomology class in $H^{2k+1}(\Gamma_{2,2k})$ which hits it under the composition

$$\begin{aligned} H^{2k+1}(\Gamma_{2,2k}) &= H^1(\text{GL}_2(\mathbb{Z}); H^{2k}(F_2^{2k})) \rightarrow H^1(P; H^{2k}(F_2^{2k})) \\ &\rightarrow H^1(P; H^{2k}(B)) = H^{2k+1}(M). \end{aligned}$$

This class evaluates nontrivially on the image $m \in H_{2k+1}(\Gamma_{2,2k})$ of $\omega \otimes \mathbf{x}^{2k-2} \in H_{2k+1}(M)$, showing that m is nontrivial.

Any permutation of the indices $\{1, \dots, 2k\}$ gives another class in $H_{2k+1}(\Gamma_{2,2k})$. The \mathfrak{S}_{2k} -submodule generated by m is isomorphic to $P_{(2,1^{2k-2})}$, which coincides with $H_{2k+1}(\Gamma_{2,2k})$ only when $k = 2, 3$. This completes the proof of the proposition for $s = 2k$.

The generalization to $s > 2k$ is straightforward, since $H^{2k+1}(\Gamma_{2,s}) \cong H^1(\text{GL}_2(\mathbb{Z}); H^{2k}(F_2^s)) \cong H^1(\text{GL}_2(\mathbb{Z}); H^{\wedge 2k}) \circ P_{(s-4k)}$. The \mathfrak{S}_s -module generated by the image of $H_{2k+1}(M_{I,J})$ in $H_*(\Gamma_{2,s})$ is isomorphic to $P_{(2,1^{2k-2})} \circ P_{(s-2k)}$. \square

As was noted when the classes $m_{I,J}$ were defined, they are invariant under transposing the two indices in I and anti-invariant under permutations of the indices in J . When $s = 2k$ we can obtain a class which is anti-invariant under a larger group of permutations by adding together signed images of $m_{I,J}$ under appropriate permutations. Specifically, let $I = \{1, 2\}$, $J = \{3, \dots, 2k\}$ and $m = m_{I,J}$ as in the proof of Proposition 4.4 and choose an index $i \in \{1, \dots, 2k\}$. Then define

$$m_i = \sum_{\sigma(1)=i} \text{sign}(\sigma) \sigma(m),$$

where the sum is over all permutations $\sigma \in \mathfrak{S}_{2k}$ which send 1 to i . The class m_i is then anti-invariant under $\text{stab}_{\mathfrak{S}_{2k}}(i)$. For example when $I = \{1, 2\}$ and $J = \{3, 4\}$, so m corresponds to $\mathbf{xyxx} - \mathbf{yxxx}$ (omitting tensor symbols for simplicity), the class m_1 corresponds to $6\mathbf{yxxx} - 2\mathbf{xyxx} - 2\mathbf{xxxy} - 2\mathbf{xxxxy}$, up to sign. The formula for m_2 is similar, and one sees that $m = \pm \frac{1}{8}(m_1 - m_2)$. For larger J there are analogous formulas.

4.3.2. The general case. Now we consider the general case $|I| = 2\ell$ for odd $\ell \geq 1$. This is more involved because for $\ell > 1$ the top-dimensional cohomology $H^{2k+1}(M_{I,J})$ is quite large and it is not immediately clear how to pick a distinguished element dual to $\omega^{\otimes \ell} \otimes \mathbf{x}^{2k-2\ell}$. We settle this by using the unique element which is invariant under the action of certain involutions. This is motivated by the case $\ell = 1$, where the element $\omega \in \hat{H}^2$ spans the invariants of \hat{H}^2 under the action of the involution (12).

Given any set T of disjoint transpositions, let N_T denote the elementary abelian subgroup that they generate.

Proposition 4.5. *Let T be a set of ℓ disjoint transpositions of the set I . For ℓ odd, the top homology $H_{2k+1}(M_{I,J})$ contains a unique (up to scalar multiple) element which is invariant under the action of $N_T \subset \mathfrak{S}_I$. The image of this element under the map $H_{2k+1}(M_{I,J}) \rightarrow H_{2k+1}(\Gamma_{2,s})$ induced by inclusion is nonzero. The \mathfrak{S}_s -module generated by this image is isomorphic to $P_{(2^\ell, 1^{2k-2\ell})} \circ P_{(s-2k)}$.*

Proof. In order to simplify the notation we will assume that $I = \{1, \dots, 2\ell\}$, $J = \{2\ell + 1, \dots, 2k\}$ and $T = \{(1, 2), (3, 4), \dots, (2\ell - 1, 2\ell)\}$, and set $B_\ell = B_{I,J}$ and $M_\ell = M_{I,J}$.

Recall that \hat{X} is the 1-dimensional subspace of $H_1(F_2)$ spanned by \mathbf{x} . The actions of P and N_T on $H_{2k}(B_\ell) = \hat{H}^{2\ell} \otimes \hat{X}^{2k-2\ell}$ commute so

$$\begin{aligned} [H_{2k+1}(M_\ell)]^{N_T} &= [H_1(P; H_{2k}(B_\ell))]^{N_T} \\ &= [[H_{2k}(B_\ell)]^P]^{N_T} \\ &= [\hat{H}^{2\ell} \otimes \hat{X}^{2k-2\ell}]^P]^{N_T} \\ &= [\hat{H}^{2\ell} \otimes \hat{X}^{2k-2\ell}]^{N_T}]^P \\ &= [\hat{H}^{2\ell}]^{N_T} \otimes \hat{X}^{2k-2\ell}]^P \end{aligned}$$

The space of invariants in $\hat{H}^{2\ell}$ under the action of N_T is 1-dimensional, spanned by $\omega^{\otimes \ell}$, so the entire space $[H_{2k+1}(M_\ell)]^{N_T}$ is at most 1-dimensional. It is exactly 1-dimensional since the element

$$m_\ell = \omega^{\otimes \ell} \otimes \mathbf{x}^{2k-2\ell} \in [\hat{H}^{2\ell} \otimes \hat{X}^{2k-2\ell}]^P$$

is invariant under the action of N_T . We will show that m_ℓ has nontrivial image in $H_{2k+1}(\Gamma_{2,s})$ if ℓ is odd.

Let D_ℓ be the subspace of $H^{2k}(F_2^{2k}) = H^{\wedge 2k}$ generated as a $GL_2(\mathbb{Z})$ -module by $(\omega^*)^{\otimes \ell} \otimes (\mathbf{x}^*)^{\otimes 2k-2\ell}$. Thus

$$D_\ell \cong \mathbb{S}_{(2k-\ell, \ell)}(H) \cong \det^\ell \otimes \text{Sym}^{2k-2\ell}(H).$$

(See Lemma 3.3.) Viewing D_ℓ as a submodule of the induced module $H^{2k}(F_2^s) = H^{2k}(F_2^{2k}) \circ P_{(s-2k)}$ we see that m_ℓ pairs nontrivially with the P -coinvariants in D_ℓ (to compute these coinvariants, note that the action of P on $H = H^1(F_2)$ is dual to its action on $H_1(F_2)$ so sends $\mathbf{x}^* \mapsto \mathbf{x}^* + \mathbf{y}^*$ and fixes \mathbf{y}^* .) Therefore m_ℓ pairs nontrivially with the cohomology class generating $H^1(P; D_\ell)$.

Since ℓ is odd Remark 3.13 shows that the map $H^1(GL_2(\mathbb{Z}); D_\ell) \rightarrow H^1(P; D_\ell)$ is surjective. Hence the class m_ℓ pairs nontrivially with a class in $H^1(GL_2(\mathbb{Z}); D_\ell)$ which is the image of a class in $H^1(GL_2(\mathbb{Z}); H^{2k}(F_2^s)) = H^{2k+1}(\Gamma_{2,s})$. This shows that the homology class m_ℓ is nonzero in $H_*(\Gamma_{2,s})$.

The last statement of the proposition follows from the Schur–Weyl decomposition of $H_{2k}(B_\ell)$:

$$H_{2k}(B_\ell) = \hat{H}^{\wedge 2\ell} \otimes \hat{X}^{\wedge 2k-2\ell} = \bigoplus_{i \leq \ell} \mathbb{S}_{(\ell+i, \ell-i)} \hat{H} \otimes P_{(2^\ell-i, 1^{2i})} \otimes \hat{X}^{\wedge 2k-2\ell}$$

The element $m_\ell = \omega^\ell \otimes \mathbf{x}^{2k-2\ell}$ is in the $i = 0$ term $\mathbb{S}_{(\ell, \ell)} \hat{H} \cong \det^\ell \otimes \text{Sym}^{2k-2\ell} \hat{H}$, so the \mathfrak{S}_{2k} -submodule of $H_{2k}(B_\ell)$ it generates is of type $P_{(2^\ell, 1^{2k-2\ell})}$, which is then induced up to $P_{(2^\ell, 1^{2k-2\ell})} \circ P_{(s-2k)}$. \square

Remark 4.6. The \mathfrak{S}_s -module generated by the element of $H^{2k+1}(\Gamma_{2,s})$ found in Proposition 4.5 must come from the term $\mathcal{M}_{2k+2-2\ell} \otimes P_{(2^\ell, 1^{2k-2-2\ell})} \circ P_{(s-2k)}$ of the computation of $H^{2k+1}(\Gamma_{2,s})$ in Theorem 3.10. In fact it comes from the map $\mathcal{M}_{2k+2-2\ell} \rightarrow \mathbb{k}$ obtained by evaluating the modular form at infinity. This is clear from the construction since we are using the parabolic subgroup P and the inclusion of P into $GL_2(\mathbb{Z})$ kills all other classes (see Section 3.5.1).

Remark 4.7. Recall that after the proof of Proposition 4.4 we defined classes m_i using a symmetrization procedure. Similarly, we can use extra symmetrization to obtain classes $m_{I', J'}$ indexed by disjoint sets I' and J' with $|I'| = \ell$ and $|J'| = 2k - \ell$ for ℓ odd which span the module $P_{(2^\ell, 1^{2k-2\ell})} \circ P_{(s-2k)}$ inside $H_{2k+1}(\Gamma_{2,s})$. These elements are anti-invariant under the action of $\mathfrak{S}_{I'} \times \mathfrak{S}_{J'}$ and invariant under permutations fixing I' and J' pointwise. These elements generate all of $H_{2k+1}(\Gamma_{2,s})$ if $k \leq 5$. For $k > 5$ they generate only the homology coming from the parabolic group $P \subset SL_2(\mathbb{Z})$. The elements $m_{I', J'}$ are not linearly independent, but a subset similar to the one described in Remark 4.3 can be used to form a basis of the corresponding \mathfrak{S}_s -module.

5. Gluing classes together

Given a graph $X_{n,s}$ we can obtain a new set of graphs $\{X_{n_i,s_i}\}$ by snipping some of the edges at their midpoints; the snipped edges will become leaves in the X_{n_i,s_i} . Conversely, suppose we have a set of graphs $\{X_{n_1,s_1}, \dots, X_{n_k,s_k}\}$ and a *gluing pattern* ϕ which pairs up some or all of the leaf vertices to form a connected graph X_ϕ . An example is shown in Figure 7. If X_ϕ has rank n and s leaves, then the gluing

$$X_{n_1,s_1} \cup \dots \cup X_{n_k,s_k} \longrightarrow X_\phi$$

induces a homomorphism

$$p_\phi: \Gamma_{n_1,s_1} \times \dots \times \Gamma_{n_k,s_k} \longrightarrow \Gamma_{n,s}.$$

This in turn induces an *assembly map* on homology via the cross product,

$$A_\phi: H_*(\Gamma_{n_1,s_1}) \otimes \dots \otimes H_*(\Gamma_{n_k,s_k}) \longrightarrow H_*(\Gamma_{n,s}).$$

In particular, if we glue all of the univalent vertices in pairs, we obtain a map to the homology of $\text{Out}(F_n)$, and if we glue all but one we obtain a map to the homology of $\text{Aut}(F_n)$. We allow leaf vertices of a single $X_{n,s}$ to be glued together. For example, gluing all four leaf vertices of $X_{2,4}$ in pairs gives an assembly map $H_*(\Gamma_{2,4}) \rightarrow H_*(\Gamma_{4,0})$ that we use in Section 5.2.



Figure 7. Making $X_{4,3}$ from $X_{1,3}$ and $X_{2,4}$ using a gluing map ϕ

Remark 5.1. Assembly maps are associative since this is obviously true for gluing graphs together, and the cross product in homology is associative. In particular, if a gluing is done in two stages, the assembly map factors through the intermediate stage.

Remark 5.2. The vcd of $\Gamma_{n,s}$ is $2n + s - 3$ if $n > 0$, and for a k -fold assembly map as above with each $n_i > 0$ this is given by the formula

$$\text{vcd}(\Gamma_{n,s}) = \text{vcd}(\Gamma_{n_1,s_1}) + \dots + \text{vcd}(\Gamma_{n_k,s_k}) + (k - 1).$$

To see this it suffices by induction to consider the case of gluing a single pair of leaves. If $k = 1$ we are gluing two leaves of the same graph together, increasing n by one and decreasing s by two, so the vcd is unchanged. If $k = 2$ and we glue a leaf of one graph to a leaf of the other we have $n = n_1 + n_2$ and $s = s_1 + s_2 - 2$, so $2n + s - 3$ is one more than the sum $(2n_1 + s_1 - 3) + (2n_2 + s_2 - 3)$. A consequence of this relation between the vcd's is that a k -fold assembly map with $k > 1$ cannot produce homology classes in the vcd of $\Gamma_{n,s}$. This holds even when some n_i 's are 0, provided we exclude trivial factors with $(n_i, s_i) = (0, 2)$.

A different gluing ϕ' may also produce a graph $X_{\phi'}$ of rank n with s leaves. As noted at the beginning of Section 2.1, a bijection between the leaf vertices of X_ϕ and $X_{\phi'}$ determines an isomorphism between the targets of both assembly maps. For $\phi' = \phi$ this gives the action of the symmetric group \mathfrak{S}_s on $H_*(\Gamma_{n,s})$.

The left-hand side of the assembly map A_ϕ is a priori an $\mathfrak{S}_{s_1} \times \cdots \times \mathfrak{S}_{s_k}$ -module. The gluing ϕ interacts with the action of $\mathfrak{S}_{s_1} \times \cdots \times \mathfrak{S}_{s_k}$ in various ways, which can be explained by the following two observations:

- (1) If t_{ij} leaves of X_{n_i, s_i} are paired with leaves of X_{n_j, s_j} , then a permutation that does the same thing to both sets of leaves does not change the result of the gluing.
- (2) If there are u_i unglued leaves in X_{n_i, s_i} , then permuting them can be done before or after gluing with the same effect.

The algebraic effect of the first observation is that the map A_ϕ factors through the coinvariants of the diagonal action of $\mathfrak{S}_{t_{ij}}$ on $H_*(\Gamma_{n_i, s_i}) \otimes H_*(\Gamma_{n_j, s_j})$. Here $\mathfrak{S}_{t_{ij}}$ acts on $H_*(\Gamma_{n_i, s_i})$ and $H_*(\Gamma_{n_j, s_j})$ by restriction of the \mathfrak{S}_{s_i} and \mathfrak{S}_{s_j} actions. For example, if the leaves of a graph with exactly s leaves are glued to the leaves of another graph with exactly s leaves, then the assembly map factors through the space of \mathfrak{S}_s coinvariants:

$$\begin{array}{ccc} H_p(\Gamma_{n_1, s}) \otimes H_q(\Gamma_{n_2, s}) & \xrightarrow{A_\phi} & H_{p+q}(\Gamma_{n_1+n_2+s-1, 0}) \\ & \searrow & \nearrow \\ & (H_p(\Gamma_{n_1, s}) \otimes H_q(\Gamma_{n_2, s}))_{\mathfrak{S}_s} & \end{array}$$

Now $(P_\lambda \otimes P_\mu)_{\mathfrak{S}_s}$ is zero unless $\lambda = \mu$, in which case it is 1-dimensional. Therefore the assembly map is trivial unless some irreducible P_λ appears in the \mathfrak{S}_s decompositions of both $H_p(\Gamma_{n_1, s})$ and $H_q(\Gamma_{n_2, s})$.

The second observation says that we can make A_ϕ into an $(\mathfrak{S}_{u_1} \times \cdots \times \mathfrak{S}_{u_k})$ -module map by realizing $\mathfrak{S}_{u_1} \times \cdots \times \mathfrak{S}_{u_k}$ as the appropriate subgroup of \mathfrak{S}_s . (Here again \mathfrak{S}_{u_i} acts on $H_*(\Gamma_{n_i, s_i})$ by restricting the \mathfrak{S}_{s_i} action.) We can sometimes obtain new information about A_ϕ by extending it to an \mathfrak{S}_s -module map, which we call \hat{A}_ϕ .

Thus the range of \hat{A}_ϕ is still $H_*(\Gamma_{n,s})$ but the domain of \hat{A}_ϕ is the module obtained by inducing $H_*(\Gamma_{n_1,s_1}) \otimes \cdots \otimes H_*(\Gamma_{n_k,s_k})$, considered as a $(\mathfrak{S}_{u_1} \times \cdots \times \mathfrak{S}_{u_k})$ -module, up to \mathfrak{S}_s . The advantage here is that \mathfrak{S}_s -modules and \mathfrak{S}_s -module maps between them are very well understood.

In the following sections we give examples of assembly maps. In particular, we show how all but one of the known nontrivial homology classes for $\text{Out}(F_n)$ and $\text{Aut}(F_n)$ are obtained by assembling classes from the homology of $\Gamma_{1,s}$ and $\Gamma_{2,s}$.

5.1. Morita's original series [33]. Recall from Section 4.1 that $H_{2k}(\Gamma_{1,2k+1}) \cong \mathbb{k}$ with generator α_k . Fix a gluing pattern $\phi : X_{1,2k+1} \cup X_{1,2k+1} \rightarrow X_{2k+2,0}$ which matches all of the leaves of the first graph with those of the second. This gives an assembly map

$$A_\phi : H_{2k}(\Gamma_{1,2k+1}) \otimes H_{2k}(\Gamma_{1,2k+1}) \longrightarrow H_{4k}(\text{Out}(F_{2k+2})).$$

The k th Morita class μ_k is the image under A_ϕ of $\alpha_k \otimes \alpha_k$. Remark 9.5 explains why this viewpoint leads to the same classes as those originally defined by Morita. The classes μ_1, μ_2 , and μ_3 are known to be nontrivial.

A lift of μ_k to $H_{4k}(\text{Aut}(F_{2k+2}))$ can be obtained via assembly maps using the gluing pattern $\phi : X_{1,2k+1} \cup X_{0,3} \cup X_{1,2k+1} \rightarrow X_{2k+2,1}$ which matches one leaf of $X_{0,3}$ with a leaf of one $X_{1,2k+1}$, another leaf of $X_{0,3}$ with a leaf of the other $X_{1,2k+1}$, and then pairs the remaining leaves of the two copies of $X_{1,2k+1}$ as before. Let $\hat{\mu}_k$ be the image of $\alpha_k \otimes \iota \otimes \alpha_k$ under the resulting assembly map $H_{2k}(\Gamma_{1,2k+1}) \otimes H_0(\Gamma_{0,3}) \otimes H_{2k}(\Gamma_{1,2k+1}) \rightarrow H_{4k}(\Gamma_{2k+2,1})$, where ι is a generator of $H_0(\Gamma_{0,3})$. The projection map $H_{4k}(\Gamma_{2k+2,1}) \rightarrow H_{4k}(\Gamma_{2k+2,0})$ then sends $\hat{\mu}_k$ to μ_k .

Remark 5.3. This argument shows more generally that every assembly map with target $H_i(\text{Out}(F_n))$ lifts to $H_i(\text{Aut}(F_n))$.

Proposition 5.4. *The Morita class $\mu_k \in H_{4k}(\text{Out}(F_{2k+2}))$ is supported in an abelian subgroup \mathbb{Z}^{4k} of $\text{Out}(F_{2k+2})$, and the analogous statement also holds for a lift to $\text{Aut}(F_{2k+2})$.*

Proof. As noted in Section 4.1, the class α_k is the top-dimensional homology class of a subgroup \mathbb{Z}^{2k} in $\Gamma_{1,2k+1}$. The assembly that produces μ_k then gives a map $\mathbb{Z}^{4k} \rightarrow \text{Out}(F_{2k+2})$ taking a generator of $H_{4k}(\mathbb{Z}^{4k})$ to μ_k . It is easy to see using the definition of $\Gamma_{n,s}$ as a group of homotopy equivalences that the image of the map $\mathbb{Z}^{4k} \rightarrow \text{Out}(F_{2k+2})$ is generated by automorphisms λ_{ij} and ρ_{ij} for $3 \leq i \leq 2k+2$, where λ_{ij} is left multiplication of the basis element x_i by x_j , and ρ_{ij} is right multiplication of x_i by x_j , with all basis elements other than x_i fixed in both cases. From this description one can see that the map $\mathbb{Z}^{4k} \rightarrow \text{Out}(F_{2k+2})$ is injective, as is its lift to $\text{Aut}(F_{2k+2})$, so the two versions of μ_k for Out and Aut are supported on \mathbb{Z}^{4k} subgroups. \square

5.2. Homology of $\Gamma_{2,4}$. If ϕ connects two copies of $X_{1,3}$ by gluing just one pair of leaves as in Figure 8, the result is a graph X_ϕ of rank 2 with 4 leaves and an assembly map

$$A_\phi: H_2(\Gamma_{1,3}) \otimes H_2(\Gamma_{1,3}) \longrightarrow H_4(\Gamma_{2,4}).$$

As before, let α_1 be a generator of $H_2(\Gamma_{1,3}) \cong P_{(1^3)}$. The image of $\alpha_1 \otimes \alpha_1$ under A_ϕ is then the nonvanishing class $\alpha_{I,J}$ in Proposition 4.1 in the case $k = 1$. By Theorem 3.10 we have $H_4(\Gamma_{2,4}) = P_{(2,2)}$, which is 2-dimensional. Since $P_{(2,2)}$ is irreducible as an \mathfrak{S}_4 -module, nontriviality of A_ϕ implies that the induced map

$$\hat{A}_\phi: \text{Res}_{\mathfrak{S}_2}^{\mathfrak{S}_3}(H_2(\Gamma_{1,3})) \circ \text{Res}_{\mathfrak{S}_2}^{\mathfrak{S}_3}(H_2(\Gamma_{1,3})) \longrightarrow H_4(\Gamma_{2,4})$$

is surjective.




Figure 8. Gluing two copies of $X_{1,3}$

5.3. Gluing two leaves of a single rank 1 graph. If ϕ glues two leaves of $X_{1,s}$ together as in Figure 9, the result is a graph X_ϕ of rank 2 with $s - 2$ leaves and an assembly map

$$A_\phi: H_k(\Gamma_{1,s}) \longrightarrow H_k(\Gamma_{2,s-2}).$$

Let us show that this A_ϕ is zero when $k > 0$.




Figure 9. A self-gluing of $X_{1,s}$

The map A_ϕ is an \mathfrak{S}_{s-2} -module map, where \mathfrak{S}_{s-2} is the subgroup of \mathfrak{S}_s which permutes the unglued leaves. For $k > 0$ either the domain or the range of A_ϕ is zero unless k is a multiple of 4, by Proposition 3.7 and Theorem 3.10. If $k = 4\ell > 0$, then $H_{4\ell}(\Gamma_{1,s}) = P_{(s-4\ell, 1^{4\ell})}$. Restriction from \mathfrak{S}_s to \mathfrak{S}_{s-2} removes two boxes from the Young diagram for $P_{(s-4\ell, 1^{4\ell})}$, so as an \mathfrak{S}_{s-2} -module the domain of A_ϕ is

$$P_{(s-4\ell-2, 1^{4\ell})} \oplus 2P_{(s-4\ell-1, 1^{4\ell-1})} \oplus P_{(s-4\ell, 1^{4\ell-2})}$$

(though if $s - 4\ell < 3$ some of the terms are not there). On the other hand, by Theorem 3.10 all partitions in $H_{4\ell}(\Gamma_{2,s-2})$ contain at least 2ℓ boxes in the second column, so none of these modules appears in $H_{4\ell}(\Gamma_{2,s-2})$ and A_ϕ must be zero.

5.4. Rank 2. For any positive s_1 and s_2 we can join X_{1,s_1} to X_{1,s_2} by connecting one pair of leaves. We obtain a graph X_ϕ of rank 2 with $s_1 + s_2 - 2$ leaves and assembly maps

$$A_\phi: H_{2k_1}(\Gamma_{1,s_1}) \otimes H_{2k_2}(\Gamma_{1,s_2}) \longrightarrow H_{2k_1+2k_2}(\Gamma_{2,s_1+s_2-2}).$$

If $k_1 + k_2$ is odd then $H_{2k_1+2k_2}(\Gamma_{2,s_1+s_2-2}) = 0$, so the map is obviously trivial. In fact this map is trivial unless $k_1 = k_2$, in which case it is nontrivial. This follows immediately from Proposition 4.1 and Remark 4.2.

In the special case $s_i = 2k_i + 1$ and $k_1 \neq k_2$ there is an alternative argument for proving the assembly map is zero using representation theory. From the discussion in the beginning of the section the assembly map induces a map

$$\begin{aligned} \text{Res}_{\mathfrak{S}_{2k_1}}^{\mathfrak{S}_{2k_1+1}}(H_{2k_1}(\Gamma_{1,2k_1+1})) \circ \text{Res}_{\mathfrak{S}_{2k_2}}^{\mathfrak{S}_{2k_2+1}}(H_{2k_2}(\Gamma_{1,2k_2+1})) \\ \longrightarrow H_{2k_1+2k_2}(\Gamma_{2,2k_1+2k_2}). \end{aligned}$$

By Proposition 3.7 and Theorem 3.10 we have $H_{2k_i}(\Gamma_{1,2k_i+1}) = P_{(1^{2k_i+1})}$ and $H_{2k_1+2k_2}(\Gamma_{2,2k_1+2k_2}) = P_{(2^{k_1+k_2})}$, so that the induced map is $P_{(1^{2k_1})} \circ P_{(1^{2k_2})} \rightarrow P_{(2^{k_1+k_2})}$. Since $k_1 \neq k_2$ there is no way to add $2k_2$ boxes to distinct rows in the Young diagram for $P_{(1^{2k_1})}$ to obtain the diagram for $P_{(2^{k_1+k_2})}$, which means that the decomposition of $P_{(1^{2k_1})} \circ P_{(1^{2k_2})}$ does not contain $P_{(2^{k_1+k_2})}$. So the map \hat{A}_ϕ (and hence the assembly map A_ϕ) must be zero. One can use a similar argument when $s_i \neq 2k_i + 1$ but this requires the full Littlewood–Richardson rule instead of the much easier Pieri rule.

5.5. Generalized Morita Classes. In [12], Morita's original series was generalized, and it is not hard to describe the generalization in terms of assembly maps arising from gluing together graphs of rank 0 and rank 1. Suppose we are given a finite connected graph G with no valence 2 vertices, along with a partition of its non-leaf vertices into two subsets V_0 and V_1 such that all vertices in V_1 have odd valence. Take a copy X_v of $X_{1,2k+1}$ for each vertex v in V_1 of valence $2k+1$, and identify the leaves of X_v with the edges of G incident to v . Similarly for each vertex in V_0 of valence k take a copy of $X_{0,k}$. The graph G then gives gluing instructions for constructing a graph $X_{n,s}$ and a corresponding assembly map. (The s leaves of $X_{n,s}$ come from the valence 1 vertices of G .) By assembling copies of $\alpha_k \in H_{2k}(\Gamma_{1,2k+1})$ at the valence $2k+1$ vertices in V_1 , along with standard generators for $H_0(\Gamma_{0,k})$ at the valence k vertices in V_0 , we obtain a *generalized Morita class* $\mu_G \in H_i(\Gamma_{n,s})$. The original

Morita classes are the case that G has two vertices, both in V_1 , with all edges going from one vertex to the other.

Vertices in V_1 are called *rank one* vertices, and vertices in V_0 are *rank zero* vertices. There is no loss of generality in assuming that all rank zero vertices are isolated, in the sense that no edge of G connects two different rank zero vertices, since such edges can be collapsed one by one without affecting the class μ_G .

If the graph G has certain “orientation-reversing” symmetries, for example if it has an edge with both ends at the same vertex in V_1 , then the class μ_G is automatically zero; this is spelled out in detail in [12] in the case $s = 0$, which is the case considered there. The following result shows that μ_G vanishes in many other cases as well.

Theorem 5.5. *If G has two rank one vertices of different valence then $\mu_G = 0$.*

Proof. The graph G contains a path connecting two rank one vertices of different valence and passing only through rank zero vertices. Using gluing instructions given by this linear subgraph we obtain a graph $X_\phi = X_{2,s}$ for some $s > 0$. We can also obtain an $X_{2,s}$ by first gluing one of the rank 1 graphs and all of the rank 0 graphs to obtain a graph X_{1,s_1} , then gluing X_{1,s_1} to the other rank 1 graph X_{1,s_2} . Call the latter gluing ψ . The assembly map A_ϕ factors through A_ψ , and we showed in Section 5.4 that A_ψ is zero since the two classes assembled by it have different homology degrees, by the hypothesis that the original two rank one vertices have different valence. \square

5.6. Eisenstein classes. Consider a gluing ϕ that attaches leaves of $X_{2,2k+2}$ to leaves of $X_{1,2k+1}$, leaving one leaf of $X_{2,2k+2}$ unpaired. This gluing determines a map $\Gamma_{2,2k+2} \times \Gamma_{1,2k+1} \rightarrow \Gamma_{2k+3,1} = \text{Aut}(F_{2k+3})$ and an assembly map

$$A_\phi : H_{2k+3}(\Gamma_{2,2k+2}) \otimes H_{2k}(\Gamma_{1,2k+1}) \longrightarrow H_{4k+3}(\text{Aut}(F_{2k+3})).$$

In Section 4.3 we constructed classes $m_{I,J} \in H_{2k+3}(\Gamma_{2,2k+2})$ corresponding to a partition of $\{1, \dots, 2k+2\}$ into sets I and J with $|I| = 2$ and $|J| = 2k$. (Note that the parameter k now corresponds to $k-1$ in Section 4.3.) Choose I to consist of the unglued leaf of $X_{2,2k+2}$ and one other leaf, with J the remaining leaves. Then the *Eisenstein class* \mathcal{E}_k is the image of $m_{I,J} \otimes \alpha_k$ under A_ϕ . This does not depend on the choice of the other leaf in I or the ordering of the leaves in J since permutations of the glued leaves in $X_{2,2k+2}$ become inner automorphisms of $\text{Aut}(F_{2k+3})$ which therefore induce the identity on homology.

We choose I to contain the unglued leaf because if we did not, then \mathcal{E}_k would automatically be zero by the following symmetry argument. If both leaves in $X_{2,2k+2}$ indexed by I were glued to leaves of $X_{1,2k+1}$ then the transposition switching these two leaves would extend to a transposition of two edges of the glued-together graph $X_{2k+3,1}$. This transposition sends $m_{I,J} \otimes \alpha_k$ to its negative since it preserves $m_{I,J}$ as we noted in Section 4.3 and it sends α_k to its negative. On the other hand, after

gluing, the transposition gives an inner automorphism of $\text{Aut}(F_{2k+3})$ inducing the identity on homology. Choosing I to contain the unglued leaf has the effect of breaking this symmetry, so \mathcal{E}_k does not vanish for any obvious reason. (Permutations of the leaves in J act trivially on \mathcal{E}_k since they act by their sign on both $m_{I,J}$ and α_k .)

An alternative construction would be to use the class m_i defined after the proof of Proposition 4.4 instead of $m_{I,J}$. Permutations of the glued leaves change m_i by the sign of the permutation, and the same is true for α_k , so $m_i \otimes \alpha_k$ is invariant under these permutations. From the definition of m_i it follows that using m_i instead of $m_{I,J}$ changes \mathcal{E}_k only by a nonzero scalar multiple.

Remark 5.6. The Eisenstein class \mathcal{E}_k maps to zero in $H_{4k+3}(\text{Out}(F_{2k+3}))$ since the map $\Gamma_{2k+3,1} \rightarrow \Gamma_{2k+3,0}$ is induced by forgetting the leaf of $X_{2k+3,1}$, and this leaf could just as well be omitted from $X_{2,2k+2}$ before the gluing, but this puts the class m_i in a dimension above the vcd of $\Gamma_{2,2k+1}$. This argument applies more generally whenever one has an assembly map with target $H_i(\Gamma_{n,1})$ and a source factor $H_{\text{vcd}}(\Gamma_{n_j,s_j})$ whose graph X_{n_j,s_j} is the one with the unglued leaf.

5.7. Odd-dimensional classes in $H_*(\text{Out}(F_n))$. The Euler characteristic calculations for $n \leq 11$ imply that there must exist odd-dimensional classes in $H_*(\text{Out}(F_n))$, probably in great abundance as n increases. However, only one such class has been found so it becomes an interesting challenge to find nontrivial odd-dimensional classes in a systematic way.




Figure 10. Left: Assembling $\gamma_1 \in H_{11}(\text{Out}(F_8))$ from $H_2(\Gamma_{1,3})$, $H_4(\Gamma_{1,5})$, and $H_5(\Gamma_{2,4})$. Right: Assembling $\gamma_2 \in H_{15}(\text{Out}(F_{10}))$ from $H_4(\Gamma_{1,5})$, $H_6(\Gamma_{1,7})$, and $H_5(\Gamma_{2,4})$

A sequence of candidates for such classes was introduced by Morita, Sakasai, and Suzuki in [35], Proposition 6.3. These are classes $\gamma_k \in H_{4k+7}(\text{Out}(F_{2k+6}))$ for $k \geq 1$. The class γ_k can be interpreted as gluing $X_{1,2k+3} \cup X_{1,2k+1} \cup X_{2,4} \rightarrow X_{2k+6,0}$ (illustrated for $k = 1$ and 2 in Figure 10) and considering the image of the class $\alpha_{k+1} \otimes \alpha_k \otimes m_i$, where m_i is the class defined after the proof of Proposition 4.4, with i labeling the leaf of $X_{2,4}$ attached to $X_{1,2k+1}$. However, since the image of

$\alpha_{k+1} \otimes \alpha_k$ in $H_{4k+2}(\Gamma_{2,4k-2})$ is trivial by Section 5.4, these classes must also be zero by associativity of the assembly map.




Figure 11. Candidates for a nontrivial class in $H_{11}(\text{Out}(F_8))$

The class $\gamma_1 \in H_{11}(\text{Out}(F_8))$ was introduced to account for the fact that $\mu_3 \in H_{12}(\text{Out}(F_8))$ is nonzero while the Euler characteristic of $\text{Out}(F_8)$ is 1, so an odd-dimensional class must exist. An alternative candidate class in $H_{11}(\text{Out}(F_8))$ can be constructed by assembling two copies of $\alpha_1 \in H_2(\Gamma_{1,3})$ with a class in $H_7(\Gamma_{2,6})$ by gluing all the leaves of two copies of $X_{1,3}$ to one copy of $X_{2,6}$ as shown in the left half of Figure 11. Another possibility is to glue all the leaves of $X_{1,1}$ and $X_{1,5}$ to the leaves of $X_{2,6}$, obtaining an assembly map $H_0(\Gamma_{1,1}) \otimes H_7(\Gamma_{2,6}) \otimes H_4(\Gamma_{1,5}) \rightarrow H_{11}(\text{Out}(F_8))$ as in the right half of Figure 11. However, this assembly map may well be zero since it produces classes that lift to classes in $H_{11}(\text{Aut}(F_8))$ that are in the image of the stabilization $H_{11}(\text{Aut}(F_7)) \rightarrow H_{11}(\text{Aut}(F_8))$ (see Section 6.2) so if they were nonzero they would give counterexamples to Conjecture 10.5.




Figure 12. Candidates for a nontrivial class in $H_{15}(\text{Out}(F_{10}))$

A few other ways to construct candidates for odd-dimensional classes are shown in Figure 12, this time for classes in $H_{15}(\text{Out}(F_{10}))$.

There are many other ways to construct candidates for odd-dimensional homology classes. If we glue $X_{1,s}$ to $X_{2,s}$ by joining all of their leaves, we obtain a rank $s + 2$ graph with no leaves and an associated assembly map

$$H_2(\Gamma_{1,s}) \otimes H_5(\Gamma_{2,s}) \longrightarrow H_7(\Gamma_{s+2,0}).$$

For $s \geq 4$ we have $H_2(\Gamma_{1,s}) = P_{(s-2,1^2)}$. The decomposition of $H_5(\Gamma_{2,s}) = W_4 \circ P_{(s-4)} = P_{(2,1^2)} \circ P_{(s-4)}$ has several terms but one of them is $P_{(s-2,1^2)}$, with multiplicity one. Thus the space of coinvariants $(H_2(\Gamma_{1,s}) \otimes H_5(\Gamma_{2,s}))_{\mathfrak{S}_s}$ is 1-dimensional and the assembly map, which factors through these coinvariants, may well be nontrivial. This construction produces potential classes in $H_7(\text{Out}(F_n))$ for all $n \geq 6$, although by homology stability these classes must be trivial for $n \geq 10$. According to the calculations in [37] and [1] the classes for $n = 6, 7$ also vanish.

We can make a similar construction with $H_{2k}(\Gamma_{1,s})$ and $H_{2k+3}(\Gamma_{2,s})$ as long as $s \geq 2k + 2$ using the partition $(s - 2k, 1^{2k})$. We have $H_{2k}(\Gamma_{1,s}) = P_{(s-2k,1^{2k})}$ and $H_{2k+3}(\Gamma_{2,s}) = W_{2k+2} \circ P_{(s-2k-2)}$. The second term of W_{2k+2} is $M_{2k+2} \otimes P_{(2,1^{2k})}$, and one term of $P_{(2,1^{2k})} \circ P_{(s-2k-2)}$ is equal to $P_{(s-2k,1^{2k})}$. Thus we obtain a map from \mathcal{M}_{2k+2} to $H_{4k+3}(\text{Out}(F_{s+2}))$. For large s this map must be trivial since the target group is trivial. In fact we suspect that the map is trivial for $s \geq 2k + 4$ but not for $s = 2k + 3$. For $s = 2k + 3$ the image lies in $H_{4k+3}(\text{Out}(F_{2k+5}))$. Since the vcd of $\text{Out}(F_{2k+5})$ is $4k + 7$, these classes lie in codimension 4.

If $s \geq 2k + 1$ there is another similar construction with $H_{2k}(\Gamma_{1,s})$ and $H_{2k+1}(\Gamma_{2,s})$, again using the partition $(s - 2k, 1^{2k})$. Here $H_{2k+1}(\Gamma_{2,s})$ contains the summand $S_{2k+2} \otimes P_{(1^{2k})} \circ P_{s-2k}$. Since $P_{(1^{2k})} \circ P_{(s-2k)}$ contains a copy of $P_{(s-2k,1^{2k})}$ we get a map from \mathcal{S}_{2k+2} to $H_{4k+1}(\text{Out}(F_{s+2}))$. The first potentially nontrivial class occurs when $k = 5$, in $H_{21}(\text{Out}(F_{13}))$; this class is in $H_{\text{vcd}-2}$.

5.8. Classes from pairs of cusp forms. Consider a gluing

$$X_{2,2m} \cup X_{2,2m} \longrightarrow X_{2m+3,0}$$

matching the leaves of the first graph with those of the second, and the associated assembly map

$$H_{2m+1}(\Gamma_{2,2m}) \otimes H_{2m+1}(\Gamma_{2,2m}) \longrightarrow H_{4m+2}(\text{Out}(F_{2m+3})).$$

There is a \mathbb{Z}_2 action on $H_{2m+1}(\Gamma_{2,2m}) \otimes H_{2m+1}(\Gamma_{2,2m})$ which switches the factors. By basic properties of the cross product (which gives the Künneth isomorphism) we have

$$a \otimes b = (-1)^{(2m+1)(2m+1)} b \otimes a = -b \otimes a,$$

so this assembly map factors through the exterior product $\bigwedge^2 H_{2m+1}(\Gamma_{2,2m})$ in addition to factoring through the \mathfrak{S}_{2m} -coinvariants.

Applying Lemma 3.8 we now have a map

$$\begin{aligned} \left(\bigwedge^2 H_{2m+1}(\Gamma_{2,2m}) \right)_{\mathfrak{S}_{2m}} &= \left(\bigwedge^2 \left(\bigoplus_{0 \leq i < m} \mathcal{X}_{2m,i} \otimes P_{(2^i, 1^{2m-2i})} \right) \right)_{\mathfrak{S}_{2m}} \\ &\longrightarrow H_{4m+2}(\text{Out}(F_{2m+3})), \end{aligned}$$

where $\mathcal{X}_{2m,i}$ is the space of either cusp forms (if i is even) or all modular forms (if i is odd) of weight $2m+2-2i$. Since the $P_{(2^i, 1^{2m-2i})}$ are pairwise non-isomorphic this gives a map

$$\bigoplus_{0 \leq i < m} \left(\bigwedge^2 \mathcal{X}_{2m,i} \right) \longrightarrow H_{4m+2}(\text{Out}(F_{2m+3})).$$

The term corresponding to $i = 0$ was first mentioned in [10].

6. Stabilization

In this section we consider the two ways to stabilize $\Gamma_{n,s}$ by letting one of the parameters n and s increase while keeping the other fixed. Both stabilizations can be viewed as special cases of assembly maps.

6.1. Stabilization with respect to s . For $s \geq 1$ the map gluing $X_{0,3}$ to $X_{n,s}$ by a single edge simply increases the number of leaves, and the associated assembly map

$$H_0(\Gamma_{0,3}) \otimes H_i(\Gamma_{n,s}) \longrightarrow H_i(\Gamma_{n,s+1})$$

gives a stabilization map $H_i(\Gamma_{n,s}) \rightarrow H_i(\Gamma_{n,s+1})$ sending a class α to $\iota \otimes \alpha$ where ι is the standard generator of $H_0(\Gamma_{0,3})$. This is the same as the map induced by the splitting of the natural projection $\Gamma_{n,s+1} \rightarrow \Gamma_{n,s}$ defined in the proof of Proposition 2.2. Strictly speaking, there are s different stabilization maps depending on which leaf of $X_{n,s}$ we attach $X_{0,3}$ to, although these stabilizations differ only by the action of \mathfrak{S}_s . The stabilization maps are always injective since they are induced by splittings of the groups. By [24] the stabilization map is surjective if $n \geq 2i + 2$, but in this case the homology groups are trivial by Galatius' theorem.

Switching from homology to cohomology, Proposition 3.7 and Theorem 3.10 demonstrate directly that for fixed i the groups $H^i(\Gamma_{1,s})$ and $H^i(\Gamma_{2,s})$ satisfy *representation stability* as s increases, i.e., for large enough s the partitions which appear in their irreducible decompositions as \mathfrak{S}_s -modules differ only by the number of boxes in the first row. This leads one to suspect that the cohomology of $\Gamma_{n,s}$ satisfies representation stability for all n . This is indeed the case and can be deduced easily from a theorem of Jiménez Rolland [26] giving the corresponding result for mapping class groups of certain manifolds with punctures.

Proposition 6.1. *For fixed i and n the groups $H^i(\Gamma_{n,s})$ satisfy representation stability as s increases.*

Proof. As described in Section 2.1 we can view $\Gamma_{n,s}$ as the quotient of the mapping class group of the 3-manifold $M_{n,s}$ by the subgroup generated by Dehn twists along 2-spheres. This subgroup is normal and is just a direct product of finitely many cyclic groups of order 2. In particular it is a finite group so the projection from the mapping class group to $\Gamma_{n,s}$ induces an isomorphism on cohomology with coefficients in \mathbb{k} by the Leray–Serre spectral sequence. Thus it suffices to prove representation stability for the mapping class group, and this was done in [26], with a specific stable range $s \geq 3i$. To apply [26] one uses the manifold $M = M_{n,0}$ as the base manifold, and one needs to check that this satisfies certain hypotheses: (1) $\pi_1 M = F_n$ is of type FP_∞ and has trivial center, which is obviously true; and (2) the mapping class group of M is of type FP_∞ , which follows from $\text{Out}(F_n)$ being FP_∞ and the kernel of the map from the mapping class group to $\text{Out}(F_n)$ being finite abelian and hence of type FP_∞ . \square

We remark that A. Saied has recently shown that $H^i(\Gamma_{n,s})$ satisfies representation stability with respect to s whenever $s \geq n + i$ [38]. If $n < 2i$ this is an obvious improvement on the stable range mentioned in the above proposition, and if $n \geq 2i + 2$ the cohomology is zero for all s .

6.2. Stabilization with respect to n . A stabilization map $H_i(\Gamma_{n,s}) \rightarrow H_i(\Gamma_{n+1,s})$ can be obtained in a similar way by gluing $X_{1,2}$ to $X_{n,s}$ along one edge. Here sufficiently many iterations take one to the stable range where the homology groups are zero, so the interest is in what happens unstably. We can describe completely what happens when rank one classes are stabilized to rank two:

Proposition 6.2. *If a class in $H_i(\Gamma_{n,s})$ is obtained from an assembly map with a factor group $H_j(\Gamma_{1,k})$, $j > 0$, then this class maps to zero under the stabilization $H_i(\Gamma_{n,s}) \rightarrow H_i(\Gamma_{n+1,s})$ obtained by gluing $X_{1,2}$ to the corresponding factor graph $X_{1,k}$.*

An immediate consequence is the following result, first proved in [13] by combinatorial arguments.

Corollary 6.3. *The Morita class μ_k , lifted from Out to Aut , vanishes under the stabilization map $H_{4k}(\Gamma_{2k+2,1}) \rightarrow H_{4k}(\Gamma_{2k+3,1})$.*

Proof of Proposition 6.2. It suffices to prove that the stabilization $H_i(\Gamma_{1,s}) \rightarrow H_i(\Gamma_{2,s})$ is trivial for $i > 0$. The extension of $A_\phi: H_i(\Gamma_{1,s}) \otimes H_0(\Gamma_{1,2}) \rightarrow H_i(\Gamma_{2,s})$ to the \mathfrak{S}_s -module map

$$\hat{A}_\phi: \text{Res}_{\mathfrak{S}_{s-1}}^{\mathfrak{S}_s}(H_i(\Gamma_{1,s})) \circ \text{Res}_{\mathfrak{S}_1}^{\mathfrak{S}_2}(H_0(\Gamma_{1,2})) \longrightarrow H_i(\Gamma_{2,s})$$

is accomplished by first restricting the \mathfrak{S}_s -action on $\Gamma_{1,s}$ to \mathfrak{S}_{s-1} and then inducing it back up to \mathfrak{S}_s . In terms of Young diagrams, restriction to \mathfrak{S}_{s-1} is accomplished by removing one box (in all possible ways), while induction adds a box (also in all possible ways). Since the diagrams of all partitions appearing in $H_i(\Gamma_{1,s})$ have boxes only in the first row and column, all resulting diagrams will have at most two boxes in the second column.

Using Theorem 3.10 we can see that the simple modules appearing in $H_i(\Gamma_{2,s})$ for i even always have at least $i/2$ boxes in the second column. Thus if $i \geq 6$ there is no partition which appears in both the domain and range of \hat{A}_ϕ , which forces \hat{A}_ϕ (and therefore A_ϕ) to be zero.

For $i = 2$ the target $H_2(\Gamma_{2,s})$ is always zero. For $i = 4$ the diagrams appearing in $H_4(\Gamma_{1,s})$ have five rows, but the diagrams appearing in $H_4(\Gamma_{2,s})$ have at most three rows, so it is not possible to obtain one from the other by changing the position of a single box, and again the assembly map must be trivial. \square

In Section 7 below we show precisely how the class α_k becomes trivial after one stabilization, using a natural geometric interpretation of this class.

7. A more geometric viewpoint

A rational model for $B\text{Out}(F_n)$ is the quotient Q_n of Outer Space for rank n graphs by the action of $\text{Out}(F_n)$ changing the marking. (One could instead use just the spine of Outer Space, but for our present purposes it is more convenient not to restrict to the spine.) Points of Q_n are thus isometry classes of finite connected graphs of rank n with no vertices of valence 1 or 2 and with lengths assigned to the edges, normalized so that the sum of the lengths of all the edges is 1. Collapsing edges to points by shrinking their lengths to zero is allowed provided this does not decrease the rank of the graph. There is a similar rational model $Q_{n,s}$ for $B\Gamma_{n,s}$ consisting of graphs in Q_n with s leaves attached to them at arbitrary points. There is no need to assign lengths to the leaf edges since they are not allowed to collapse to points. An assembly map is induced from a map $Q_{n_1,s_1} \times \cdots \times Q_{n_k,s_k} \rightarrow Q_{n,s}$ where edge lengths on a glued-together graph $X_{n,s}$ are obtained by first assigning a fixed length, say 1, to the new edges created by the leaf pairings, then renormalizing the lengths of all the nonleaf edges of $X_{n,s}$.

7.1. Geometric Morita cycles. The classes $\alpha_k \in H_{2k}(\Gamma_{1,2k+1})$ are particularly easy to describe from this perspective, and hence also the Morita classes and their generalizations. The class α_k is the image of the top-dimensional homology class of a $2k$ -dimensional torus under a map $f: T^{2k} \rightarrow Q_{1,2k+1}$ described as follows. Consider graphs $X_{1,2k+1}$ obtained from a circle c by attaching $2k + 1$ leaves. By rotating the circle if necessary, we can assume the first leaf attaches at a fixed

basepoint of c . The other leaves attach at $2k$ arbitrary points of c which need not be distinct. Letting these points vary independently around c then gives the map $f: T^{2k} \rightarrow Q_{1,2k+1}$. This is surjective but not injective since graphs differing by a reflection of c fixing the basepoint are identified in $Q_{1,2k+1}$, so $Q_{1,2k+1}$ is T^{2k} modulo the action of \mathbb{Z}_2 reflecting each circle factor. From this point of view one can see why we require the total number of leaves to be odd, because if it were $2k$ instead of $2k + 1$ then the \mathbb{Z}_2 -action would reverse the orientation of the torus T^{2k-1} and hence the map $T^{2k-1} \rightarrow Q_{1,2k}$ would induce the trivial map on the top-dimensional homology of the torus.

For the map $Q_{1,2k+1} \times Q_{1,2k+1} \rightarrow Q_{2k+2,0}$ used to construct the Morita class $\mu_k = \alpha_k \otimes \alpha_k$ we glue all the leaves of the first copy of $X_{1,2k+1}$ to the leaves of the second copy. Thus we have two circles joined by $2k + 1$ edges. One of these edges serves as a “basepoint” edge, and then by varying where the remaining $2k$ edges attach we obtain a family of graphs $X_{2k+2,0}$ corresponding to a map $T^{4k} \rightarrow Q_{2k+2,0}$ taking a generator of $H_{4k}(T^{4k})$ to μ_k . (The basepoint edge could be collapsed to a point, giving a map $T^{4k} \rightarrow Q_{2k+2,0}$ homotopic to the original one.) The generalized Morita classes have similar geometric descriptions as maps from a torus to the appropriate $Q_{n,s}$.

For the Morita class μ_k the map $T^{4k} \rightarrow Q_{2k+2,0}$ is invariant under certain symmetries. To start, there is the $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry coming from the symmetries of the two α_k factors reflecting each of the two circles. There is another \mathbb{Z}_2 symmetry from interchanging the two circles. Finally, there is an \mathfrak{S}_{2k} symmetry group permuting the $2k$ arcs connecting the two circles. Altogether this gives a symmetry group G_k of order $8(2k)!$ with the map $T^{4k} \rightarrow Q_{2k+2,0}$ factoring through the quotient T^{4k}/G_k . One can regard G_k as acting on choices of an ordering and orientations of the two circles and an ordering of the $2k$ connecting arcs. This makes it clear that the induced map $T^{4k}/G_k \rightarrow Q_{2k+2,0}$ is injective.

In the case $k = 1$ the quotient T^4/G_1 can be determined explicitly.

Proposition 7.1. $T^4/G_1 = S^4$.

Proof. The quotient of T^4 by the reflections of the two circles gives $S^2 \times S^2$ since the quotient of T^2 by reflection of its two circle factors is the familiar 2-sheeted branched covering space $T^2 \rightarrow S^2$. Next, factor out the \mathbb{Z}_2 -action interchanging the two circles, corresponding to interchanging the two factors of $S^2 \times S^2$. This gives the two-fold symmetric product $SP_2(S^2)$ which is well-known to be $\mathbb{C}P^2$. Explicitly, $\mathbb{C}P^2$ can be identified with nonzero polynomials $a_2 z^2 + a_1 z + a_0$ in $\mathbb{C}[z]$ up to scalar multiplication, and these are determined by their unordered pair of roots in $S^2 = \mathbb{C} \cup \infty$ where linear factors corresponding to roots at ∞ are deleted. (See e.g. [21, Example 4K.4]) Finally we need to factor out by the \mathbb{Z}_2 -action interchanging the two connecting arcs. This corresponds to reflecting each torus T^2 across its diagonal. In the quotient $S^2 \times S^2$ of $T^2 \times T^2$ this is equivalent to reflecting each S^2 across its equator. In the space of quadratic polynomials this is given by complex

conjugation of the roots, hence also of the coefficients. Thus we are forming the quotient of $\mathbb{C}P^2$ by complex conjugation. This quotient is S^4 by a classical result of Massey [32] and Kuiper [30]. \square

The quotient T^{4k}/G_k for $k > 1$ cannot be a sphere since one can compute that its rational homology consists of a copy of \mathbb{Q} in each dimension $4i \leq 4k$.

We can use the geometric viewpoint to give another proof that Morita classes and their generalizations vanish after one stabilization of the rank:

Geometric proof that α_k vanishes after one stabilization. The idea is to see how the commutator relation $[e_{ij}, e_{jk}] = e_{ik}$ among elementary matrices can be translated into a two-parameter family of graphs.

We can reinterpret the stabilization map as the map $Q_{1,2k+1} \rightarrow Q_{2,2k+1}$ obtained by attaching both ends of a new edge b at the basepoint of the graphs $X_{1,2k+1}$ described above consisting of a circle c with $2k + 1$ leaves attached, where by the basepoint we mean the point of c where the fixed leaf attaches. Let a be any one of the remaining $2k$ leaves. Sliding a around c gives one of the S^1 factors of the torus T^{2k} whose map to $Q_{1,2k+1}$ sends a generator of $H_{2k}(T^{2k})$ to α_k .

Figure 13 describes a two-parameter family of graphs in which one end of the arc b moves across c while one end of the arc a moves across b and c . As we proceed from left to right in the sequence of four pictures we see one end of b sliding around c . The dotted arc denotes the path followed by the attaching point of a . Initially it just goes across b , then when b has moved partway around c the end of a must backtrack across part of c after it crosses b , in order to return to the basepoint. In the last picture we see that a crosses both b and c .




Figure 13. A 2-parameter family of graphs $X_{2,1}$

On the boundary of the parameter square for this two-parameter family one thus has five slides of one arc over another, as indicated in Figure 14. The quotient space of the square obtained by identifying the two a/b edges and the two b/c edges is a surface $S_{1,1}$ of genus one with one boundary circle, where this boundary circle parametrizes the a/c slide. The a/c slide was the restriction of the map $f: T^{2k} \rightarrow Q_{2,2k+1}$ representing the stabilization of α_k to one of the circle factors of T^{2k} , so we can extend f to a map $T^{2k-1} \times S_{1,1} \rightarrow Q_{2,2k+1}$. This implies that f induces the zero map $H_{2k}(T^{2k}) \rightarrow H_{2k}(Q_{2,2k+1})$. \square




Figure 14. The parameter space, a punctured torus

7.2. Representing Eisenstein classes geometrically. Let us describe how the Eisenstein classes $\mathcal{E}_k \in H_{4k+3}(\text{Aut}(F_{2k+3}))$ can be realized as families of graphs parametrized by certain manifolds E_k that are analogous to the tori representing Morita classes, but a little more complicated. Consider first the case $k = 1$, so E_1 will be a closed orientable 7-manifold. This is the product of T^4 with a 3-manifold N , where N splits along a 2-torus into submanifolds N_1 and N_2 each homeomorphic to a product of S^1 with a compact surface $S_{1,1}$ of genus 1 with one boundary component. We obtain N from $N_1 \sqcup N_2$ by gluing the two boundary tori via a homeomorphism of $\partial N_1 = \partial N_2 = T^2$ switching the two circle factors of T^2 . From this description one can see that $\pi_1 E_1$ is the product of \mathbb{Z}^4 with two copies of $\mathbb{Z} \times F_2$ amalgamated along $\mathbb{Z} \times \mathbb{Z}$ where the second \mathbb{Z} is generated by the commutator of the generators of F_2 and the amalgamation interchanges the two factors of $\mathbb{Z} \times \mathbb{Z}$. Also from the construction of E_1 one can easily see that it is a $K(\pi, 1)$.

Now we describe a map $E_1 \rightarrow Q_{5,1}$ corresponding to a 7-dimensional family of rank 5 graphs with one leaf. To construct these graphs, start with the family of graphs parametrized by $S_{1,1}$ indicated in Figure 13, consisting of a circle c with edges a and b attached. One end of b is attached to the basepoint of c and the other end to a point moving around c . The arc a attaches at one end to a point that moves across b and then returns to the basepoint along an arc of c . Next we attach one end of another arc a' at a point that moves only around c , independently of how a and b attach. This gives a family of graphs parametrized by $S^1 \times S_{1,1}$. Reversing the roles of a and a' gives another family parametrized by $S^1 \times S_{1,1}$. On $S^1 \times \partial S_{1,1}$ both families consist of graphs in which $b \cup c$ is $S^1 \vee S^1$ with a and a' attached to arbitrary points of c . The two families parametrized by $S^1 \times S_{1,1}$ then fit together to form a family parametrized by N , so we have a map $N \rightarrow Q_{2,2}$. Attaching two more arcs d and e at an endpoint of each that moves freely around c gives a family parametrized by $N \times T^2$ and so a map $N \times T^2 \rightarrow Q_{2,4}$, hence a class $m \in H_5(\Gamma_{2,4})$. Finally, we assemble this class m with the torus T^2 representing $\alpha_1 \in H_2(\Gamma_{1,3})$ by adjoining another circle c' and attaching the free ends of a , d , and e at points that move around c' , where by

rotating c' we can assume that a attaches just at the basepoint of c' . The arc a' has one end unattached, so it is a leaf. Figure 15 shows a graph in the resulting 7-parameter family in the case that a' attaches to c , but a' could also attach to b when a attaches to c . These two possibilities correspond to the two submanifolds N_1 and N_2 of N .




Figure 15. A graph in the \mathcal{E}_1 family

To relate this to the earlier construction of Eisenstein classes, note first that the group $M_{I,J}$ in Proposition 4.4 is the semidirect product $\mathbb{Z} \ltimes (F_2^2 \times \mathbb{Z}^2)$ in the case at hand when $|I| = 2 = |J|$. A $K(\pi, 1)$ for this group is the mapping torus T_f of a map f from $(S^1 \vee S^1)^2 \times T^2$ to itself that is the identity on the T^2 factor, and on each $S^1 \vee S^1$ induces the automorphism of $\pi_1(S^1 \vee S^1) = F_2$ fixing the first basis element x and sending the second basis element y to xy . We can compute $H_5(T_f)$ geometrically as the elements of $H_4((S^1 \vee S^1)^2 \times T^2)$ fixed by f_* using the exact sequence in Example 2.48 of [21]. This H_4 has dimension 4 with basis corresponding to the 4-cells of $(S^1 \vee S^1)^2 \times T^2$ by the Künneth formula. The basis elements can be written (omitting tensor product symbols for simplicity) as $xxzz$, $xyzz$, $yxzz$, and $yyzz$. (It would make sense to simplify the notation further by replacing z by x throughout.) The map f_* fixes x and z and takes y to $x + y$. A short calculation shows that the elements invariant under f_* form a 2-dimensional subspace with basis $xxzz$ and $xyzz - yxzz$. The element $xxzz$ is not of interest to us since it can be shown to give the trivial element of $H_5(\Gamma_{2,4})$. For the element $xyzz - yxzz$ the four letters of each of these two words correspond to the edges a, a', d, e in that order. For example the four letters of $xyzz$ correspond to a moving around c , a' moving across b , and d and e moving around c . Interchanging a and a' gives the other word $yxzz$. To go from $H_4((S^1 \vee S^1)^2 \times T^2)$ to $H_5(T_f)$ involves a fifth parameter, and this corresponds to one end of the edge b moving around c . The whole mapping torus T_f corresponds to a 5-dimensional family of graphs in which a and a' move freely around both b and c , while d, e , and one end of b move around c . The reason for restricting to the graphs parametrized by $N \times T^2$ is to get a manifold as parameter space and thus reduce the dimension of H_5 from two to one.

To generalize from \mathcal{E}_1 to \mathcal{E}_k is easy since all one has to do is replace the arcs d and e by $2k$ arcs d_1, \dots, d_k and e_1, \dots, e_k that behave in exactly the same way as d and e . Thus $E_k = N \times T^{4k}$. Incidentally, the manifold N belongs to the class of 3-manifolds known as graph manifolds, which seems an especially appropriate name in the present context.

One can see that $\mathcal{E}_k \in H_{4k+3}(\Gamma_{2k+3,1})$ maps to zero in $H_{4k+3}(\Gamma_{2k+3,0})$ since ignoring the leaf a' replaces the 3-manifold N by a 2-dimensional quotient, namely $S_{1,1}$, so the composition $E_k \rightarrow Q_{2k+3,1} \rightarrow Q_{2k+3,0}$ factors through $S_{1,1} \times T^{4k}$ which has one lower dimension than E_k .

8. Homological triviality of a standard “maximal torus” for $\text{Aut}(F_n)$

Let A be the subgroup of $\text{Aut}(F_n)$ generated by the automorphisms λ_i and ρ_i for $1 \leq i \leq n-1$, where λ_i sends the basis element x_i to $x_n x_i$ and fixes x_j for $j \neq i$, and ρ_i is defined similarly but multiplies x_i by x_n on the right. Thus A is isomorphic to \mathbb{Z}^{2n-2} , realizing the maximal rank of an abelian subgroup since the vcd of $\text{Aut}(F_n)$ is $2n-2$.

Theorem 8.1. *The inclusion $A \hookrightarrow \text{Aut}(F_n)$ of the standard free abelian subgroup of maximal rank induces the trivial map on rational homology in all positive dimensions.*

Note that the $\mathbb{Z}^{2n-4} \subset \text{Aut}(F_n)$ realizing the Morita class μ_k is not contained in this $\mathbb{Z}^{2n-2} \subset \text{Aut}(F_n)$ or any subgroup conjugate to this by permuting basis elements for F_n , and the theorem gives a good reason why this must be the case.

We will give three different proofs of this theorem, each with its own advantages. The first proof is probably the most elementary.

Algebraic proof. We can enlarge A to a subgroup $G \subset \text{Aut}(F_n)$ by adjoining the automorphisms that permute the basis elements x_1, \dots, x_{n-1} and send a subset of them to their inverses. These automorphisms form a copy of the signed permutation group \mathfrak{S}_{n-1}^\pm in $\text{Aut}(F_n)$, and G is the semidirect product $\mathfrak{S}_{n-1}^\pm \ltimes A$. It will suffice to show $H_i(G) = 0$ for $i > 1$ since $H_1(\text{Aut}(F_n)) = 0$ from the classical presentations of $\text{Aut}(F_n)$.

Passing from homology to cohomology and applying the usual argument with transfer homomorphisms, we can compute $H^*(G)$ as the invariants of $H^*(A)$ under the action of \mathfrak{S}_{n-1}^\pm induced by conjugation. The cohomology ring $H^*(A)$ is an exterior algebra on generators a_i and b_i corresponding to λ_i and ρ_i . Conjugation by the map inverting x_i sends λ_i to ρ_i^{-1} and ρ_i to λ_i^{-1} , so in $H^*(A)$ this sends a_i to $-b_i$ and b_i to $-a_i$. Conjugation by a permutation of the x_i 's has the effect of permuting the subscripts on the a_i 's and b_i 's.

Elements of $H^k(A)$ are linear combinations of degree k monomials in the a_i 's and b_i 's. We will show that for any monomial m of degree $k > 1$ there exists $\sigma \in \mathfrak{S}_{n-1}^\pm$ such that $\sigma m = -m$. This implies that m cannot appear in any element of $H^k(A)$ that is invariant under the action of \mathfrak{S}_{n-1}^\pm , and hence $H^k(G) = 0$ for $k > 1$. There are three cases: if m contains both a_i and b_i for some i , then inverting x_i changes the sign of $a_i \wedge b_i$ and thus the sign of m ; if m contains a_i and a_j but not either of b_i or b_j we use the involution in G interchanging x_i and x_j ; and if m contains a_i and b_j but not either of b_i or a_j we use the involution in G

interchanging x_i and x_j^{-1} . (This argument does not apply when $k = 1$, but it is easy to check that $H^1(G) = \mathbb{k}$ generated by $\sum_i a_i - \sum_i b_i$.) \square

Geometric proof. As in the previous section we consider the rational model $Q_{n,1}$ for $\text{Aut}(F_n)$. The inclusion $\mathbb{Z}^{2n-2} \hookrightarrow \text{Aut}(F_n)$ corresponds to a map $f: T^{2n-2} \rightarrow Q_{n,1}$ of the $(2n-2)$ -torus to $Q_{n,1}$. This specifies a family of graphs parametrized by T^{2n-2} constructed as follows. Start with a basepointed circle c , then attach $n-1$ arcs a_i by identifying their endpoints with points s_i and t_i in c . The s_i and t_i are the coordinates on T^{2n-2} , and we can write f as a function $f(s_1, t_1, \dots, s_{n-1}, t_{n-1})$.

The map f is not injective since there are some symmetries present. One can interchange s_i and t_i , switching the ends of a_i , without changing the graph, and one can permute the arcs a_i . Switching s_i and t_i gives a quotient of the i th 2-torus factor of $T^{2n-2} = (T^2)^{n-1}$. The quotient of a 2-torus by interchanging the two circle factors is a triangle with two edges identified. This deformation retracts to a single circle, say the s_i circle. The quotient of T^{2n-2} by these coordinate transpositions thus has the homotopy type of T^{n-1} . This already implies that the inclusion $A \hookrightarrow \text{Aut}(F_n)$ induces the trivial map on H_k for $k > n-1$.

Now we can factor out the permutations of the $n-1$ factors of this T^{n-1} , producing the $(n-1)$ -fold symmetric product of S^1 . This is well known to have the homotopy type of a single circle. (See for example the end of Example 4K.4 in [21].) Thus the map f factors through a space homotopy equivalent to S^1 so it induces the trivial map on H_i for $i > 1$. It also induces the trivial map on H_1 since $H_1(\text{Aut}(F_n)) = 0$ as noted in the first proof. \square




Figure 16. The gluings $X_{1,2n-1} \rightarrow X_{2,2n-3} \rightarrow X_{n,1}$

Proof via representation theory. We have inclusions $A \subset \Gamma_{1,2n-1} \subset \text{Aut}(F_n)$ where the second inclusion corresponds to the self-gluing $X_{1,2n-1} \rightarrow X_{n,1}$. (In fact $\Gamma_{1,2n-1}$ is contained in the subgroup G used in the first proof above). The gluing $X_{1,2n-1} \rightarrow X_{n,1}$ factors as the composition of two gluings $X_{1,2n-1} \rightarrow X_{2,2n-3} \rightarrow X_{n,1}$ as shown in Figure 16, so the map $H_*(A) \rightarrow H_*(\text{Aut}(F_n))$ factors as $H_*(A) \rightarrow H_*(\Gamma_{1,2n-1}) \rightarrow H_*(\Gamma_{2,2n-3}) \rightarrow H_*(\Gamma_{n,1})$. The middle of these three maps is an

assembly map which we showed is trivial in Section 5.3 (unless the degree is 0), which implies that $H_k(A) \rightarrow H_k(\text{Aut}(F_n))$ is trivial if $k > 0$. \square

One of the advantages of this last proof is that it also works in degree 1, so one does not need a separate argument for this case.

Remark 8.2. The composition $A \rightarrow \text{Aut}(F_n) \rightarrow \text{Out}(F_n)$ has kernel \mathbb{Z} and is injective when restricted to a suitable subgroup \mathbb{Z}^{2n-3} realizing the vcd of $\text{Out}(F_n)$. The theorem implies that this inclusion $\mathbb{Z}^{2n-3} \hookrightarrow \text{Out}(F_n)$ is also trivial on homology since it factors through $A \hookrightarrow \text{Aut}(F_n)$.

9. Connections with hairy graph homology and the Lie algebra of symplectic derivations

9.1. Hairy graph homology. In this section we note the connection between our calculations and the hairy graph homology theory of [10]. As above \mathfrak{S}_k denotes the symmetric group on k letters and Sym^k the k -th symmetric power functor on vector spaces. The following lemma is an immediate consequence of Proposition 3.6.

Lemma 9.1. *Let $H = H^1(F_n) \cong \mathbb{k}^n$. For any \mathbb{k} -vector space V*

$$H^{2n-3+s}(\Gamma_{n,s}) \otimes_{\mathfrak{S}_s} V^{\wedge s} \cong H^{2n-3}(\text{Out}(F_n); \text{Sym}^s(H \otimes V)).$$

Proof. The Künneth isomorphism $H^s(F_n^s) \cong H^1(F_n)^{\wedge s} = H^{\wedge s}$ is $\text{Out}(F_n) \times \mathfrak{S}_s$ -equivariant. So

$$\begin{aligned} H^{2n-3+s}(\Gamma_{n,s}) \otimes_{\mathfrak{S}_s} V^{\wedge s} &\cong H^{2n-3}(\text{Out}(F_n); H^s(F_n^s)) \otimes_{\mathfrak{S}_s} V^{\wedge s} \quad (\text{Prop. 3.6}) \\ &\cong H^{2n-3}(\text{Out}(F_n); H^{\wedge s}) \otimes_{\mathfrak{S}_s} V^{\wedge s} \\ &\cong H^{2n-3}(\text{Out}(F_n); H^{\wedge s} \otimes_{\mathfrak{S}_s} V^{\wedge s}) \\ &\cong H^{2n-3}(\text{Out}(F_n); H^{\otimes s} \otimes_{\mathfrak{S}_s} V^{\otimes s}) \\ &\cong H^{2n-3}(\text{Out}(F_n); \text{Sym}^s(H \otimes V)). \end{aligned} \quad \square$$

Let $\mathcal{H}_V^{n,s}$ denote the hairy Lie graph complex (see [10, 11]), where graphs have rank n and s hairs labeled by vectors from V . In [10] the following theorem, with the twist accidentally omitted, was proved by a direct analysis of the chain complex.

Theorem 9.2 ([10], Theorem 11.1). *There is an isomorphism*

$$H_k(\mathcal{H}_V^{n,s}) \cong H^{2n+s-2-k}(\Gamma_{n,s}) \otimes_{\mathfrak{S}_s} V^{\wedge s}.$$

Combining this with Lemma 9.1 gives a shorter proof of the following theorem from [11] relating the first homology of the hairy Lie graph complex with the cohomology of $\text{Out}(F_n)$ with twisted coefficients.

Theorem 9.3 ([10], Theorem 8.8). *For $n \geq 2, s \geq 0$ there is an isomorphism*

$$H_1(\mathcal{H}_V^{n,s}) \cong H^{2n-3}(\text{Out}(F_n); \text{Sym}^s(\mathbb{H} \otimes V)).$$

This theorem has been slightly restated here to be more compatible with current notation.

9.2. The Lie algebra of symplectic derivations. According to Theorem 9.2 hairy graph homology is obtained by twisting the homology of $\Gamma_{n,s}$ with $V^{\wedge s}$. In [10, 11], it was shown that if the dimension of V is sufficiently large, the k -th homology of the Lie algebra \mathfrak{h}_V of positive degree symplectic derivations embeds in hairy Lie graph homology:

$$H_k(\mathfrak{h}_V) \subset H_k(\mathcal{H}_V) = \bigoplus_{n,s} H_k(\mathcal{H}_V^{n,s}) \cong \bigoplus_{n,s} H^{2n+s-2-k}(\Gamma_{n,s}) \otimes_{\mathfrak{S}_s} V^{\wedge s}.$$

Furthermore, every irreducible $\text{GL}(V)$ -module $\mathbb{S}_\lambda V$ in the decomposition of hairy graph homology corresponds to an irreducible $\text{Sp}(V)$ -module $\mathbb{S}_{\langle \lambda \rangle} V$ in the decomposition of $H_*(\mathfrak{h}_V)$. Thus the cohomology classes found in this paper produce homology classes for \mathfrak{h}_V . These classes can be used to show that $H_k(\mathfrak{h})$ contains infinitely many different Sp -modules, as we now show.

Theorem 9.4. *$H_{3n+d-2}(\mathfrak{h})$ contains the Sp -module $\mathbb{S}_{\langle ((2m+1)^n, 1^d) \rangle}$ for all m .*

Proof. Let $s = 2n(m+1) + d$. By Theorem 3.14, $H^{2nm}(\Gamma_{n,s})$ contains the \mathfrak{S}_s -submodule $P_{(n+d, n^{2m})}$ with multiplicity 1. Setting $k = 3n + d - 2$ we have

$$\begin{aligned} H_{3n+d-2}(\mathcal{H}_V^{n,s}) &\cong H^{2nm}(\Gamma_{n,s}) \otimes_{\mathfrak{S}_s} V^{\wedge s} \\ &\supset P_{(n+d, n^{2m})} \otimes_{\mathfrak{S}_s} V^{\wedge s} \end{aligned}$$

Since $V^{\wedge s} = \bigoplus_\lambda P_\lambda \otimes \mathbb{S}_\lambda(V)$ and $P_\lambda \otimes_{\mathfrak{S}_s} P_\mu = 0$ unless $\lambda = \mu$, in which case it is the trivial module \mathbb{k} , we see that for $\dim(V) \geq \max(2m+1, n+d)$, $H_{3n+d-2}(\mathcal{H}_V^{n,s})$ contains the $\text{GL}(V)$ -submodule $\mathbb{S}_{(n+d, n^{2m})'}(V) = \mathbb{S}_{\langle ((2m+1)^n, 1^d) \rangle}(V)$. Therefore for $\dim(V)$ sufficiently large $H_{3n+d-2}(\mathfrak{h}_V)$ contains the corresponding $\text{Sp}(V)$ -submodule $\mathbb{S}_{\langle ((2m+1)^n, 1^d) \rangle}(V)$. Taking the limit as $\dim(V)$ goes to infinity, gives that $H_k(\mathfrak{h})$ contains $\mathbb{S}_{\langle ((2m+1)^n, 1^d) \rangle}$ for all m . \square

Remark 9.5. In [12] the original Morita classes in the cohomology of $\text{Out}(F_n)$ were re-interpreted in terms of hairy graphs and this point of view was then used to construct more classes, called generalized Morita classes. The fact that the classes described in Sections 5.1 and 5.5 represent the same classes stems from the identification

$$H_1(\mathcal{H}_V^{1,2k+1}) \cong H^{2k}(\Gamma_{1,2k+1}) \otimes_{\mathfrak{S}_{2k+1}} V^{\wedge 2k+1} = \mathbb{S}_{2k+1} V = \text{Sym}^{2k+1} V$$

from Theorem 9.2. A generator of $H_1(\mathcal{H}_V^{1,2k+1})$ is a linear Lie tree with two ends joined by an edge and $2k + 1$ commuting V -labeled leaves (“hairs”), whereas a

generator of $H^{2k}(\Gamma_{1,2k+1})$ is the same thing, but with unlabeled leaves. Call this generator α_k^* , as it is dual to α_k . The graphical cocycle of [12] is nonzero only if the graph is the union of $\alpha_{k_1}^*, \dots, \alpha_{k_m}^*$ with the hairs connected up by edges. In that case the cocycle evaluates to \pm the graph obtained by shrinking each α_k to a point. Further projecting the graphical cocycle into the subspace spanned by a single graph gives the Morita cocycle μ_G . By construction, this process is dual to the gluing map defined by G .

Remark 9.6. In [11] it was explained that hairy graph homology can be viewed as the Feynman transform of a cyclic operad. In light of Theorem 9.2 this implies that the cohomology groups $H^*(\Gamma_{n,s})$ can be combined into a (twisted) modular co-operad [18]. Therefore the duals $H_*(\Gamma_{n,s})$ of these groups form a (twisted) modular operad. The assembly maps defined in Section 5 are the structure maps of this modular operad.

10. Open Questions

We finish with several questions and conjectures related to results in the paper. The conjectures have been verified for all but the most recently discovered nontrivial classes in $H_k(\Gamma_{n,s})$, but there are not enough of these verified cases to provide overwhelming evidence for the conjectures.

In what follows we always exclude trivial assembly maps, those that involve a component graph $X_{0,2}$ where the gluing involves only one leaf, because the map $\beta : H_k(\Gamma_{n,s}) \rightarrow H_k(\Gamma_{n,s})$ induced by the assembly map $\alpha : H_k(\Gamma_{n,s}) \otimes H_0(\Gamma_{0,2}) \rightarrow H_k(\Gamma_{n,s})$ is equal to the identity.

10.1. Surjectivity of the assembly maps below the virtual cohomological dimension. An assembly map $H_*(\Gamma_{n_1,s_1}) \otimes \dots \otimes H_*(\Gamma_{n_k,s_k}) \rightarrow H_*(\Gamma_{n,s})$ for $k \geq 2$ can never have nontrivial image in the virtual cohomological dimension (vcd) of $\Gamma_{n,s}$ since in that case the vcd of $\Gamma_{n,s}$ is strictly greater than the sum of the vcd's of the factors. An assembly map induced by gluing pairs of leaves of a single graph preserves the vcd, so such an assembly map might conceivably be nontrivial in that dimension, though it seems unlikely that such a map can be surjective (with nontrivial image). By contrast, in other dimensions we expect that all classes are constructed by assembly from lower-rank graphs.

Conjecture 10.1. *Suppose that $k \neq 2n + s - 3$, the vcd of $\Gamma_{n,s}$. Then $H_k(\Gamma_{n,s})$ is generated by the images of the assembly maps A_ϕ over all possible gluings which give the graph $X_{n,s}$.*

The examples in Section 5 confirm this conjecture for $n = 1$ and $n = 2$. It can be seen from the description of the cohomology groups that $H_k(\Gamma_{n,s+1})$ is generated by the images of the assembly maps $H_k(\Gamma_{n,s}) \otimes H_0(\Gamma_{0,3}) \rightarrow H_k(\Gamma_{n,s+1})$ in all cases

except when $k = 2n + s - 3$, or when $n = 2$ and $k = s$ is divisible by 4. The first exception is excluded from the conjecture since these classes are in the vcd and the second one is covered by Section 5.4.

The first three Morita classes and the first two Eisenstein classes are obtained as images of assembly maps. By Bartholdi's calculations there are nontrivial classes in $H_{11}(\text{Out}(F_7))$ and $H_8(\text{Out}(F_7))$. The first of these is in the vcd , and there is a natural candidate for the second one, obtained by assembling four copies of $\alpha_1 \in H_2(\Gamma_{1,3})$ in a tetrahedral pattern. By Theorem 2.4 the class in $H_{11}(\text{Out}(F_7))$ also produces a class in $H_{11}(\text{Aut}(F_7))$, which cannot be the Eisenstein class by Remark 5.6. It is possible that this class can be obtained by assembling classes in $H_{11}(\Gamma_{6,2})$ and $H_0(\Gamma_{0,3})$.

As mentioned in Remark 9.6, the homology groups $H_*(\Gamma_{n,s})$ for $n \geq 1$ form a twisted modular operad. Conjecture 10.1 together with computations in Section 5 imply that this modular operad is generated by $H_0(\Gamma_{0,3})$ and $H_{2n+s-3}(\Gamma_{n,s})$.

10.2. Injectivity of classes constructed from modular forms. In Section 5.8 we constructed maps from spaces of modular forms to $H_{4m+2}(\text{Out}(F_{2m+3}))$ by gluing two rank 2 graphs together along all of their leaves. These maps take the form

$$\chi : \bigoplus_{0 \leq i < m} \left(\bigwedge^2 \mathcal{X}_{2m,i} \right) \longrightarrow H_{4m+2}(\text{Out}(F_{2m+3}))$$

The first of these maps which could be nontrivial has target $H_{42}(\text{Out}(F_{23}))$. We do not know how to show that the image is nontrivial and it is beyond the reach of computer calculations. Nevertheless, we make the following conjecture:

Conjecture 10.2. *The restriction of χ to the term $\bigwedge^2 \mathcal{X}_{2m,0}$ is injective.*

This conjecture appeared as a question in [10]. It seems unlikely that χ is injective on $\bigwedge^2 \mathcal{X}_{2m,i}$ for i close to m , but injectivity is still plausible for small values of i , so we ask the following question:

Question 10.3. *For which i is the restriction of χ to the term $\bigwedge^2 \mathcal{X}_{2m,i}$ injective?*

We remark that Conjecture 10.2 would contradict a conjecture made by Church, Farb, and Putman [8, Conjecture 12] concerning a stability property of the groups $H_{2n-3-k}(\text{Out}(F_n))$ for fixed k as $n \rightarrow \infty$.

10.3. Odd-dimensional classes. In Section 5.7 we constructed maps from the space \mathcal{M}_{2k+2} of modular forms of weight $2k + 2$ to $H_{4k+3}(\text{Out}(F_{s+2}))$ for all k and s . For small s this map is trivial because the homology dimension is above the vcd and for large s the map must be trivial by homology stability, but for $s = 2k + 2$ we conjecture that it is highly nontrivial:

Conjecture 10.4. *The map $\mathcal{M}_{2k+2} \rightarrow H_{4k+3}(\text{Out}(F_{2k+4}))$ constructed in Section 5.7 is injective for all k .*

This too is incompatible with the conjecture of Church, Farb, and Putman. Since the vcd of $\text{Out}(F_{2k+4})$ is $4k + 5$, these classes lie in codimension 2, and Conjecture 10.4 implies that the dimension of $H_{\text{vcd}-2}(\text{Out}(F_n))$ grows at least linearly with n when n is even.

10.4. Vanishing under stabilization. For $s > 0$ there are two natural ways to stabilize $\Gamma_{n,s}$, by increasing n or by increasing s , as described in Section 6. Here we consider the stabilization increasing n .

Conjecture 10.5. *If ϕ is any gluing, then all positive-dimensional classes in the image of the assembly map A_ϕ vanish after a single stabilization with respect to n .*

The condition of positive dimension is of course necessary to exclude the classes in $H_0(\Gamma_{n,s})$ which clearly do not vanish after stabilization. In Section 6 we showed that the conjecture is true for classes in the image of an assembly map where stabilization is done using a rank 1 factor.

We point out that Conjectures 10.1 and 10.5 imply the following for classes in dimensions below the vcd :

Conjecture 10.6. *All classes in $H_k(\Gamma_{n,s})$ for $0 < k < 2n + s - 3$ vanish after a single stabilization with respect to n .*

Since the vcd of $\Gamma_{n,s}$ increases with n , this conjecture implies that two stabilizations kill *all* homology classes.

11. Tables of results

In this section we write out tables of the cohomology of $\Gamma_{n,s}$ for small values of s . The space above the diagonal in each table is left blank since it represents terms above the virtual cohomological dimension, where the cohomology must vanish.

| | H^0 | H^1 | H^2 | H^3 | H^4 | H^5 | H^6 | H^7 |
|----------------|--------------|-------|---------------|-------|---------------|-------|---------------|-------|
| $\Gamma_{1,0}$ | \mathbb{k} | | | | | | | |
| $\Gamma_{1,1}$ | $P_{(1)}$ | | | | | | | |
| $\Gamma_{1,2}$ | $P_{(2)}$ | 0 | | | | | | |
| $\Gamma_{1,3}$ | $P_{(3)}$ | 0 | $P_{(1^3)}$ | | | | | |
| $\Gamma_{1,4}$ | $P_{(4)}$ | 0 | $P_{(2,1^2)}$ | 0 | | | | |
| $\Gamma_{1,5}$ | $P_{(5)}$ | 0 | $P_{(3,1^2)}$ | 0 | $P_{(1^5)}$ | | | |
| $\Gamma_{1,6}$ | $P_{(6)}$ | 0 | $P_{(4,1^2)}$ | 0 | $P_{(2,1^4)}$ | 0 | | |
| $\Gamma_{1,7}$ | $P_{(7)}$ | 0 | $P_{(5,1^2)}$ | 0 | $P_{(3,1^4)}$ | 0 | $P_{(1^7)}$ | |
| $\Gamma_{1,8}$ | $P_{(8)}$ | 0 | $P_{(6,1^2)}$ | 0 | $P_{(4,1^4)}$ | 0 | $P_{(2,1^6)}$ | 0 |

Table 1. $H^i(\Gamma_{1,s}; \mathbb{k})$ for $s \leq 8$

| | H^0 | H^1 | H^2 | H^3 | H^4 | H^5 | H^6 | H^7 |
|----------------|-------|-------|-------|-------|-------|-------|-------|-------|
| $\Gamma_{1,0}$ | 1 | | | | | | | |
| $\Gamma_{1,1}$ | 1 | | | | | | | |
| $\Gamma_{1,2}$ | 1 | 0 | | | | | | |
| $\Gamma_{1,3}$ | 1 | 0 | 1 | | | | | |
| $\Gamma_{1,4}$ | 1 | 0 | 3 | 0 | | | | |
| $\Gamma_{1,5}$ | 1 | 0 | 6 | 0 | 1 | | | |
| $\Gamma_{1,6}$ | 1 | 0 | 10 | 0 | 5 | 0 | | |
| $\Gamma_{1,7}$ | 1 | 0 | 15 | 0 | 15 | 0 | 1 | |
| $\Gamma_{1,8}$ | 1 | 0 | 21 | 0 | 35 | 0 | 7 | 0 |

Table 2. Dimensions of $H^i(\Gamma_{1,s}; \mathbb{k})$ for $s \leq 8$

| | H^0 | H^1 | H^2 | H^3 | H^4 | H^5 | H^6 | H^7 | H^8 | H^9 | H^{10} |
|-----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----------|
| $\Gamma_{2,0}$ | 1 | 0 | | | | | | | | | |
| $\Gamma_{2,1}$ | 1 | 0 | 0 | | | | | | | | |
| $\Gamma_{2,2}$ | 1 | 0 | 0 | 0 | | | | | | | |
| $\Gamma_{2,3}$ | 1 | 0 | 0 | 0 | 0 | | | | | | |
| $\Gamma_{2,4}$ | 1 | 0 | 0 | 0 | 2 | 3 | | | | | |
| $\Gamma_{2,5}$ | 1 | 0 | 0 | 0 | 10 | 15 | 0 | | | | |
| $\Gamma_{2,6}$ | 1 | 0 | 0 | 0 | 30 | 45 | 0 | 5 | | | |
| $\Gamma_{2,7}$ | 1 | 0 | 0 | 0 | 70 | 105 | 0 | 35 | 0 | | |
| $\Gamma_{2,8}$ | 1 | 0 | 0 | 0 | 140 | 210 | 0 | 140 | 14 | 35 | |
| $\Gamma_{2,9}$ | 1 | 0 | 0 | 0 | 252 | 378 | 0 | 420 | 126 | 315 | 0 |
| $\Gamma_{2,10}$ | 1 | 0 | 0 | 0 | 420 | 630 | 0 | 1050 | 630 | 1575 | 0 |

Table 4. Dimensions of $H^i(\Gamma_{2,s}; \mathbb{k})$ for $i, s \leq 10$

| | H^0 | H^1 | H^2 | H^3 | H^4 | H^5 | H^6 | H^7 | H^8 | H^9 | H^{10} |
|-----------------|--|-------|-------|-------|--|---|-------|-------|--|-------|----------|
| $\Gamma_{2,0}$ | \mathbb{k} | 0 | | | | | | | | | |
| $\Gamma_{2,1}$ | $P_{(1)}$ | 0 | 0 | | | | | | | | |
| $\Gamma_{2,2}$ | $P_{(2)}$ | 0 | 0 | 0 | | | | | | | |
| $\Gamma_{2,3}$ | $P_{(3)}$ | 0 | 0 | 0 | 0 | | | | | | |
| $\Gamma_{2,4}$ | $P_{(4)}$ | 0 | 0 | 0 | 0 | $P_{(2^2)}$ | | | $P_{(2,1^2)}$ | | |
| $\Gamma_{2,5}$ | $P_{(5)}$ | 0 | 0 | 0 | 0 | $P_{(3,2)} \oplus P_{(2^2,1)}$ | | | $P_{(3,1^2)} \oplus P_{(2^2,1)} \oplus P_{(2,1^3)}$ | | |
| $\Gamma_{2,6}$ | $P_{(6)}$ | 0 | 0 | 0 | 0 | $P_{(4,2)} \oplus P_{(3,2,1)} \oplus P_{(2^3)}$ | | | $P_{(4,1^2)} \oplus P_{(3,2,1)} \oplus P_{(3,1^3)} \oplus P_{(2^2,1^2)}$ | | |
| $\Gamma_{2,7}$ | $P_{(7)}$ | 0 | 0 | 0 | 0 | $P_{(5,2)} \oplus P_{(4,2,1)} \oplus P_{(3,2^2)}$ | | | $P_{(5,1^2)} \oplus P_{(4,2,1)} \oplus P_{(4,1^3)} \oplus P_{(3,2,1^2)}$ | | |
| $\Gamma_{2,8}$ | $P_{(8)}$ | 0 | 0 | 0 | 0 | $P_{(6,2)} \oplus P_{(5,2,1)} \oplus P_{(4,2^2)}$ | | | $P_{(6,1^2)} \oplus P_{(5,2,1)} \oplus P_{(5,1^3)} \oplus P_{(4,2,1^2)}$ | | |
| $\Gamma_{2,9}$ | $P_{(9)}$ | 0 | 0 | 0 | 0 | $P_{(7,2)} \oplus P_{(6,2,1)} \oplus P_{(5,2^2)}$ | | | $P_{(7,1^2)} \oplus P_{(6,2,1)} \oplus P_{(6,1^3)} \oplus P_{(5,2,1^2)}$ | | |
| $\Gamma_{2,10}$ | $P_{(10)}$ | 0 | 0 | 0 | 0 | $P_{(8,2)} \oplus P_{(7,2,1)} \oplus P_{(6,2^2)}$ | | | $P_{(8,1^2)} \oplus P_{(7,2,1)} \oplus P_{(7,1^3)} \oplus P_{(6,2,1^2)}$ | | |
| | H^0 | H^1 | H^2 | H^3 | H^4 | H^5 | H^6 | H^7 | H^8 | H^9 | H^{10} |
| $\Gamma_{2,5}$ | 0 | | | | | | | | | | |
| $\Gamma_{2,6}$ | 0 | | | | $P_{(2,1^4)}$ | | | | | | |
| $\Gamma_{2,7}$ | 0 | | | | $P_{(3,1^4)} \oplus P_{(2^2,1^3)} \oplus P_{(2,1^5)}$ | 0 | | | | | |
| $\Gamma_{2,8}$ | $P_{(4,1^4)} \oplus P_{(3,2,1^3)} \oplus P_{(3,1^5)} \oplus P_{(2^2,1^4)}$ | | | | $P_{(2^4)}$ | $P_{(2,1^6)} \oplus P_{(2^3,1^2)}$ | | | | | |
| $\Gamma_{2,9}$ | 0 | | | | $P_{(5,1^4)} \oplus P_{(4,2,1^3)} \oplus P_{(4,1^5)} \oplus P_{(3,2,1^4)}$ | $P_{(3,2^3)} \oplus P_{(2^4,1)}$ | | | $P_{(3,1^6)} \oplus P_{(2^2,1^5)} \oplus P_{(2,1^7)} \oplus P_{(3,2^2,1^2)} \oplus P_{(2^4,1)} \oplus P_{(2^3,1^3)}$ | | 0 |
| $\Gamma_{2,10}$ | 0 | | | | $P_{(6,1^4)} \oplus P_{(5,2,1^3)} \oplus P_{(5,1^5)} \oplus P_{(4,2,1^4)}$ | $P_{(4,2^3)} \oplus P_{(3,2^3,1)} \oplus P_{(2^5)}$ | | | $P_{(4,1^6)} \oplus P_{(3,2,1^5)} \oplus P_{(3,1^7)} \oplus P_{(2^2,1^6)} \oplus P_{(4,2^2,1^2)} \oplus P_{(2^4,1^2)} \oplus P_{(3,2^2,1^3)} \oplus P_{(3,2^3,1)}$ | | 0 |

Table 3. $H^i(\Gamma_{2,s}; \mathbb{k})$ for $i, s \leq 10$

References

- [1] L. Bartholdi, The rational homology of the outer automorphism group of F_7 , *New York J. Math.*, **22** (2016), 191–197. MR 3484680 Zbl 06553443
- [2] M. Bestvina and M. Feighn, The topology at infinity of $\text{Out}(F_n)$, *Invent. Math.*, **140** (2000), no. 3, 651–692. MR 1760754 Zbl 0954.55011
- [3] J. Birman, Mapping class groups and their relationship to braid groups, *Comm. Pure Appl. Math.*, **22** (1969), 213–238. MR 0243519 Zbl 0167.21503
- [4] M. R. Bridson and K. Vogtmann, Abelian covers of graphs and maps between outer automorphisms of free groups, *Math. Ann.*, **353** (2012), no. 4, 1069–1102. MR 2944024 Zbl 1261.20035
- [5] M. Chan, Topology of the tropical moduli spaces $M_{2,n}$, 2015. arXiv:1507.03878
- [6] M. Chan, S. Galatius and S. Payne, The tropicalization of the moduli space of curves II: Topology and applications, 2016. arXiv:1604.03176
- [7] T. Church and B. Farb, Representation theory and homological stability, *Adv. in Math.*, **245** (2013), 250–314. MR 3084430 Zbl 1300.20051
- [8] T. Church, B. Farb and A. Putman, A stability conjecture for the unstable cohomology of $SL_n\mathbb{Z}$, mapping class groups, and $\text{Aut}(F_n)$, in *Algebraic topology: applications and new directions*, 55–70, Contemp. Math., 620, Amer. Math. Soc., Providence, RI, 2014. MR 3290086 Zbl 06520459
- [9] T. Church, B. Farb and A. Putman, The rational cohomology of the mapping class group vanishes in its virtual cohomological dimension, *Int. Math. Res. Not.*, **2012** (2012), no. 21, 5025–5030. MR 2993444 Zbl 1260.57033
- [10] J. Conant, M. Kassabov and K. Vogtmann, Hairy graphs and the unstable homology of $\text{Mod}(g, s)$, $\text{Out}(F_n)$, and $\text{Aut}(F_n)$, *Journal of Topology*, **6** (2013), no. 1, 119–153. MR 3029423 Zbl 06272197
- [11] J. Conant, M. Kassabov and K. Vogtmann, Higher hairy graph homology, to appear in *Geometriae Dedicata*. MR 3347586 Zbl 06446920
- [12] J. Conant and K. Vogtmann, Morita classes in the homology of automorphism groups of free groups, *Geom. Topol.*, **8** (2004), 1471–1499 (electronic). MR 2119302 Zbl 1121.20036
- [13] J. Conant and K. Vogtmann, Morita classes in the homology of $\text{Aut}(F_n)$ vanish after one stabilization, *Groups, Geometry, and Dynamics*, **2** (2008), no. 1, 121–138. MR 2367210 Zbl 1148.20036
- [14] M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups, *Inventiones*, **84** (1986), 91–119. MR 0830040 Zbl 0589.20022

- [15] W. Fulton and J. Harris, *Representation theory. A first course*, Graduate Texts in Mathematics, 129, Readings in Mathematics, Springer-Verlag, New York, 1991. MR 1153249 Zbl 0744.22001
- [16] S. Galatius, Stable homology of automorphism groups of free groups, *Ann. of Math.*, **173** (2011), 705–768. MR 2784914 Zbl 1268.20057
- [17] F. Gerlitz, Ph. D. thesis, Cornell University, 2002.
- [18] E. Getzler and M. Kapranov, Modular operads, *Compositio Math.*, **110** (1998), no. 1, 65–126. MR 1601666 Zbl 0894.18005
- [19] J. Gray, *On the homology of automorphism groups of free groups*, Ph. D thesis, University of Tennessee, 2011.
- [20] K. Haberland, Perioden von Modulformen einer Variabler und Gruppencohomologie. I, [Periods of modular forms of one variable and group cohomology. I], *Math. Nachr.*, **112** (1983), 245–282. MR 0726861 Zbl 0526.10024
- [21] A. Hatcher, *Algebraic Topology*, Cambridge University Press, Cambridge, 2002. MR 1867354 Zbl 1044.55001
- [22] A. Hatcher, Homological stability for automorphism groups of free groups, *Comment. Math. Helv.*, **70** (1995), no. 1, 39–62. MR 1314940 Zbl 0836.57003
- [23] A. Hatcher and K. Vogtmann, Rational homology of $\text{Aut}(F_n)$, *Math. Res. Let.*, **5** (1998), 759–780. MR 1671188 Zbl 1040.20042
- [24] A. Hatcher and K. Vogtmann, Homology stability for outer automorphism groups of free groups, *Algebr. Geom. Topol.*, **4** (2004), 1253–1272; Erratum (with Nathalie Wahl), **6** (2006), 573–579. MR 2113904 Zbl 1093.20020
- [25] A. Hatcher and N. Wahl, Stabilization for mapping class groups of 3-manifolds, *Duke Math. J.*, **155** (2010), 205–269. MR 2736166 Zbl 1223.57004
- [26] R. Jiménez Rolland, On the cohomology of pure mapping class groups as FI-modules, *J. Homotopy Relat. Struct.*, **10** (2015), no. 3, 401–424. MR 3385692 Zbl 06483843
- [27] N. Kawazumi, Cohomological aspects of Magnus expansions, 2006. arXiv:math/0505497v3
- [28] M. Kontsevich, Formal (non) commutative symplectic geometry, *The Gelfand Mathematical Seminars, 1990–1992*, 173–187. MR 1247289 Zbl 0821.58018
- [29] M. Kontsevich, Feynman diagrams and low-dimensional topology, *First European Congress of Mathematics*, vol. 2, 97–121, 1992. MR 1341841 Zbl 0872.57001
- [30] N. H. Kuiper, The quotient space of $CP(2)$ by complex conjugation is the 4-sphere, *Math. Ann.*, **208** (1974), 175–177. MR 0346817 Zbl 0265.52002

- [31] S. Lang, *Introduction to modular forms*, With appendixes by D. Zagier and Walter Feit. Corrected reprint of the 1976 original, Grundlehren der Mathematischen Wissenschaften, 222, Springer-Verlag, Berlin, 1995. MR 0429740 Zbl 0344.10011
- [32] W. S. Massey, The quotient space of the complex projective plane under conjugation is a 4-sphere, *Geometriae Dedicata*, **2** (1973), 371–374. MR 0341511 Zbl 0273.57019
- [33] S. Morita, Structure of the mapping class groups of surfaces: a survey and a prospect, in *Proceedings of the Kirbyfest*, 349–406, Geom. Topol. Monogr., 2, Geom. Topol. Publ., Coventry, 1999 (electronic). MR 1734418 Zbl 0959.57018
- [34] S. Morita, T. Sakasai and M. Suzuki, Abelianizations of derivation Lie algebras of the free associative algebra and the free Lie algebra, *Duke Math. J.*, **162** (2013), 965–1002. MR 3047471 Zbl 1308.17021
- [35] S. Morita, T. Sakasai and M. Suzuki, Computations in formal symplectic geometry and characteristic classes of moduli spaces, *Quantum Topol.*, **6** (2015), no. 1, 139–182. MR 3335007 Zbl 06436533
- [36] S. Morita, T. Sakasai and M. Suzuki, Integral Euler characteristic of $\text{Out}(F_{11})$, 2014. arXiv:1405.4063
- [37] R. Ohashi, The rational homology group of $\text{Out}(F_n)$ for $n \leq 6$, *Experiment. Math.*, **17** (2008), 167–179. MR 2433883 Zbl 1191.20057
- [38] A. Saied, On the FI-module structure of $H_i(\Gamma_{n,s})$, 2016. arXiv:1506.05861

Received September 7, 2015

J. Conant, Department of Mathematics, University of Tennessee,
Knoxville, TN 37996, USA

E-mail: jim.conant@gmail.com

A. Hatcher, Mathematics Department, Malott Hall, Cornell University,
Ithaca, NY 14853, USA

E-mail: hatcher@math.cornell.edu

M. Kassabov, Mathematics Department, Malott Hall, Cornell University,
Ithaca, NY 14853, USA

E-mail: kassabov@math.cornell.edu

K. Vogtmann, Mathematics Institute, Zeeman Building, University of Warwick,
Coventry CV4 7AL, UK

E-mail: k.vogtmann@warwick.ac.uk