

# Preface

Autor(en): **[s.n.]**

Objekttyp: **Preface**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **57 (1964)**

Heft 2

PDF erstellt am: **09.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

delle zone a Foraminiferi planctonici nella Tetide. Per ragioni storiche tutte le località-tipo dei piani del Paleogene si trovano in serie epicontinentali con faune di tipo boreale. Di conseguenza il loro valore e significato per quanto riguarda la stratigrafia del Terziario antico non dovrebbe essere sopravvalutato.

Vengono descritte tre serie, di cui due nei dintorni di Gubbio (Perugia) ed una presso Fossombrone (Pesaro).

### РЕЗЮМЕ

В данной работе изучены эволюция и стратиграфическое распространение некоторых Глобороталий в палеоцене и нижнем эоцене в формации Сгаля центральной части Апеннин (Италия).

Выбор ограничивается "коническими" Глобороталиями и видами, которые могут-быть рассматриваемы как их предки. 22 вида описаны и изображены.

Зональное расчленение, разработанное Болли (1957) в Тринидаде, может-быть с некоторыми незначительными изменениями применено также к формациям Сгаля.

Автор убежден, что стратиграфия палеогена будет основываться на зональной шкале по фауне планктонных фораминифер Средиземноморской области. По историческим причинам все стратотипические разрезы яруса палеогена лежат в эпиконтинентальных отложениях бореальной фаунистической области: Их значение для стратиграфии палеогена не должно быть поэтому преувеличено.

### ABSTRACT

The taxonomy and stratigraphic distribution of some *Globorotalia* from the Paleocene and Lower Eocene of the Central Apennines are studied. The systematic descriptions are restricted to the «conical» *Globorotalia* and to species which are thought to be their predecessors. 22 species are described and figured.

In the Scaglia of the Central Apennines, the same zonal succession as established by BOLLI (1957) in Trinidad is observed. A few minor modifications are due mainly to taxonomic reasons.

An attempt to correlate the planktonic foraminiferal zonation with the respective European stages is discussed.

### PREFACE

The present paper is a doctoral thesis submitted to the Faculty of Science of the University of Basel. It has been prepared at the Geological Institute of this University.

Investigations on Lower Tertiary *Globorotalia* were started in late summer 1960 and finished in spring 1964. For different reasons, they had to be interrupted several times.

Three sections within the Central Apennines have been measured and sampled during two excursions in autumn 1961 and spring 1962. For comparison, classical

localities of uppermost Cretaceous and Paleogene in the Netherlands, Belgium, Southern France, Northern Spain and Northern Italy have been visited. A visit at the Geological Institute of the Academy of Science of the Soviet Union in Moscow during August 1963 has been very instructive.

#### ACKNOWLEDGMENTS

I wish to express my sincere thanks to Professor Dr. M. REICHEL for suggesting the present thesis, and mainly for his invaluable help and advice during the investigations and the preparation of the manuscript and the illustrations. I am greatly indebted to him for introducing me to the methods and problems of micropaleontology.

Professor Dr. L. VONDERSCHMITT, Director of the Geological Institute of the University of Basel, is acknowledged for granting all kinds of facilities during the study at the Institute, for the interest he has taken in my work and the financial aid he granted me.

I want to thank Professor Dr. H. SCHAUB, Director of the Museum of Natural History, Basel, for introducing me to the stratigraphy of the Paleogene, for helpful suggestions and criticism, for giving me numerous samples and for his guidance during several excursions.

Dr. O. RENZ, at the Museum of Natural History, Basel, has kindly read and corrected the manuscript. It was a great honour for me to discuss many problems with him, who is the author of the fundamental work on the micropaleontology and stratigraphy of the Central Apennines.

I am greatly indebted to Dr. H. G. KUGLER, at the Museum of Natural History, Basel, for his stimulating interest and help in my studies, for much advice based on his vast experience and for having placed at my disposal his type-collection of Foraminifera from Trinidad.

I am pleased to thank Dr. M. B. CITA, Professor at the University of Milan, for discussing with me several problems concerning my investigations, for her aid in procuring literature not available in Basel and for having shared with me her vast experience in micropaleontological problems of the Scaglia.

Much is owed to Dr. I. PREMOLI SILVA, University of Milan, for her collaboration, for her guidance to numerous localities in Northern and Central Italy and for her help in obtaining literature and samples for comparison.

Professor Dr. W. W. HAY, University of Illinois, has contributed important samples from the Paleogene of the United States and Mexico and much information on stratigraphic problems related to them.

I want to express my gratitude to the Presidium of the Academy of Science of the Soviet Union for an invitation to the Geological Institute of the Academy, to Professor Dr. D. M. RAUZER-CHERNOUSOVA and to Professor Dr. V. V. MENNER for their generous help, and especially to Dr. V. A. KRASHENNINIKOV for his introduction to the Paleogene of Southern Soviet Union, for numerous samples and literature and for all the trouble he took to help me during my stay in Moscow.

Professor Dr. N. N. SUBBOTINA, VNIGRI Leningrad, and Dr. E. K. SHUTZKAYA, VNIGNI Moscow, have kindly discussed many taxonomic and stratigraphic prob-