

III. A NEW METHOD FOR NOT NECESSARILY MONOTONIC $f(x)$

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **11 (1965)**

Heft 2-3: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **02.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

and therefore, using (13)

$$\int_{\Psi(x_0)}^{\gamma\psi(b_v)} f(x) dx \leq \alpha \int_{\psi(x_0)}^{\psi(b_v)} f(x) dx = \alpha \int_{\psi(x_0)}^{\gamma\psi(b_v)} f(x) dx + \alpha \int_{\gamma\psi(b_v)}^{\psi(b_v)} f(x) dx .$$

But the last right hand integral is, by (14), $\leq c \log \frac{1}{\gamma}$, so that we obtain:

$$(1 - \alpha) \int_{\Psi(x_0)}^{\gamma\psi(b_v)} f(x) dx \leq \int_{\psi(x_0)}^{\Psi(x_0)} f(x) dx + c \log \frac{1}{\gamma} .$$

The convergence of (2) follows now immediately from $\psi(b_v) \rightarrow \infty$.

13. Suppose that we have, on the other hand, for an $a > x_0$:

$$\Psi(a) \leq \psi(a) .$$

Proceeding then as in the proof of the Theorem 1 we have, as from $\psi(b_v) \rightarrow \infty$ and the total continuity of $\psi(x)$ follows $b_v \rightarrow \infty$, for $b_v \geq a$:

$$\int_{\Psi(a)}^{\Psi(b_v)} f(x) dx \leq \alpha \int_{\psi(a)}^{\psi(b_v)} f(x) dx ,$$

and, for $v \rightarrow \infty$:

$$\int_{\Psi(a)}^{\infty} f(x) dx \leq \alpha \int_{\psi(a)}^{\infty} f(x) dx .$$

But here the left hand integral is > 0 , the right hand integral is majorized by it and the relation is impossible for $\alpha < 1$.³⁾

III. A NEW METHOD FOR NOT NECESSARILY MONOTONIC $f(x)$

14. THEOREM 4. Assume that $\Psi(x)$ is for $x \geq x_0$ a positive and monotonically increasing differentiable function for which

³⁾ Observe that in Ermakof's paper [1] the criteria are given in the following form:
 $\sum_{\infty} f(v)$ for a monotonic $f(x)$ is convergent or divergent according as

$$\lim_{x \rightarrow \infty} \frac{f(\Psi(x))\Psi'(x)}{f(\psi(x))\psi'(x)}$$

is < 1 or > 1 . In the note [2] Ermakof takes $\Psi(x) \equiv x$ which is no essential specialisation. However, the conditions (5) for convergence and (9) for divergence (with the specialisation $\Psi(x) \equiv x$) are already found in the textbooks, see e.g. [3].

$\Psi'(x)$ is also monotonically increasing and that we have:

$$\Psi(x) > x \quad (x \geq x_0). \quad (16)$$

Suppose further that $f(x)$ is > 0 for $x \geq x_0$ and integrable and bounded from below by a positive number in any finite subinterval of $\langle x_0, \infty \rangle$. If we have for all $x \geq x_0$:

$$f(\Psi(x)) \Psi'(x) \geq f(x), \quad (17)$$

the sum

$$\sum_{v \geq x_0} f(v) \quad (18)$$

is divergent.

15. *Proof.* Introduce the function

$$F(x) = \inf_{x_0 \leq u \leq x} f(u); \quad (19)$$

then $F(x)$ is monotonically decreasing and we have for each $x \geq x_0$:

$$F(x) = \lim_{\kappa \rightarrow \infty} f(u_\kappa)$$

for a convenient sequence u_κ from the interval $\langle x_0, x \rangle$.

We can write therefore for a certain sequence v_κ from the interval $\langle x_0, x \rangle$:

$$F(\Psi(x)) \Psi'(x) = \lim_{\kappa \rightarrow \infty} f(\Psi(v_\kappa)) \Psi'(x) \geq \overline{\lim}_{\kappa \rightarrow \infty} f(\Psi(v_\kappa)) \Psi'(v_\kappa).$$

This is, however, by (17) $\geq \overline{\lim}_{\kappa \rightarrow \infty} f(v_\kappa) \geq F(x)$.

It follows

$$F(\Psi(x)) \Psi'(x) \geq F(x),$$

so that the integral $\int_{x_0}^{\infty} F(x) dx$ is divergent. Since $F(x)$ is monotonic, the same follows for the series $\sum_{v \geq x_0} F(v)$ which has (18) as a majorant. The Theorem 4 is proved.

16. THEOREM 5. Assume that $\Psi(x)$ is for $x \geq x_0$ a positive and monotonically increasing differentiable function for which (16) holds. Assume further that $\Psi'(x)$ is either, from a certain x on, monotonically increasing or, for $x \rightarrow \infty$, convergent to a finite

limit ω . Assume finally that $f(x)$ is ≥ 0 for $x \geq x_0$, measurable and bounded in each interval $x_0 \leq x \leq a$ and satisfies for all $x \geq x_0$ and for a certain constant $\delta < 1$ the inequality:

$$f(\Psi(x)) \Psi'(x) \leq \delta f(x) \quad (x \geq x_0). \quad (20)$$

Then the series (18) is convergent.

17. *Proof.* Take a number β with $1 > \beta > \delta$. Observe that $\Psi'(x)$ certainly cannot have for $x \rightarrow \infty$ a limit $\omega < 1$. For otherwise we would have, with $x \rightarrow \infty$,

$$(\Psi(x) - x)' \rightarrow \omega - 1 < 0, \quad \Psi(x) - x \rightarrow -\infty$$

contrary to (16).

We have therefore in any case, from a certain x on, $\Psi'(x) \geq \delta$, and, by (20), $f(\Psi(x)) \leq f(x)$. We can therefore assume, changing x_0 if necessary, that we have:

$$f(\Psi(x)) \leq f(x) \quad (x \geq x_0). \quad (21)$$

Further, if we have $\Psi(x) \rightarrow \omega \geq 1$ and if ω is finite there certainly exists an x_1 such that we have, if $x \geq x_1$, $y \geq x_1$,

$$\frac{\delta}{\beta} \leq \frac{\Psi'(x)}{\Psi'(y)} \leq \frac{\beta}{\delta}.$$

We can therefore assume, increasing x_0 if necessary, that we have:

$$\Psi'(x) \leq \frac{\beta}{\delta} \Psi'(y) \quad (y \geq x \geq x_0), \quad (22)$$

and this is obviously also true if $\Psi'(x)$ is monotonically increasing, so that we can now assume (22) as being true under the conditions of our Theorem.

18. Put

$$x_0 = a_0, \quad \Psi(a_0) = a_1, \dots, \Psi(a_v) = a_{v+1}, \dots$$

The sequence a_v is monotonically increasing. If $\lim a_v = \tau$ were finite, we would have $\Psi(\tau) = \tau$, contrary to (16). Therefore we have $a_v \uparrow \infty$.

We have therefore for any $x \geq x_0$ an index ν such that $a_\nu \leq x < a_{\nu+1}$.

Denoting by c an upper bound for $f(x)$ in the interval $\langle a_0, a_1 \rangle$ it follows then from (21):

$$f(x) \leq c \quad (x \geq x_0).$$

19. Put

$$G(x) = \sup_{u \geq x} f(u). \quad (23)$$

$G(x)$ is finite and monotonically decreasing and we have:

$$f(x) \leq G(x) \quad (x \geq x_0). \quad (24)$$

By (23), there exists for any $x \geq x_0$ a sequence of numbers u_κ , $u_\kappa \geq x$ such that $G(\Psi(x)) = \lim_{\kappa \rightarrow \infty} f(\Psi(u_\kappa))$ and by (22)

$$G(\Psi(x)) \Psi'(x) = \lim_{\kappa \rightarrow \infty} f(\Psi(u_\kappa)) \Psi'(x) \leq \overline{\lim}_{\kappa \rightarrow \infty} f(\Psi(u_\kappa)) \frac{\beta}{\delta} \Psi'(u_\kappa).$$

But this is, by (20),

$$\leq \frac{\beta}{\delta} \delta \overline{\lim}_{\kappa \rightarrow \infty} f(u_\kappa) \leq \beta G(x).$$

20. We have therefore

$$G(\Psi(x)) \Psi'(x) \leq \beta G(x),$$

so that $\int^\infty G(x) dx$ is convergent. But then, since $G(x)$ is monotonically decreasing, the series $\sum_{\nu} G(\nu)$ is convergent too, and, by (24), the same holds for the series (18). The Theorem 5 is proved.

21. THEOREM 6. Assume that $\Psi(x)$ is for $x \geq x_0$ a positive and monotonically increasing differentiable function for which we have (16). Suppose further that $f(x)$ is > 0 for $x \geq x_0$, is integrable and bounded from below by a positive number in any finite subinterval of $\langle x_0, \infty \rangle$ and satisfies for a constant $\beta > 1$ and for all $x \geq x_0$ the condition

$$f(\Psi(x)) \Psi'(x) \geq \beta f(x), \quad x \geq x_0. \quad (25)$$

Finally assume that there exists an $x_1 \geq x_0$ such that we have for all x, u with $x \geq u \geq x_1$:

$$\frac{\Psi'(x)}{\Psi'(u)} \geq \frac{1}{\beta} (x \geq u \geq x_1). \quad (26)$$

Then the series (18) is divergent.

22. Observe that the condition (26) is certainly satisfied from a certain x_1 on, if $\Psi(x)$ has a finite limit ω ,

$$\Psi'(x) \rightarrow \omega < \infty (x \rightarrow \infty). \quad (27)$$

23. *Proof of the Theorem 6.* Since x_0 can be replaced by any greater number we can assume, without loss of generality, that $x_1 = x_0$. Then we proceed as in the proof of the Theorem 4 defining $F(x)$ by (19) and obtain, as in the section 15, using (26):

$$\begin{aligned} F(\Psi(x)) \Psi'(x) &= \lim_{\kappa \rightarrow \infty} f(\Psi(v_\kappa)) \Psi'(x) \geq \frac{1}{\beta} \overline{\lim}_{\kappa \rightarrow \infty} f(\Psi(v_\kappa)) \Psi'(v_\kappa) \\ &\geq \overline{\lim}_{\kappa \rightarrow \infty} f(v_\kappa) \geq F(x). \end{aligned}$$

24. We see that $F(x)$ satisfies the conditions of the Theorem 2; therefore the integral $\int_0^\infty F(x) dx$ is divergent and the same holds for the series $\sum_0^\infty F(\nu)$, as $F(x)$ is monotonically decreasing. But then the series (18) is also divergent since $f(x)$ is a majorant of $F(x)$. The Theorem 6 is proved.

IV. ANOTHER METHOD IN THE CASE OF DIVERGENCE

25. **THEOREM 7.** *The assertion of the Theorem 4 remains valid if the assumption that $\Psi'(x)$ is monotonically increasing is replaced by the assumption that $\Psi'(x)$ is monotonically decreasing.*

26. *Proof.* Since in any case $\Psi'(x) \geq 0$ there exists a finite ω such that

$$\Psi'(x) \downarrow \omega \quad (x \rightarrow \infty)$$

and, as in the sec. 17, we see that this limit is ≥ 1 .