

# ON THE CLOSED CONVEX HULL OF A COMPACT SET

Autor(en): **Goodner, Dwight B.**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **16 (1970)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-43862>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# ON THE CLOSED CONVEX HULL OF A COMPACT SET

by Dwight B. GOODNER

Although it is known that the closed convex hull of a compact set may not be compact, the literature contains few, if any, reasonably easy proofs of this theorem, and existing proofs seem to be difficult to find. Unfortunately many students encounter this result before their training and experience have prepared them to prove it. The purpose of this note is to give a proof that seems to be minimal in terms of the knowledge and mathematical sophistication required to understand it.

*Theorem.* There is a linear topological space in which the closed convex hull of a certain compact set is not compact.

*Proof.* Let  $m$  be the space of bounded real-valued sequences with the usual supremum norm. We proceed by identifying a compact set  $S$  [2, p. 135] contained in a linear subspace  $L$  [1, p. 12] of  $m$  such that in  $L$  the convex hull [1, p. 10] of  $S$  is closed but is not compact.

Let  $\omega$  be the directed system of positive integers. For fixed  $i \in \omega$  let  $\delta^i = \{ \delta^i_j \}_{j \in \omega}$  be the sequence defined by  $\delta^i_i = 1$  and  $\delta^i_j = 0$  for  $i \neq j$ ;

$\delta^i_j$  is the Kronecker delta. We note that  $\delta^i \in m$ . Let  $S = \{ \Theta \} \cup \{ \frac{\delta^i}{i} : i \in \omega \}$

where  $\Theta$  is the zero element in  $m$ , and let  $K(S)$  be the convex hull of  $S$  in  $m$ .

Since  $\lim_{i \rightarrow \infty} \frac{\delta^i}{i} = \Theta$ , each open neighborhood of  $\Theta$  contains all but finitely

many elements of the sequence  $\{ \frac{\delta^i}{i} \}_{i \in \omega}$ , and we see that  $S$  is compact.

$$\text{Let } x_1 = 1 \frac{\delta^1}{1}$$

$$x_2 = \frac{1}{2} \frac{\delta^1}{1} + \frac{1}{2} \frac{\delta^2}{2}$$

$$\begin{aligned}
 x_3 &= \frac{1}{2} \frac{\delta^1}{1} + \frac{1}{4} \frac{\delta^2}{2} + \frac{1}{4} \frac{\delta^3}{3} \\
 &\quad \cdot \quad \cdot \quad \cdot \\
 x_i &= \frac{1}{2} \frac{\delta^1}{1} + \frac{1}{4} \frac{\delta^2}{2} + \cdots + \frac{1}{2^{i-1}} \frac{\delta^{i-1}}{i-1} + \frac{1}{2^{i-1}} \frac{\delta^i}{i} \\
 &\quad \cdot \quad \cdot \quad \cdot
 \end{aligned}$$

For each  $i$ ,  $x_i \in K(S)$  [1, p. 10]. If

$$x_0 = \lim_{i \rightarrow \infty} x_i = \lim_{i \rightarrow \infty} \left( \sum_{n=1}^{i-1} \frac{1}{2^n} \frac{\delta^n}{n} + \frac{1}{2^{i-1}} \frac{\delta^i}{i} \right) = \lim_{i \rightarrow \infty} \sum_{n=1}^i \frac{1}{2^n} \frac{\delta^n}{n},$$

then  $x_0 \notin K(S)$  but  $x_0 \in \text{cl } K(S)$  where  $\text{cl } K(S)$  is the closure of  $K(S)$  in  $m$ . We remark that this shows that in a linear topological space the convex hull of a convex set may not be closed and, hence, may not be compact [2, p. 141]. We note but do not prove that the closed convex hull,  $\text{cl } K(S)$ , of  $S$  is compact.

Let  $L$  be the set of all linear combinations of all finite subsets of  $S$ . Then  $L$  with the relative topology of  $m$  [2, p. 51] is the smallest linear subspace [1, p. 2] of  $m$  containing  $K(S)$ , and  $K'(S) = L \cap \text{cl } K(S)$  is the closed convex hull of  $S$  in  $L$ . To complete our proof it suffices to show that  $K'(S)$  is not a compact subset of  $L$ .

The sequence  $\{x_i\}_{i \in \omega}$  defined above is a closed subset of  $L$  since its only cluster point  $x_0 \notin L$ . For each  $k \in \omega$  let  $A_k = \{x \in L : x \notin \{x_i\}_{i=k}^\infty\}$ . Each  $A_k$  is an open subset of  $L$  and  $A = \{A_k : k \in \omega\}$  is an open cover of  $K'(S)$ . However, if  $A_{k(1)}, A_{k(2)}, \dots, A_{k(n)}$  is a finite subset of  $A$  and if  $j$  is any integer greater than  $\max\{k(1), k(2), \dots, k(n)\}$ , then  $x_j \notin \bigcup_{i=1}^n A_{k(i)}$ . Since  $x_j \in \{x_i\}_{i \in \omega} \subset K'(S)$  it follows that  $A$  does not contain a finite subset which covers  $K'(S)$  and, hence, that  $K'(S)$  is not a compact subset of  $L$  which completes the proof.

#### REFERENCES

- [1] M. M. DAY, *Normed linear spaces*. Academic Press, New York and Springer, Berlin, 1962.
- [2] J. L. KELLEY, *General topology*. D. Van Nostrand Co., Princeton, 1955.

D. B. GOODNER

Dept. of Math.  
 Florida State University  
 Tallahassee, Florida 32306