

Introduction

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **27 (1981)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **29.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

ON THE GENUS OF GENERALIZED FLAG MANIFOLDS

by Henry H. GLOVER and Guido MISLIN

INTRODUCTION

Let X be a nilpotent space of finite type. We denote by $G(X)$ the genus of X , i.e. the set of all homotopy types Y (nilpotent, of finite type) with p -localizations $Y_p \simeq X_p$ for all primes p , (cf. [HMR]). The set $G(X)$ has been studied extensively in case of X an H -space. In particular it is known that for the special unitary group $SU(n)$ one has

$$|G(SU(n))| \geq \prod_{1 < m < n} (\phi(m!)/2)$$

where ϕ is the Euler function [Z, p. 152]. We are interested in this note in finding non-trivial examples X with $G(X) = \{[X]\}$ and we call such spaces *generically rigid*. A large family of such generically rigid spaces is provided by certain generalized flag manifolds. Let

$$G = U(n_1 + n_2 + \dots + n_k)$$

and

$$H = U(n_1) \times U(n_2) \times \dots \times U(n_k),$$

embedded in G in the obvious way. Then

$$M = M(n_1, n_2, \dots, n_k) = G/H$$

is a generalized flag manifold (generalizing the standard complex flag manifold $U(n)/T^n$ which corresponds to $M(1, 1, \dots, 1)$). We will show essentially that whenever the homotopy rigidity result for linear actions holds for M (cf. [L1], [L2], [EL]), then M is also generically rigid. These two seemingly unrelated rigidity results are tied up by certain results on $E(X)$ and $E(X_0)$, the groups of homotopy classes of self equivalences of X and X_0 , X_0 the rationalization of X .

To make our result more precise, we need some further notation. For

$$M = M(n_1, \dots, n_k) = G/H$$

as above, we write $N(H)$ for the normalizer of H in G . The finite group $N(H)/H$ acts on M in an obvious way and it is well known that through that action, $N(H)/H$ is faithfully represented in $H^*(M; \mathbf{Q})$. We can therefore consider $N(H)/H$ as a subgroup of $E(M)$ or $E(M_0)$. By Theorem 1.1 of [GH2] the canonical map

$$E(M_0) \rightarrow \text{Aut}_{\text{alg}} H^*(M; \mathbf{Q})$$

is a group isomorphism. In particular, the grading automorphisms

$$g(q): H^*(M; \mathbf{Q}) \rightarrow H^*(M; \mathbf{Q})$$

defined by $g(q)x = q^i x$ for $x \in H^{2i}(M; \mathbf{Q})$ and $q \in \mathbf{Q}^*$, lift to unique self equivalences of M_0 (which we denote also by $g(q)$), and thus

$$Gr(M_0) = \{g(q) \mid q \in \mathbf{Q}^*\} \subset E(M_0)$$

is a central subgroup isomorphic to \mathbf{Q}^* .

In all cases of generalized flag manifolds for which $E(M_0)$ has been computed, the subgroup generated by $Gr(M_0)$ and $N(H)/H$,

$$\langle Gr(M_0), N(H)/H \rangle \subset E(M_0)$$

is all of $E(M_0)$. The following conjecture is thus plausible.

Conjecture C. Let $M = M(n_1, n_2, \dots, n_k)$ be a generalized flag manifold. Then

$$E(M_0) = \langle Gr(M_0), N(H)/H \rangle.$$

A similar conjecture appears in [L1, Conjecture C] but the relationship between the two conjectures is not entirely clear.

The Conjecture C has been verified in the following cases:

- 1) $n_1 = n_2 = \dots = n_k = 1$ (compare the proof of Thm. 1 in [EL])
- 2) $n_1 = n_2 = \dots = n_{k-1} = 1, n_k \geq k - 1$ (compare the proof of Theorem 9 in [L1])
- 3) $n_1 = 2$ and $k = 2$ (follows from [O])
- 4) $n_2 > n_1$ and $k = 2$ ([GH1], [Br])
- 5) $n_1 = 1, n_2 > 1, n_3 \geq 2n_2^2 - 1$ and $k = 3$ ([GH2])

The Conjecture C holds therefore for instance for all complex Grassmann manifolds $G_p(\mathbf{C}^{p+q}) = M(p, q)$ with $p \neq q$ (since $M(p, q) \simeq M(q, p)$), and for the classical flag manifolds $U(n)/T^n$.

Our main theorem may be stated as follows.

THEOREM. Let $M = M(n_1, \dots, n_k)$ be a generalized flag manifold for which the Conjecture C holds. Then

$$G(M) = \{[M]\}.$$

In particular the Grassmann manifolds $G_p(\mathbb{C}^{p+q})$ for $p \neq q$ and the flag manifolds $U(n)/T^n$ are all generically rigid.

§1. GENUS AND SELF MAPS

Let P denote a fixed set of primes. Two P -sequences

$$S_1, S_2 : P \rightarrow E(X_0)$$

are called *equivalent*, if there exist maps $h(0) \in E(X_0)$ and

$$h(p) \in \text{im}(E(X_p) \xrightarrow{\text{can}} E(X_0))$$

such that for all $p \in P$ one has

$$h(0) S_1(p) = S_2(p) h(p).$$

Definition 1.1. We denote by $P\text{-Seq}(E(X_0))$ the set of equivalence classes of P -sequences in $E(X_0)$.

If P is a finite set of primes and X a nilpotent space of finite type, then there is a canonical map

$$\theta : G(X) \rightarrow P\text{-Seq}(E(X_0)).$$

It is defined as follows. Let $Y \in G(X)$ and $P = \{p_1, \dots, p_n\}$. Then the localization Y_P is a pull-back of maps $X_{p_i} \xrightarrow{\lambda_i} X_0$, i.e. $Y_P \simeq \text{hoinvlim} \{X_{p_i} \xrightarrow{\lambda_i} X_0\}$. The maps λ_i induce equivalences $\bar{\lambda}_i \in E(X_0)$ and we put

$$\theta(Y) = \{[\bar{\lambda}_1, \bar{\lambda}_2, \dots, \bar{\lambda}_n]\}.$$

If Y_P may also be represented by $\text{hoinvlim} \{X_{p_i} \xrightarrow{\mu_i} X_0\}$, then there exist maps $h(0) \in E(X_0)$ and $\tilde{h}(p_i) \in E(X_{p_i})$, $i \in \{1, \dots, n\}$ rendering the diagrams