

2. Statement of Tits' theorem

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **03.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

In 1914, this example allowed Hausdorff to show that there does not exist any finitely additive rotation-invariant measure defined on all subsets of the sphere S^2 . See [H], and [DE] for subsequent history. While discussing this, let us mention the following open problem (brought to my attention by M. Keane): does there exist a finitely additive probability measure on the Borel subsets of S^2 , vanishing on meagre sets, invariant under rotations? (The answer for countably additive measures is no, and follows from the unicity of Haar measure on a compact group; see e.g. §9 in [Wi].)

Remark. Let G be a connected real Lie group. Then G contains at least one subgroup isomorphic to the free group on two generators F_2 if and only if G is not solvable, as results from standard Lie theory as follows.

To check the non trivial implication, we assume that G is not solvable, so that G contains a semi-simple subgroup S by a theorem of Levi and Mal'cev. Consider a Cartan decomposition $\mathfrak{s} = \mathfrak{k} \oplus \mathfrak{p}$ of the Lie algebra of S . If $\mathfrak{k} \neq \{0\}$, root theory shows that the semi-simple compact algebra \mathfrak{k} contains a subalgebra isomorphic to $\mathfrak{su}(2)$, so that G contains a subgroup isomorphic to one of $SU(2)$, $SO(3)$. If $\mathfrak{k} = \{0\}$, then \mathfrak{s} is split and root theory again shows that \mathfrak{s} contains a copy of $\mathfrak{sl}(2, \mathbf{R})$, so that G contains a subgroup isomorphic to a covering of $PSL(2, \mathbf{R})$. In all cases, examples above show that G contains a copy of F_2 .

So, let G be a connected Lie group containing a copy of F_2 . For $w \in F_2 - \{1\}$ and $g, h \in G$, let $w(g, h)$ be the element of G obtained when replacing the two generators of F_2 by g and h in w . Then

$$X_w = \{ (g, h) \in G \times G \mid w(g, h) = 1 \}$$

has empty interior (think of analytic continuation). It follows from Baire's theorem that the set $G \times G - \bigcup X_w$ (union over $w \in F_2 - \{1\}$) of those $(g, h) \in G \times G$ such that g and h generate a free group is dense and has full measure in $G \times G$ [E]. (If G is moreover semi-simple, it follows from a note by Kuranishi and from Tits' theorem that there exist $g, h \in G$ generating a subgroup of G which is both free and dense [Ku].)

2. STATEMENT OF TITS' THEOREM

Recall that, given a group Γ , its derived group $D\Gamma$ is the subgroup generated by elements of the form $ghg^{-1}h^{-1}$ and that Γ is *solvable* if $D(\dots D(\Gamma) \dots) = \{1\}$ for sufficiently many D 's. We say that Γ is *almost solvable* (other people say *virtually solvable*) if it contains a solvable subgroup of finite index. For example, groups of

triangular matrices are solvable and non abelian free groups are not almost solvable. By "free group", we mean hereafter *non abelian free group*.

A *linear group* over a field \mathbf{K} is a group which has at least one faithful finite dimensional representation over \mathbf{K} , namely a group isomorphic to a subgroup of $GL(n, \mathbf{K})$ for some n . Groups are far from being all linear, even under the hypothesis of finite generation. Famous examples of non linear groups are the quotients F_2/F_2^m for m odd and large enough, where F_2^m is the subgroup of the free group F_2 generated by elements of the form g^m . (Novikov's negative solution to the Burnside problem; in the original paper, m large enough means $m \geq 4381$.)

Easier examples are provided by finitely generated infinite simple groups (there is such a group, discovered by G. Higman, which is described in [S], n° I.1.4). They are not linear, because it is a result of Mal'cev that a finitely generated linear group Γ is residually finite [M]. (This means that, for any $\gamma \in \Gamma - \{1\}$, there exists a homomorphism φ of Γ onto a finite group with $\varphi(\gamma) \neq 1$; instructive and easy exercise: check that $SL(n, \mathbf{Z})$ is residually finite.)

Also, any finitely generated non hopfian group cannot be linear (Γ is non hopfian if there exists a non injective homomorphism of Γ onto itself); an example of such a group is that generated by two elements g, h submitted to the relation $h^{-1}g^2h = g^3$ (see [LS], page 197).

TITS' THEOREM. *A linear group Γ over a field \mathbf{K} of characteristic 0 which is not almost solvable contains a free group.*

This theorem has been conjectured by Bass and Serre, and proved in [T] together with other results, some concerning positive characteristics.

The following precision has been added by Wang [Wa]: there exists for each positive integer n a constant $\lambda(n)$ such that any subgroup of $GL(n, \mathbf{K})$ without free subgroup contains a solvable subgroup of index smaller than $\lambda(n)$.

Let Γ be a group having a finite set of generators S which is a subgroup of $GL(n, \mathbf{K})$ for some n . If k is the subfield of \mathbf{K} generated by entries of elements of S , then $\Gamma \subset GL(n, k)$. As k is finitely generated of characteristic zero, there exists an embedding of k in \mathbf{C} and one may assume that Γ lies in $GL(n, \mathbf{C})$. For finitely generated groups (and also in the general case by [Wh]), it is consequently sufficient to prove Tits' theorem for $\mathbf{K} = \mathbf{C}$ (or $\mathbf{K} = \mathbf{R}$ because $GL(n, \mathbf{C})$ is a subgroup of $GL(2n, \mathbf{R})$). But this apparent simplification (?) is deceptive, because the proof does require other fields than fields of complex numbers.

It follows from the theorem that a linear group over a field of characteristic zero which is not amenable contains a free group; this answers for linear groups a question formulated by J. von Neumann [vN]. Another famous result whose

proof requires Tits' theorem is due to Gromov: a finitely generated group has polynomial growth if and only if it is almost nilpotent [G].

The analogue of Tits' theorem for division rings does not hold as such [L1], but conjectural statements have been formulated [L2]. Another generalisation of the theorem is proposed as a research problem in remark 1.4.2 of [BL].

3. DIGRESSION ON HYPERBOLIC GEOMETRY

Let n be an integer, $n \geq 1$. The hyperbolic space H^{n+1} of dimension $n + 1$ is the open unit ball of the euclidean space \mathbf{R}^{n+1} . Hyperbolic lines (called lines below) in H^{n+1} are traces on H^{n+1} of circles and euclidean lines in \mathbf{R}^{n+1} which are orthogonal to \mathbf{S}^n . Two distinct points $P, Q \in H^{n+1}$ are on a unique line which determines two points $P_\infty, Q_\infty \in \mathbf{S}^n$, say with P, Q, Q_∞, P_∞ arranged in cyclic order on the euclidean circle defining this line. The (hyperbolic) distance between P and Q is given by a cross-ratio of euclidean distances; more precisely, it is defined to be

$$d(P, Q) = \text{Log}(P, Q, Q_\infty, P_\infty) = \log \left(\frac{|P - Q_\infty|}{|P - P_\infty|} : \frac{|Q - Q_\infty|}{|Q - P_\infty|} \right).$$

The *proper Mæbius group* $GM(n)_0$ is the group of orientation preserving isometries of \mathbf{R}^{n+1} for this distance. Any $g \in GM(n)_0$ extends to a homeomorphism of the closed ball $H^{n+1} \cup \mathbf{S}^n$. One may check that $GM(1)_0$ is isomorphic to $PGL(2, \mathbf{R})$ and $GM(2)_0$ to $PGL(2, \mathbf{C})$.

There is an equivalent description with H^{n+1} the half space $\mathbf{R}^n \times \mathbf{R}_+^*$. The set of "points at infinity" is then $\mathbf{R}^n \cup \{\infty\}$ rather than \mathbf{S}^n .

For all this, see e.g. [A] or [Si].

An isometry $g \in GM(n)_0$ is said to be

elliptic if there is some point in H^{n+1} fixed by g ,

parabolic if there is in \mathbf{S}^n exactly one point fixed by g ,

hyperbolic if there is a line in H^{n+1} invariant by g on which g has no fixed point.

(Following Thurston [Th], we call "hyperbolic" elements which are "loxodromic" in classical literature, such as in [Gr].)

PROPOSITION. *Elliptic, parabolic and hyperbolic elements define a partition of the proper Mæbius group in three disjoint classes.*

Proof. Let us first check that the three classes do not overlap in $GM(n)_0$. If g is hyperbolic, it has two fixed points in \mathbf{S}^n and thus cannot be parabolic; if g was