

SOME ALMOST HOMOGENEOUS GROUP ACTIONS ON SMOOTH COMPLETE RATIONAL SURFACES

Autor(en): **Moser-Jauslin, Lucy**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **34 (1988)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-56601>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

SOME ALMOST HOMOGENEOUS GROUP ACTIONS ON SMOOTH COMPLETE RATIONAL SURFACES

by Lucy MOSER-JAUSLIN

In this article we are interested in certain actions of a Borel subgroup of $SL(2)$ on rational surfaces. More specifically, let X be a complete smooth rational surface over an algebraically closed field k of characteristic zero. Let B be the linear algebraic group defined by

$$B = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \mid \alpha \in k^*, \beta \in k \right\}.$$

We study all the actions of B on X such that there is an *open* orbit. This orbit is necessarily isomorphic to B/Γ where Γ is a finite cyclic subgroup of B .

Any complete smooth rational surface is obtained by blowing up one of the minimal rational surfaces, which are well-known (see for example [Har], [Beau] or [Saf]). In section 1, we generalize this result to surfaces with a B -action: that is, any smooth complete rational surface with an action of B is obtained by blowing up one of the minimal surfaces with a B -action. Thus we are reduced to studying actions on the minimal rational surfaces.

In section 2 we state the main result of the article. We give a complete list of B -actions on each of the minimal models which have an open orbit. There are two methods to do this. First, one can find all possible homomorphisms of B into the automorphism group of the surface which yield the desired actions. Secondly one can study *geometrically* the complement to the open orbit with the action of B . In this article we use the latter approach.

The problem considered here is useful for the study of $SL(2)$ -embeddings. This is explained in section 3.

The minimal rational surfaces are *almost homogeneous*. That is, they contain an open dense orbit with respect to the action of its automorphism group. In [Pot] all complex analytic almost homogeneous surfaces are classified. It is shown that any such surface is either a rational surface,

a topologically trivial \mathbf{P}^1 -bundle over a one-dimensional complex torus, a Hopf surface with abelian fundamental group, or a two-dimensional complex torus. (See also [H-O].) There have been other studies of almost homogeneous surfaces. For example in [Pop] the author describes those which are affine and such that the complement to the open orbit is a finite set of points. In these studies one is primarily interested in the surfaces. In this article, however, we are given the surface and the group, and we are interested in the action.

I would like to thank Th. Vust, D. Luna, H. Kraft, and M. Brion for helpful discussions and comments.

§ 1. MINIMAL EMBEDDINGS: DEFINITIONS AND PRELIMINARY REMARKS

Let G be a connected algebraic group and let H be an algebraic subgroup.

Definition. An *embedding* of the homogeneous space G/H is a reduced irreducible algebraic variety X endowed with a regular action of G having an open orbit isomorphic to G/H . Two embeddings are equivalent if they are G -isomorphic.

In this paper we study all smooth complete embeddings of B/Γ , where Γ is a finite subgroup of B (any such Γ is cyclic, and two finite subgroups of the same order are conjugate). Since B/Γ is rational and two-dimensional, the underlying variety of such an embedding is a smooth complete rational surface.

Given a smooth complete B/Γ -embedding X with fixed point x , the action of B on X induces an action on \tilde{X} , the variety obtained by blowing up x in X , giving \tilde{X} the structure of a B/Γ -embedding. (This is a consequence of the universal property of blowing up. See e.g. [Har], p. 164. See also [O-W], pp. 48-49.) We say that X is a *minimal B/Γ -embedding* if it is not the blow up of another smooth B/Γ -embedding. If X is a minimal model as a variety (that is, if the underlying variety of X is not the blow up of another smooth variety), then clearly X is a minimal embedding. We will now prove the converse.

LEMMA 1.1. *Suppose X is a smooth complete surface on which a connected linear algebraic group H acts regularly. Suppose also that X contains an irreducible curve C with a strictly negative self-intersection number. Then C is stable by H .*

Proof. Let $s \in H$. Then since H is connected and the action is regular, sC is linearly equivalent to C . (See e.g. [Gro], p. 5-06, Lemme 1 or [Kam]. See also [O-W], p. 49 and [Ful] for related results.) Thus the intersection number $sC \cdot C$ equals the self-intersection number $C \cdot C$. Since sC is irreducible, the assumption $sC \neq C$ implies that $sC \cdot C$ is non-negative, since these curves intersect in a finite number of points, each counted with positive multiplicity. \square

PROPOSITION 1.2. *Suppose X is a minimal B/Γ -embedding. Then it is a minimal model as a variety; that is, X is a rational minimal model.*

Proof. If X is not a minimal model as a variety, then it contains an irreducible curve C isomorphic to \mathbf{P}^1 with self-intersection -1 . (Castelnuovo criterion. See e.g. [Har] p. 414.) If we apply Lemma 1.1 to the case $H = B$, we see that C is stable by B . By Zariski's Main Theorem (projective-smooth case) (see [Mum], p. 52), the action of B on X induces an action on the surface obtained by blowing down C . Thus this new surface is also a B/Γ -embedding, and X is not a minimal embedding. Also X must be rational, because B/Γ is rational. \square

We recall the description of the set of minimal models of rational surfaces (see for example [Har] Section V.2, [Beau] Ch. IV, or [Saf] Ch. V). For any integer $n \geq 0$, define $F_n = \mathbf{P}(\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(n))$. (For $k = \mathbf{C}$ these surfaces are known as the Hirzebruch surfaces.) Then F_n is a ruled surface over \mathbf{P}^1 . For example, $F_0 \cong \mathbf{P}^1 \times \mathbf{P}^1$ and F_1 is the blow up of \mathbf{P}^2 in one point. The set of minimal rational models is given by \mathbf{P}^2 and $F_n, n \neq 1$.

Let us review some elementary properties of the surfaces F_n . These facts can be found in the references above. As mentioned above, F_n is a ruled surface over \mathbf{P}^1 ; that is, it is a \mathbf{P}^1 -fibre bundle over \mathbf{P}^1 . We restrict to the case $n \geq 1$. Then there is exactly one ruling of F_n , i.e. there is exactly one morphism $\pi_n: F_n \rightarrow \mathbf{P}^1$ with fibres isomorphic to \mathbf{P}^1 . The bundle $\pi_n: F_n \rightarrow \mathbf{P}^1$ has a unique section E_n with self-intersection $-n$, and E_n is the only irreducible curve of F_n with strictly negative self-intersection. The fibres of π_n are all linearly equivalent, and they are the only irreducible curves with self-intersection 0. So any automorphism of F_n stabilizes E_n and permutes the fibres. Now $F_n - E_n$ is the total space of the vector bundle $\mathcal{O}(n)$ over \mathbf{P}^1 . All the sections of $\mathcal{O}(n)$ are linearly equivalent (as divisors of F_n) with self-intersection n .

If one contracts the section E_n of $F_n, n \geq 1$, one obtains a surface X_n (nonsingular if and only if $n = 1$) contained in \mathbf{P}^{n+1} . In fact X_n is the closure

of the affine cone over the n -tuple embedding $\mathbf{P}^1 \rightarrow \mathbf{P}^n$ (see [Beau] Ch. IV, Ex. 1 or [G-H], p. 523). That is,

$$X_n = \{(z_0 : s^n : s^{n-1}t : \dots : t^n) \mid z_0, s, t \in k\} \subset \mathbf{P}^{n+1}.$$

The vertex of the cone X_n is $(1:0:\dots:0)$. The image of a general fibre of F_n in X_n is given by choosing s and t such that $\alpha s = \beta t$ for some $(\alpha:\beta) \in \mathbf{P}^1$.

One can also construct the surfaces F_n inductively: given $F_n, n \geq 1$, one blows up a point x on E_n and then blows down the strict transform of the fibre containing x to obtain F_{n+1} . The rational map thus obtained from F_n to F_{n+1} is sometimes called an *elementary transformation*. (See e.g. [Saf] Ch. V.)

Also, for $n \geq 1$, we have an exact sequence

$$1 \rightarrow k^* \rtimes H^0(\mathbf{P}^1, \mathcal{O}(n)) \rightarrow \text{Aut } F_n \xrightarrow{\Phi} \text{PGL}(2) \rightarrow 1$$

where Φ is the restriction of an automorphism to $E_n \cong \mathbf{P}^1$, and k^* acts on $H^0(\mathbf{P}^1, \mathcal{O}(n))$ by multiplication. The kernel of Φ is the subgroup of automorphisms that fix the fibres of π_n . (See [Beau] Ch. V, Ex. 4.)

We define an action of $\text{Aut } F_n$ on $H^0(\mathbf{P}^1, \mathcal{O}(n))$ as follows. If $\varphi \in \text{Aut } F_n$ and s is a global section of $\mathcal{O}(n)$, then φs is the section given by $(\varphi s)(x) = \varphi(s(\varphi^{-1}x))$, where $x \in \mathbf{P}^1$ and the action of φ^{-1} on \mathbf{P}^1 is given by its action on $E_n \cong \mathbf{P}^1$. Thus $(\varphi s)(\mathbf{P}^1) = \varphi(s(\mathbf{P}^1))$.

LEMMA 1.3. *Let $\varphi \in \text{Aut } F_n, n \geq 1$; then the action of φ on the vector space $H^0(\mathbf{P}^1, \mathcal{O}(n))$ given above is an affine transformation.*

Proof. One has to check that for $s_1, s_2 \in H^0(\mathbf{P}^1, \mathcal{O}(n))$ and $t \in k^*$ we have that $\varphi(ts_1 + (1-t)s_2) = t(\varphi s_1) + (1-t)(\varphi s_2)$. We use that given $x \in \mathbf{P}^1$ the restriction of φ to the fibre $\varphi^{-1}(\pi_n^{-1}x)$ gives an isomorphism

$$k \cong \varphi^{-1}(\pi_n^{-1}x) \xrightarrow{\sim} \pi_n^{-1}x \cong k;$$

this transformation is affine. Now suppose we have s_1, s_2 , and t as above; let $s = ts_1 + (1-t)s_2$. Then for any $x \in \mathbf{P}^1$ we have

$$\begin{aligned} (\varphi s)x &= \varphi(s(\varphi^{-1}x)) = \varphi(ts_1(\varphi^{-1}x) + (1-t)s_2(\varphi^{-1}x)) \\ &= t\varphi(s_1(\varphi^{-1}x)) + (1-t)\varphi(s_2(\varphi^{-1}x)) = t(\varphi s_1)x + (1-t)(\varphi s_2)x. \end{aligned}$$

This proves the lemma. □

Thus for $n \geq 1$, there is a homomorphism $\text{Aut } F_n \rightarrow \text{Aff}(H^0(\mathbf{P}^1, \mathcal{O}(n)))$ given by $\varphi \rightarrow (s \mapsto \varphi s)$.

To describe a B/Γ -embedding with underlying variety X , we must give a homomorphism $B \rightarrow \text{Aut } X$ such that X has an open orbit B -isomorphic to B/Γ . Two such homomorphisms give rise to equivalent embeddings if and only if they are conjugate.

In the following section we will use the information given here to study the possible B/Γ -embeddings into \mathbf{P}^2 , $\mathbf{P}^1 \times \mathbf{P}^1$, and \mathbf{F}_n , $n \geq 1$.

§ 2. THE MINIMAL B/Γ -EMBEDDINGS

THEOREM 2.1. *Let Γ be a finite subgroup of B , and let X be the projective plane \mathbf{P}^2 or a rational ruled surface \mathbf{F}_n (with $n \geq 0$, where $\mathbf{F}_0 = \mathbf{P}^1 \times \mathbf{P}^1$).*

(i) *The number $\text{emb}(X)$ of equivalence classes of B/Γ -embeddings into X with at least two fixed points is*

$$\text{emb}(\mathbf{P}^2) = 2, \quad \text{emb}(\mathbf{P}^1 \times \mathbf{P}^1) = 1, \quad \text{and} \quad \text{emb}(\mathbf{F}_n) = n + 3, \quad n \geq 1.$$

We call these the "ordinary" embeddings.

(ii) *Moreover, for any such surface X , there is exactly one subgroup Γ and an "exceptional" B/Γ -embedding into X with only one fixed point (up to equivalence), and the corresponding order $\text{ord}(X)$ of this group Γ is*

$$\text{ord}(\mathbf{P}^2) = 4, \quad \text{ord}(\mathbf{P}^1 \times \mathbf{P}^1) = 2, \quad \text{and} \quad \text{ord}(\mathbf{F}_n) = 2(n+1), \quad n \geq 1.$$

(iii) *The complement to the open orbit consists of two (for \mathbf{P}^2) resp. three (for the \mathbf{F}_n) smooth rational curves, intersecting transversely, except in the "exceptional" case with $X = \mathbf{P}^2$, in which case the two curves are tangent.*

(In this theorem we include the case \mathbf{F}_1 even though it is not minimal.)

To be more precise, we indicate the form of the complement Z to the open orbit in each case. Also to distinguish the embeddings where Z has the same form, we indicate how the action of B differs on Z . Let U be the unipotent radical of B and T be a maximal torus. (That is, U is the subgroup of elements of B where both eigenvalues are 1, and T can be chosen to be the subgroup of diagonal elements.) Then B is $T \ltimes U$, and the characters of B are the characters of T . We denote the character group of B by $\{\alpha^n : n \in \mathbf{Z}\}$.

Denote by c the order of the group Γ .

Embeddings into \mathbf{P}^2 :

(i) “Ordinary” embeddings: We find that for each Γ there are two embeddings where $Z = L_1 \cup L_2$ and L_1 and L_2 are lines in \mathbf{P}^2 . The group B acts on L_1 in the standard manner and on L_2 by the character α^{2+c} or α^{2-c} . There are two fixed points except in one embedding for the case $c = 2$, where L_2 is a line of fixed points. See Fig. 1a.

(ii) The “exceptional” embedding: If $c = 4$, we also find an embedding where $Z = L_1 \cup C$ and C is a smooth conic which is tangent to L_1 at the unique fixed point. See Fig. 1b.




FIGURE 1.

Embeddings into $\mathbf{P}^1 \times \mathbf{P}^1$:

In this case, Z is always the union of three curves. Let $p_i: \mathbf{P}^1 \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$, $i = 1, 2$ be the two projections.

(i) "Ordinary" embeddings: For each Γ there is an embedding where $Z = F_1 \cup F'_1 \cup F_2$ and F_1, F'_1 are fibres of p_1 and F_2 is a fibre of p_2 . There are two fixed points. See Fig. 1c.

(ii) The "exceptional" embedding: Also, if $c = 2$, we find another embedding into $\mathbf{P}^1 \times \mathbf{P}^1$ where $Z = F_1 \cup F_2 \cup C$, and C is a section of p_1 and p_2 which intersects F_1 and F_2 transversely in the unique fixed point. See Fig. 1d.

Embeddings into \mathbf{F}_n , $n \geq 1$:

Again Z is always the union of three curves. Let $\pi_n: \mathbf{F}_n \rightarrow \mathbf{P}^1$ be the unique ruling of \mathbf{F}_n , and let E_n be the irreducible curve of \mathbf{F}_n with self-intersection $-n$.

(i) "Ordinary" embeddings: For each Γ we find $n + 1$ cases where $Z = E_n \cup F \cup F'$ and F and F' are fibres of π_n . The torus T acts on F by the character α^{cp+2} and on F' by the character $\alpha^{-c(n-p)+2}$, $p = 0, \dots, n$. There are either 3 or 4 fixed points (depending on the action of U on F and F'), or, if T acts trivially on F' , then F' is a curve of fixed points. See Fig. 1e.

There are also two other embeddings in \mathbf{F}_n for each Γ where $Z = F \cup E_n \cup D$ and F is a fibre as before and D is a section of π_n which does not intersect E_n . The group B acts on F by the character $\alpha^{2n \pm c}$. There are two fixed points except in one of the embeddings in the case where $c = 2n$, in which case F consists entirely of fixed points. See Fig. 1f.

(ii) "Exceptional" embeddings: Also if $c = 2(n+1)$, there is one more embedding where $Z = E_n \cup F \cup C$ and C is a section which intersects E_n and F transversely in the unique fixed point. See Fig. 1g. This embedding is obtained as follows. Consider the embedding into \mathbf{F}_{n+1} of the previous type where the fibre F consists of fixed points. Blow up a point of F which is not on E_{n+1} or D and contract the strict transform of F . This gives the required embedding into \mathbf{F}_n .

The explicit matrix representations of the different B -actions are given in the proof of the theorem.

Proof of the Theorem. Throughout the proof we denote the order of the group Γ by c .

Recall that to give an embedding of B/Γ into a variety X , we must find a homomorphism $\varphi: B \rightarrow \text{Aut } X$ such that under the induced action of B on X , there is an open orbit isomorphic to B/Γ . Two such embeddings are equivalent if and only if the homomorphisms are conjugate.

We have $B = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \mid \alpha \in k^* \text{ and } \beta \in k \right\}$, $U = \left\{ \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} \mid \beta \in k \right\}$,

and set $T = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \mid \alpha \in k^* \right\}$.

We consider separately the embeddings into \mathbf{P}^2 , $\mathbf{P}^1 \times \mathbf{P}^1$ and \mathbf{F}_n , $n \geq 1$.

Embeddings into \mathbf{P}^2 :

If B acts on \mathbf{P}^2 , it has a fixed point o since \mathbf{P}^2 is complete and B is solvable (see e.g. [Bor], p. 242). Also B acts on the linear system $S = \{\text{lines of } \mathbf{P}^2 \text{ passing through } o\}$. Since we have $S \cong \mathbf{P}^1$, B stabilizes one such line, which we call L . We can choose homogeneous coordinates $(z_0:z_1:z_2)$ of \mathbf{P}^2 such that $o = (1:0:0)$ and $L = (z_0:z_1:0)$; thus $\varphi(B) \subset PGL(3)$ is upper triangular.

CASE 1. U acts trivially on L .

Then there is another point $o' \in L$ fixed by B . By choosing an appropriate basis, we can assume that for $\begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} \in U$ we have

$$\varphi \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} = \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array} \right] \in PGL(3).$$

The brackets indicate the class of the matrix in $PGL(3)$. All the lines passing through o' are stable by U . By a change of basis we can also assume that $\varphi(T)$ is diagonal. Then for φ to be a homomorphism, it is necessary that

$$\varphi \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} = \left[\begin{array}{ccc} \alpha^m & 0 & 0 \\ 0 & \alpha & \beta \\ 0 & 0 & \alpha^{-1} \end{array} \right] \in PGL(3), \quad m \in \mathbf{Z}.$$

For $m = -1 \pm c$, this gives two embeddings of B/Γ with $|\Gamma| = c$. The group B acts on L by the character $\alpha^{2 \pm c}$. There is another stable line

$\{(0:z_1:z_2) \mid z_i \in k\}$ on which B acts in the standard manner. This gives the two "ordinary" B/Γ -embeddings mentioned earlier for \mathbf{P}^2 .

CASE 2. U acts non-trivially on L .

(i) U acts trivially on the linear system S .

Then B stabilizes another line L' passing through o . Since we have that $\mathbf{P}^2 - \{L \cup L'\} \cong k \times k^* \cong B/\Gamma$, and since $k \times k^*$ contains no proper open subvariety isomorphic to itself, we must have that the complement to the open orbit is $Z = L \cup L'$. We will show that U acts trivially on L' . Indeed, let $x \in L' \setminus L$ and D be a line of \mathbf{P}^2 passing through x but not o , and let $u \in U$, $u \neq e$; then $uD \cap D$ is a point fixed by u since U acts trivially on S ; therefore it must belong to Z , but it is not in L ; thus it is in L' , hence it is x . So by exchanging L and L' , we are in Case 1.

(ii) U acts non-trivially on the linear system S .

Then T stabilizes a line L' in $S - L$.

Fix $u \in U$, $u \neq e$. We can choose a basis such that $\varphi(u)$ is in Jordan

normal form $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$. Now by a change of basis we can assume

$$\varphi \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} = \begin{bmatrix} 1 & 2\beta & \beta^2 \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{bmatrix} \in PGL(3).$$

Let S' be the linear system of conics passing through the point o . Now B acts on S' , and one can easily check that ${}^US'$, the set of conics stable by U is isomorphic to \mathbf{P}^1 . In fact it is the set of conics of the form

$$\{(z_0:z_1:z_2) \mid a(z_0z_2 - z_1^2) + bz_2^2 = 0\}, \quad (a:b) \in \mathbf{P}^1.$$

Also T acts on ${}^US'$; it must leave two conics invariant: the double line $L = \{(z_0:z_1:0)\}$ and a non-degenerate conic C . Since $\mathbf{P}^2 - \{L \cup C\}$ is isomorphic to $k \times k^*$, the complement to the open orbit is $L \cup C$. By a change of basis one can choose

$$C = \{(z_0:z_1:z_2) \mid z_0z_2 - z_1^2 = 0\} \quad \text{and} \quad L' = \{(z_0:0:z_1)\}.$$

By checking the action of T on $\mathbf{P}^2 - L$, one finds there is just one possibility which yields:

$$\varphi \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} = \begin{bmatrix} \alpha^2 & 2\alpha\beta & \beta^2 \\ 0 & 1 & \alpha^{-1}\beta \\ 0 & 0 & \alpha^{-2} \end{bmatrix} \in PGL(3) .$$

(So φ is obtained from the irreducible representation of $SL(2)$ of dimension 3.) This homomorphism gives rise to a B/Γ -embedding for $c = 4$. Note that there is exactly one fixed point: $(1:0:0)$. This is the “exceptional” embedding.

Embeddings into $\mathbf{P}^1 \times \mathbf{P}^1$:

The two projections $\mathbf{P}^1 \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ give the two different rulings of $\mathbf{P}^1 \times \mathbf{P}^1$. Any automorphism of $\mathbf{P}^1 \times \mathbf{P}^1$ either leaves the two rulings invariant or exchanges them. In other words,

$$\text{Aut}(\mathbf{P}^1 \times \mathbf{P}^1) = (PGL(2) \times PGL(2)) \rtimes \mathbf{Z}/2\mathbf{Z} .$$

Since B is connected, the image of $\varphi(B) \subset \text{Aut}(\mathbf{P}^1 \times \mathbf{P}^1)$ is connected; thus we consider homomorphisms $\varphi: B \rightarrow PGL(2) \times PGL(2)$. Up to conjugation, the only homomorphisms of B to $PGL(2)$ are

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \rightarrow \begin{bmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{bmatrix} \in PGL(2)$$

or
$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \rightarrow \begin{bmatrix} \alpha^m & 0 \\ 0 & 1 \end{bmatrix} \in PGL(2), \quad m = 0, 1, 2, \dots$$

To obtain an embedding, U cannot act trivially on $\mathbf{P}^1 \times \mathbf{P}^1$. So the possibilities (up to conjugation) are

$$\varphi \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} = \left\{ \begin{bmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{bmatrix}, \begin{bmatrix} \alpha^m & 0 \\ 0 & 1 \end{bmatrix} \right\} \in \text{Aut}(\mathbf{P}^1 \times \mathbf{P}^1), \quad m = 1, 2, 3, \dots$$

or

$$\varphi \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} = \left\{ \begin{bmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{bmatrix}, \begin{bmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{bmatrix} \right\} \in \text{Aut}(\mathbf{P}^1 \times \mathbf{P}^1) .$$

In the first case, we get an “ordinary” embedding of B/Γ with $c = m$ with two fixed points. The second induces a B/Γ -embedding with $c = 2$, and the complement to the open orbit consists of three curves isomorphic to \mathbf{P}^1 all intersecting transversely in the unique fixed point. This is the “exceptional” embedding.

Embeddings into $F_n, n \geq 1$:

Remember from section 1 that we can consider F_n as the union of E_n and the total space of the line bundle $\mathcal{O}_{\mathbf{P}^1}(n)$. Suppose we have a homomorphism $\varphi: B \rightarrow \text{Aut } F_n$ which gives rise to a B/Γ -embedding. Since $\text{Aut } F_n$ stabilizes E_n , we know that B fixes E_n . We consider three cases.

CASE 1. U acts trivially on E_n .

We will find $n + 1$ inequivalent "ordinary" embeddings of this type for each Γ .

In this case, consider the action of T on E_n . It cannot act trivially (because then each B -orbit would be contained in a fibre of $\pi_n: F_n \rightarrow \mathbf{P}^1$) and has therefore exactly two fixed points, x and y . By possibly exchanging x and y , we can assume that T acts by a character $\alpha^m, m > 0$ on $E_n \cong \mathbf{P}^1$ (i.e. for $z \in E_n - \{x, y\}$, we choose $x = \lim_{t \rightarrow 0} tz$ and $y = \lim_{t \rightarrow \infty} tz, t \in T$).

The fibres F_x and F_y of x and y , respectively, are stable by B . Let Z be the complement of the open orbit in F_n . Then we have $E_n \cup F_x \cup F_y \subset Z$. Since we know that $F_n - \{E_n \cup F_x \cup F_y\} \cong k \times k^* \cong B/\Gamma$, and since, as noted earlier, $k \times k^*$ contains no proper open subvariety isomorphic to itself, we must have $Z = E_n \cup F_x \cup F_y$.

Now by Lemma 1.3, we have $T \hookrightarrow B \rightarrow \text{Aut } F_n \rightarrow \text{Aff}(H^0(\mathbf{P}^1, \mathcal{O}(n)))$. Since T is reductive, T must fix a section D of $\mathcal{O}(n)$.

We also have that U acts on the space $H^0(\mathbf{P}^1, \mathcal{O}(n))$. Consider the orbit UD . First note that $UD \cong k$ (we could not have $UD = D$, because then D would be in the complement of the open orbit). Now let $u \in U, u \neq e$; then I claim that $uD \cap D \subset \{x', y'\}$, where $x' = F_x \cap D$ and $y' = F_y \cap D$. To see this, note that since U acts trivially on E_n , it stabilizes the fibres of π_n . Thus if z belongs to $uD \cap D$, then u belongs to the isotropy group of z , and therefore z must be in Z . The intersection number

$uD \cdot D$ is n ; so $UD \subset D \cup \bigcup_{p=0}^n A_p$, where A_p is the set of sections D' of $\mathcal{O}(n)$ such that $D \cap D' = px' + (n-p)y'$ counted with multiplicity. Now $D \cup A_p$ is isomorphic to $k, p = 0, \dots, n$; so $UD = D \cup A_p$ for some $p = 0, \dots, n$. We call p the *contact index* of the embedding. See Fig. 2.




FIGURE 2.

LEMMA 2.2. *Up to equivalence, there is at most one B/Γ -embedding into F_n of a given contact index p , with $p = 0, \dots, n$. Also, for such an embedding B acts on E_n by the character α^c , where c is the order of Γ .*

Proof. Suppose we have two B/Γ -embeddings into F_n with the same contact index p . Fix $u \in U$, $u \neq e$. For the first (resp. second) action denote by x, y (resp. \tilde{x}, \tilde{y}) the fixed points in E_n and D (resp. \tilde{D}) the section fixed by T . Set $D_u := uD$ (resp. $\tilde{D}_u := u\tilde{D}$).

Remember from section 1 we know that there is an exact sequence

$$1 \rightarrow k^* \rtimes H^0(\mathbf{P}^1, \mathcal{O}(n)) \rightarrow \text{Aut } F_n \rightarrow PGL(2) \rightarrow 1.$$

Since $PGL(2)$ acts doubly transitively on \mathbf{P}^1 , we can conjugate by an automorphism of F_n which sends x to \tilde{x} and y to \tilde{y} ; thus we can assume $x = \tilde{x}$ and $y = \tilde{y}$. Then by conjugating by an element of $H^0(\mathbf{P}^1, \mathcal{O}(n))$, which translates the sections, we can assume $D = \tilde{D}$. Finally, since the two embeddings have the same contact index, by conjugating by an automorphism that fixes the fibres and which is a homothety centered at D , we can assume $D_u = \tilde{D}_u$.

Now I claim that for a fixed Γ , there is at most one possible action of B on F_n which induces a B/Γ -embedding with the quadruple $\{x, y, D, D_u\}$. Indeed U acts by translation on each of the fibres of $\mathcal{O}(n)$; so D and D_u determine how U must act. Now check the action of T on D , which is the same as its action on E_n . Choose $z \in D$ in the open orbit. The order of the isotropy group B_z is c , the order of Γ , and $B_z \subset T$. So T acts on D by a character $\alpha^{\pm c}$. Since we chose x and y such that the action of T on E_n is given by a positive character, we must have that T acts on D by the character α^c . This proves the second statement of the lemma. Now let v be an element of the open orbit and $t \in T$. Choose $u \in U$ such that $(t^{-1}ut)v = v' \in D$. Then $tv = u^{-1}tv'$. So this fixes the action of T on the open orbit, which is dense in F_n . So the claim is true, and this finishes the proof of the lemma. \square

By this lemma, we have at most $n + 1$ inequivalent embeddings of this type for each Γ . Now we must show that these actually exist.

LEMMA 2.3. *Let n be a positive integer and p be an integer such that $0 \leq p \leq n$. Then for each finite $\Gamma \subset B$, there exists a B/Γ -embedding into F_n with contact index p .*

Proof. Let X_n be the surface obtained by contracting E_n in F_n as explained in section 1. Suppose we have an embedding of B/Γ into X_n

which fixes the vertex of the cone (if $n > 1$, this condition is always satisfied, because this point is singular). Then by blowing up the vertex, we obtain an embedding into F_n .

For each p with $0 \leq p \leq n$, we will exhibit an action of B on X_n which induces a B/Γ -embedding with contact index p . To do this we give a linear action of B on k^{n+2} which induces an action of B on \mathbf{P}^{n+1} stabilizing X_n and its vertex.

B acts on k^2 in the standard way:

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \begin{pmatrix} s \\ t \end{pmatrix} = \begin{pmatrix} \alpha s + \beta t \\ \alpha^{-1} t \end{pmatrix}.$$

Also for $i \in \mathbf{Z}$, we denote by (k, α^i) the vector space k with the action of B by the character α^i :

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} z = \alpha^i z.$$

Consider the B -module

$$k^2 \otimes (k, \alpha^{cp+1}) \oplus \bigoplus_{\substack{j=0 \\ j \neq p}}^n (k, \alpha^{cj}), \quad p = 0, \dots, n.$$

We have $B \rightarrow PGL(n+2) = \text{Aut } \mathbf{P}^{n+1}$ by

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \mapsto \begin{bmatrix} \alpha^{cp+2} & \alpha^{cp+1} & \beta & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & \alpha^{cp} & & & & & & \\ & & 1 & & & & 0 & \\ \cdot & & & \alpha^c & & & & \cdot \\ & & & & \cdot & & & \cdot \\ \cdot & & & & & \cdot & & \cdot \\ \cdot & 0 & & & & \cdot & \hat{\alpha}^{cp} & \cdot \\ & & & & & & \cdot & \cdot \\ & & & & & & & \cdot \\ 0 & & & & & & 0 & \alpha^{cn} \end{bmatrix}$$

We change the basis so that the image of $\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix}$ is

$$\begin{bmatrix} \alpha^{cp+2} & 0 & \dots & 0 & \alpha^{cp+1} & \beta & 0 & \dots & 0 \\ 0 & 1 & & & & & & & \\ . & & \alpha^c & & & & 0 & . & \\ . & & & \ddots & & & & . & \\ . & 0 & & & \alpha^{cp} & & & . & \\ & & & & & \ddots & & 0 & \\ 0 & & \dots & & & 0 & \alpha^{cn} & & \end{bmatrix}$$

Let X_n be as given in section 1. Clearly X_n and the vertex of the cone $(1:0:\dots:0)$ are fixed by this action. In X_n all the "fibres" are stable by U , and the two "fibres" $F_x = \{(z_0:z_1:0:\dots:0)\}$ and $F_y = \{(z_0:0:\dots:0:z_{n+1})\}$ are stable by B . It is easy to check that the isotropy group of $(0:1:\dots:1)$ is the finite subgroup of T of order c . So this induces an embedding of B/Γ into X_n which by blowing up the vertex gives a B/Γ -embedding into F_n where U acts trivially on E_n .

Let $D = \{(0:s^n:s^{n-1}t:\dots:t^n)\} \subset X_n$. Then D is a "section" stable by T . Fix $u = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in U$. Then $uD = \{(s^{n-p}t^p:s^n:s^{n-1}t:\dots:t^n)\} \subset X_n$. We check the multiplicity of the intersection of D and uD at $x' = (0:1:0:\dots:0)$. The local ring of x' in X_n is $k[z_0, t]_{(t, z_0)}$, and the local equation of D (resp. uD) is $z_0 = 0$ (resp. $z_0 = t^p$); thus this multiplicity is p , and the contact index of the embedding is p . This finishes the proof of the lemma. \square

Remark. By checking the induced torus actions on the fibres F_x and F_y , one finds the results about the structure of the action stated after Theorem 2.1.

CASE 2. U acts non-trivially on E_n and B fixes a section D of $\mathcal{O}(n)$.

We will find two "ordinary" embeddings of this type for each Γ .

In this case, U has one fixed point x on E_n . Then T must also fix x , and it also fixes another point $y \in E_n$. As before, we call Z the complement to the open orbit. Then we have $Z = E_n \cup D \cup F_x$, where F_x is the fibre of π_n containing x . Now look at the action of T on F_y , the fibre of y . Choose $z \in F_y$ in the open orbit. Then the order of the isotropy group B_z

is c , the order of Γ , and $B_z \subset T$. So T acts on F_y by the character $\alpha^{\pm c}$. For each such embedding, call this character the *sign* of the embedding. See Fig. 3.




FIGURE 3.

LEMMA 2.4. *Up to equivalence, there is at most one B/Γ -embedding into F_n with a given sign $\sigma = \alpha^{\pm c}$.*

Proof. Suppose we had two actions of B on F_n which yield two B/Γ -embeddings with the same sign σ . For the first (resp. second) action, let ψ (resp. $\tilde{\psi}$): $B \times E_n \rightarrow E_n$ be the induced action on E_n and D (resp. \tilde{D}) be the section of $\mathcal{O}(n)$ fixed by B .

Up to conjugacy there is only one action of B on $E_n \cong \mathbf{P}^1$ for which U acts non-trivially. So we can assume $\psi = \tilde{\psi}$. By conjugating by an appropriate automorphism of F_n which fixes the fibres and translates the sections, we can assume $D = \tilde{D}$.

Now I claim there is at most one action of B on F_n which yields a B/Γ -embedding with the triple $\{\psi, D, \sigma\}$. To see this, consider first the action of U on F_n . Now x is the fixed point of E_n , and F_x is its fibre. Let S be the set of sections of $\mathcal{O}(n)$ which are not D and intersect D with multiplicity n at the fixed point $x' = F_x \cap D$. This set is isomorphic to k^* (by the map $D' \rightarrow D' \cap F_y$) and is stable by B , so U acts trivially on S . Since the action of U on $D' \in S$ is identical to its action on E_n , the action of U on F_n is determined by ψ and D . As for the action of T , remember that T stabilizes the set S . The action on this set is equivalent to its action on F_y , the fibre of the point of E_n fixed by T and not fixed by U . This action is given by σ . So $\{\psi, D, \sigma\}$ determines the action of T on F_n . This proves the claim. \square

From this lemma, we see that for each Γ , there is at most two B/Γ -embeddings of this type. Now we must show that these embeddings actually exist.

LEMMA 2.5. Let Γ be a finite subgroup of B of order c and σ be $\alpha^{\pm c}$. Then there exists a B/Γ -embedding into F_n with sign σ .

Proof. We use the same notation as in Lemma 2.3. Consider the B -module

$$(k, \alpha^{-n \pm c}) \oplus S^n(k^2)$$

where $S^n(k^2)$ is the vector space of homogeneous polynomials of degree n over k with two variables, and the action of B on $S^n(k^2)$ is induced from the natural action on k^2 of B as a subgroup of $SL(2)$. We have $B \rightarrow PGL(n+2)$ by

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \mapsto \begin{bmatrix} \alpha^{-n \pm c} & 0 & \dots & 0 \\ 0 & & & \\ \cdot & \rho_n \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} & & \\ \cdot & & & \\ 0 & & & \end{bmatrix}$$

where ρ_n is the $(n+1)$ -dimensional irreducible matrix representation of $SL(2, k)$ corresponding to the basis $\left\{ \binom{n}{i} x^i y^{n-i} \right\}_{i=0, \dots, n}$ of $S^n(k^2)$.

As in Lemma 2.3, let $X_n = \{(z_0 : s^n : s^{n-1}t : \dots : t^n) \mid z_0, s, t \in k\} \subset \mathbf{P}^{n+1}$. Then X_n and its vertex $(1:0:\dots:0)$ are fixed by the action above. In X_n the "section" $\{(0:s^n:\dots:t^n)\}$ and the "fibre" $\{(z_0:z_1:0:\dots:0)\}$ are stable. The other "fibres" are not stable by U . The isotropy group of $(1:0:\dots:0:1)$ is the finite subgroup of T of order c . So this action gives an embedding of B/Γ into X_n which by blowing up the vertex gives an embedding into F_n where U acts non-trivially on E_n and B fixes a section.

The "fibre" $\{(z_0:0:\dots:0:z_{n+1})\}$ is stable by T and not by U . Also T acts on this fibre by the character $\alpha^{\pm c}$, so the sign of the embedding is $\alpha^{\pm c}$. This proves the lemma. \square

Remark. The group B acts on the fixed fibre of the B/Γ -embedding with sign $\alpha^{\pm c}$ by the character $\alpha^{2n \mp c}$. In particular, for each n , there is exactly one embedding of this type with $c = 2n$ where B acts trivially on the fixed fibre. We will use this remark for the following case.

CASE 3. U acts non-trivially on E_n and B does not fix any section of $\mathcal{O}(n)$.

For each n , we find one such case where $c = 2(n+1)$. These are the "exceptional" embeddings.

As in the previous case, B fixes one element $x \in E_n$. So Z , the complement to the open orbit, contains E_n and F_x , the fibre of x . Now $\mathbf{F}_n - \{E_n \cup F_x\}$ is isomorphic to $k \times k$; so Z must have another component. Suppose $z \in Z - \{E_n \cup F_x\}$; then $C = \overline{Bz}$ is contained in Z . Clearly C is a section of $\pi_n: \mathbf{F}_n \rightarrow \mathbf{P}^1$, and by hypothesis it is not a section of $\mathcal{O}(n)$; thus it is a section of π_n which intersects E_n at the point x . We have $Z = E_n \cup F_x \cup C$; since $\mathbf{F}_n - \{E_n \cup F_x \cup C\} \cong k \times k^*$.

LEMMA 2.6.

(i) Suppose $c = 2(n+1)$. Then there is exactly one embedding of B/Γ into \mathbf{F}_n of Case 3 with $C \cdot E_n = 1$. Also for this embedding there is a unique fixed point.

(ii) If $c \neq 2(n+1)$ there is no such embedding with $C \cdot E_n = 1$.

Proof. Recall from section 1 that one obtains \mathbf{F}_{n+1} from \mathbf{F}_n by blowing up a point x on E_n and contracting the strict transform of the fibre containing x .




FIGURE 4.

Now suppose we have such an embedding with $C \cdot E_n = 1$. We blow up the point x . (See Fig. 4.) Now there are three fixed points on the exceptional divisor F_0 , so B acts trivially on F_0 . Blow down \tilde{F}_x ; We obtain an embedding into \mathbf{F}_{n+1} as in Case 2, where B acts trivially on the fixed fibre. As we have seen in the remark of Case 2, this happens in exactly one case with $c = 2(n+1)$. Conversely, given this embedding into \mathbf{F}_{n+1} , by doing the reverse procedure, one obtains exactly one embedding of this type. (By changing the fixed point which is blown up first one obtains an equivalent embedding.) This proves everything except the unicity of the fixed point.

Now we exhibit explicitly the embedding of (i). We use the notation of Lemmas 2.3 and 2.5. Consider the B -module $S^{n+1}(k^2)$. We have $B \rightarrow PGL(n+2)$ by

$$\begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \mapsto \left[\rho_{n+1} \begin{pmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{pmatrix} \right]$$

where ρ_{n+1} is the $(n+2)$ -dimensional irreducible representation of $SL(2, k)$. Consider the closure of the orbit of $x^{n+1} + y^{n+1}$ by B using the basis $\left\{ \binom{n+1}{i} x^i y^{n+1-i} \right\}_{i=0, \dots, n+1}$. This is exactly

$$X_n = \{ (z_0 : s^n : s^{n-1}t : \dots : t^n) \mid z_0, s, t \in k \}.$$

The vertex $(1:0:\dots:0)$ is fixed by this action. The two stable curves in X_n are the "fibre" $\{(z_0 : z_1 : 0 : \dots : 0)\}$ and $\{(s^{n+1} : s^n t : \dots : t^{n+1})\}$, the image of the $(n+1)$ -uple embedding of \mathbf{P}^1 in \mathbf{P}^{n+1} . It is easy to see that the isotropy group of $(1:0:\dots:0:1)$ is the finite subgroup of T of order c ; so this action gives a B/Γ -embedding into X_n which induces an embedding into F_n . Since the only fixed point on X_n is the vertex and there is only one fixed "fibre", we have exactly one fixed point for the action on F_n . It is easily checked that the intersection number of E_n with the other stable section in F_n is 1. Thus the lemma is proven. \square

LEMMA 2.7. *Any embedding of Case 3 must have $C \cdot E_n = 1$.*

Proof. The intersection number $C \cdot E_n = p$ is strictly positive. Suppose that $p > 1$. Now blow up x and then contract the strict transform of F_x ; we obtain an embedding into F_{n+1} . Let C_1 be the strict transform of C in F_{n+1} ; then the intersection number $C_1 \cdot E_{n+1}$ is $p - 1$. Also, this new embedding has at least two fixed points: one on E_{n+1} and the other the image of the strict transform of F_x in F_{n+1} . By doing this process $p - 1$ times, we get an embedding into F_{n+p-1} of Case 3 with $C_{p-1} \cdot E_{n+p-1} = 1$ and at least two fixed points. By Lemma 2.6 this is impossible. Therefore we must have $p = 1$. (See Fig. 5.) \square




FIGURE 5.

This finishes Case 3. Thus we know all the embeddings into \mathbf{P}^2 , $\mathbf{P}^1 \times \mathbf{P}^1$ and \mathbf{F}_n , $n \geq 1$. The comments after Theorem 2.1 are easily verified by checking each embedding. This finishes the proof of the theorem. \square

Remarks.

(1) Note that — as to be expected — all the embeddings into \mathbf{F}_1 are obtained by blowing up the embeddings into \mathbf{P}^2 at fixed points.

(2) The “exceptional” embeddings, i.e. those with only one fixed point, are of special interest, because this phenomenon does not occur for smooth complete embeddings of tori. (See [KKMS] for a reference on torus embeddings.)

§ 3. APPLICATION TO $SL(2)$ -EMBEDDINGS

In [LV] a combinatorial method is presented in order to classify all normal $SL(2)$ -embeddings. A natural question is how to classify those which are smooth and complete to obtain a *geometrical* realization. We now sketch how the result of this article is useful for this. (For further details see [JM].)

Given a B/Γ -embedding X , we construct an $SL(2)/\Gamma$ -embedding in the following way. Consider the B -action on $SL(2) \times X$ given by

$$b \cdot (s, x) = (sb^{-1}, bx)$$

where $b \in B$, $s \in SL(2)$, and $x \in X$. Denote by $SL(2)*_BX$ the variety obtained by quotienting by this action. The action of $SL(2)$ on this variety by left multiplication endows it with the structure of an $SL(2)/\Gamma$ -embedding. The projection $SL(2) \times X \rightarrow SL(2)$ induces a locally trivial fibre bundle $SL(2)*_BX \xrightarrow{p} SL(2)/B \cong \mathbf{P}^1$. The morphism p is $SL(2)$ -equivariant, and the fibre of p is B -isomorphic to X . So we see that for studying the geometry of the $SL(2)/\Gamma$ -embeddings of this form it is useful to study the B/Γ -embeddings.

As for general $SL(2)/\Gamma$ -embeddings one finds the following essential result. Let Γ be a finite cyclic subgroup of $SL(2)$. Let V be a smooth $SL(2)/\Gamma$ -embedding with orbit Y . Then there exists a Borel subgroup B of $SL(2)$ containing Γ and an $SL(2)$ -stable open neighborhood of Y in V which is of the form $SL(2)*_BX$ for some smooth B/Γ -embedding X . Thus all smooth $SL(2)/\Gamma$ -embeddings are *locally* of the form above. Also any smooth B/Γ -embedding can be completed to a smooth embedding. Thus it is enough to study the complete ones.

We can use this fact, for example, to study blow-ups of orbits, since blowing up is a local property. Thus we can find the minimal $SL(2)/\Gamma$ -embeddings. This is done in [JM], Chapter IV, for $\Gamma = \{e\}$ and $\Gamma = \{\pm e\}$.

REFERENCES

- [Beau] BEAUVILLE, A. *Surfaces Algébriques Complexes*. Soc. Math. de France, (Astérisque 54) Paris, 1978.
- [Bor] BOREL, A. *Linear Algebraic Groups*. W. A. Benjamin, New York, 1969.
- [Ful] FULTON, W. *Intersection Theory*. Springer-Verlag, Berlin, Heidelberg, 1984.
- [G-H] GRIFFITHS, P. and J. HARRIS. *Principles of Algebraic Geometry*. Wiley and Sons, New York, 1978.
- [Gro] GROTHENDIECK, A. Torsion, homologie et sections rationnelles. *Séminaire Chevalley, Anneaux de Chow et Applications*, Secrétariat Math., Paris, 1958.
- [Har] HARTSHORNE, R. *Algebraic Geometry*. Springer-Verlag, New York, 1977.
- [H-O] HUCKLEBERRY, A. and E. OELJEKLAUS. *Classification Theorems for Almost Homogeneous Spaces*. Revue de l'Institut Elie Cartan 9, Université de Nancy, 1984.
- [JM] JAUSLIN-MOSER, L. *Normal Embeddings of $SL(2)/\Gamma$* . Thesis, Université de Genève, Geneva, 1987.
- [Kam] KAMBAYASHI, T. Projective Representation of Algebraic Linear Groups of Transformations. *Amer. J. Math.* 88 (1966), 199-205.
- [KKMS] KEMPF, G., F. KNUDSEN, D. MUMFORD and B. SAINT-DONAT. *Toroidal Embeddings 1*. Lect. Notes in Math. #339, Springer-Verlag, 1974.
- [LV] LUNA, D. et Th. VUST. Plongements d'espaces homogènes. *Comment. Math. Helv.* 58 (1983), 186-245.
- [Mum] MUMFORD, D. *Algebraic Geometry I, Complex Projective Varieties*. Springer-Verlag, New York, 1976.
- [O-W] ORLIK, P. and Ph. WAGREICH. Algebraic Surfaces with k^* -actions. *Acta Mathematica* 138 (1977), 43-81.
- [Pop] POPOV, V. L. Classification of Affine Algebraic Surfaces that are Quasi-homogeneous with respect to an Algebraic Group. *Izv. Akad. Nauk SSSR, Ser. Mat.* 37 (No. 5) (1973), 1039-1055.
- [Pot] POTTERS, J. On Almost Homogeneous Compact Complex Analytic Surfaces. *Inv. Math.* 8 (1969), 244-266.
- [Saf] ŠAFAREVIČ, I. R. *Algebraic Surfaces*. Proceedings of the Steklov Institute of Math. No. 75, 1965.

(Reçu le 23 février 1988)

Lucy Moser-Jauslin

Section de Mathématiques
 Université de Genève
 Case postale 240
 CH-1211 Genève 24
 (Suisse)