

# 4.1. Results

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **30.04.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## §4. ESTIMATES FOR FOURIER COEFFICIENTS OF SIEGEL CUSP FORMS

### 4.1. RESULTS

Very recently, it has turned out that Jacobi forms can be used in a rather simple way to prove growth estimates for Fourier coefficients of Siegel cusp forms of genus 2. The bounds one obtains in this way, in fact, are somewhat better than those obtained previously by different methods.

Let  $F$  be a Siegel cusp form of integral weight  $k$  on  $\Gamma_2$  and let  $a(T)$  be its Fourier coefficients. The classical Hecke argument immediately gives

$$(7) \quad a(T) \ll_F (\det T)^{k/2} .$$

If one applies a classical theorem of Landau [25, 32] to the Rankin-Dirichlet series

$$\sum_{\{T > 0\}/GL_2(\mathbf{Z})} |a(T)|^2 (\det T)^{-s}$$

where the summation extends over a complete set of representatives for the usual left-action of  $GL_2(\mathbf{Z})$  on the set of positive definite symmetric half-integral  $(2, 2)$ -matrices  $T$ , one can sharpen (7) and show that

$$a(T) \ll_{\varepsilon, F} (\det T)^{k/2 - 3/32 + \varepsilon} \quad (\varepsilon > 0) .$$

(Recall that Landau's theorem roughly speaking asserts that if a Dirichlet series has a meromorphic continuation to  $\mathbf{C}$  and satisfies an appropriate functional equation, then one can deduce a “good” upper bound for the growth of its coefficients.) For details we refer to [5] and also [11] where the argument is slightly different; note that the authors prove an estimate for arbitrary genus  $n$ .

Let us mention the following

**THEOREM 1** (Kitaoka [16]). *Suppose that  $k$  is even. Then*

$$a(T) \ll_{\varepsilon, F} (\det T)^{k/2 - 1/4 + \varepsilon} \quad (\varepsilon > 0) .$$

The proof of Theorem 1 uses Poincaré series of exponential type on  $\Gamma_2$  and estimates for generalized matrix-argument Kloosterman sums and can be viewed as a generalization to genus 2 of a well-known method how to obtain “good” bounds for the Fourier coefficients of elliptic cusp forms.

Let us explain now briefly how Jacobi forms can be brought into play (for full details cf. [20, 21]). Let  $\phi \in J_{k, m}^{\text{cusp}}$  with Fourier coefficients  $c(n, r)$ . Then for  $k > 2$  one shows that

$$(8) \quad c(n, r) \ll_{\varepsilon, k} (m + |D|^{1/2+\varepsilon})^{1/2} \frac{|D|^{k/2-3/4}}{m^{(k-1)/2}} \|\phi\| \quad (\varepsilon > 0)$$

where  $D := r^2 - 4mn$  and the bound in  $\ll$  only depends on  $\varepsilon$  and  $k$ .

For the proof one carries over the method of Poincaré series and Kloosterman sums from the one-variable situation already mentioned above to the case of the Jacobi group. Note that Poincaré series on  $\Gamma_1^J$  were studied in [14, II, § 2]. The Kloosterman sums that occur in their Fourier coefficients can be related to Salié sums and therefore can easily be estimated (a similar phenomenon happens in the case of modular forms of half-integral weight, cf. [15]). The proof of (8) for  $D$  a fundamental discriminant (i.e. the discriminant of a quadratic field) is given in [20, § 1] and for arbitrary  $D$  is given in [21, § 1].

On the other hand, if one applies Landau's theorem to the Dirichlet series  $D_{F, F}(s)$  discussed in § 3, one finds that

$$(9) \quad \|\phi_m\| \ll_{\varepsilon, F} m^{k/2-2/9+\varepsilon} \quad (\varepsilon > 0).$$

The estimates (8) and (9) now imply the following

**THEOREM 2** [20, 21]. *One has*

$$(10) \quad a(T) \ll_{\varepsilon, F} (\det T)^{k/2-13/36+\varepsilon} \quad (\varepsilon > 0).$$

In fact, both sides of (10) are invariant under  $T \mapsto U' TU$  ( $U \in GL_2(\mathbf{Z})$ ), hence if in (10) we write  $T = \begin{pmatrix} n & r/2 \\ r/2 & m \end{pmatrix}$ , then we may assume that  $m = \min T$ , where  $\min T$  denotes the least positive integer represented by  $T$ . If we use (8) and (9) together with the fact that  $\min T \ll (\det T)^{1/2}$  which is well-known from reduction theory, we obtain (10).

#### 4.2. PROBLEMS

- i) In [15], Iwaniec using some sophisticated arguments for certain sums of Salié sums showed that the Fourier coefficients  $a(n)$  ( $n \in \mathbf{N}$ ) of a cusp form  $f$  of weight  $k - 1/2$  for  $k > 0$  and  $n$  squarefree satisfy

$$a(n) \ll_K \sigma_0(n) (\log 2n)^2 n^{k/2-15/28} \|f\|,$$

where  $\sigma_0(n)$  is the number of positive divisors of  $n$  and  $\|f\|$  is the appropriately normalized Petersson norm of  $f$ . We wonder if it is possible to prove an analogous estimate for the Fourier coefficients  $c(n, r)$  ( $D = r^2 - 4mn$  a fundamental discriminant) of a function  $\phi \in J_{k, m}^{\text{cusp}}$  for  $k > 2$  which also is