

Systole and $2g-2$ of closed hyperbolic surfaces of genus g

Autor(en): **Mondal, Sugata**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **60 (2014)**

PDF erstellt am: **29.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-515843>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Systole and λ_{2g-2} of closed hyperbolic surfaces of genus g

Sugata MONDAL

Abstract. We apply topological methods to study eigenvalues of the Laplacian on closed hyperbolic surfaces. For any closed hyperbolic surface S of genus g , we get a geometric lower bound on $\lambda_{2g-2}(S)$: $\lambda_{2g-2}(S) > 1/4 + \epsilon_0(S) > 0$, where $\epsilon_0(S)$ is an explicit constant which depends only on the *systole* of S .

Mathematics Subject Classification (2010). Primary 30F; Secondary 35P05, 35P15.

Keywords. Hyperbolic surfaces, eigenfunctions.

1. Introduction

Here a *hyperbolic surface* is a complete two-dimensional Riemannian manifold with curvature equal to -1 . Any hyperbolic surface is isometric to a quotient \mathbb{H}/Γ , where \mathbb{H} is the Poincaré upper halfplane and Γ is a *Fuchsian group*, i.e. a discrete torsion-free subgroup of $\mathrm{PSL}(2, \mathbb{R})$. The *Laplacian* on \mathbb{H} is the differential operator which associates to a C^2 -function f the function

$$\Delta f(z) = y^2 \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} \right).$$

It induces a differential operator on $S = \mathbb{H}/\Gamma$ which extends to a self-adjoint operator Δ_S densely defined on $L^2(S)$. Its domain is the Sobolev space $\mathcal{H}(S)$ consisting of the functions $\phi \in L^2(S)$ whose gradient in the sense of distributions is a measurable vector field which satisfies $\int_S \|\nabla \phi\|^2 dv < \infty$. The Laplacian is a non-positive operator whose spectrum is contained in a smallest interval $(-\infty, -\lambda_0(S)] \subset \mathbb{R}^- \cup \{0\}$ with $\lambda_0(S) \geq 0$. The *Rayleigh quotients* allow us to characterize the *bottom of the spectrum* of S :

$$\lambda_0(S) = \inf \frac{\int_S \|\nabla \phi\|^2 dv}{\int_S \phi^2 dv},$$

where the infimum is taken over all non-constant smooth functions ϕ with compact support. Recall that the bottom of the spectrum on \mathbb{H} is $\lambda_0(\mathbb{H}) = 1/4$ (cf. [Cha], p. 46, Theorem 5).

Definition 1.1. Let $\lambda > 0$. A non-zero function $f: S \rightarrow \mathbb{R}$ is a λ -eigenfunction if $f \in L^2(S)$ and satisfies $\Delta f + \lambda f = 0$. One calls λ an *eigenvalue*. When $0 < \lambda \leq 1/4$, λ is called a *small eigenvalue* and f is called a *small eigenfunction*.

When S is a compact hyperbolic surface, the *spectrum* of S is a discrete set:

$$0 = \lambda_0(S) < \lambda_1(S) \leq \lambda_2(S) \leq \dots \leq \lambda_n(S) \leq \dots$$

where in the above sequence each number is repeated according to its multiplicity as an eigenvalue and $\lambda_i(S)$ denotes the i -th non-zero eigenvalue of S for $i \geq 1$.

Definition 1.2. For a hyperbolic surface S the *systole* $s(S)$ of S is defined to be the minimum of the lengths of closed geodesics on S .

The main result of this paper is

Theorem A. *Let S be a closed hyperbolic surface of genus g . Then there exists an explicit constant $\epsilon_0(S) > 0$, which depends only on the systole of S , such that $\lambda_{2g-2}(S) > 1/4 + \epsilon_0(S)$.*

We recall some general facts about the behavior of λ_{2g-2} as a function on \mathcal{M}_g , the moduli space of closed hyperbolic surfaces of genus g . Any eigenvalue λ_i , in particular λ_{2g-2} , is a continuous function on \mathcal{M}_g (see for instance [C-C]). The moduli space \mathcal{M}_g is the space of all closed hyperbolic surfaces of genus g up to isometry. Recall that the set $\mathcal{I}_\epsilon = \{S \in \mathcal{M}_g : s(S) \geq \epsilon\}$ is compact ([Bu], p. 163). By [O-R], we have $\lambda_{2g-2}(S) > 1/4$ for all $S \in \mathcal{M}_g$. Hence there exists a non-zero constant $\eta(\epsilon)$ such that $\lambda_{2g-2}(S) > 1/4 + \eta(\epsilon)$ for all $S \in \mathcal{I}_\epsilon$. This proves the theorem with $\epsilon_0(S) = \eta(s(S))$. The content of Theorem A is to make this constant explicit in terms of the geometry of S . We shall see that we can take $\epsilon_0(S)$ to be any positive number smaller than

$$\min \left\{ \frac{1}{4(g-1)}, \frac{1}{4} \left(\left(\frac{\cosh \rho_0}{\sinh \rho_0} \right)^2 - 1 \right) \right\}$$

where $2s(S)\sinh \rho_0 = |S|$, the area of the surface S . Observe that $\frac{1}{4(g-1)} = \frac{\pi}{|S|}$ and $\frac{1}{4} \left(\left(\frac{\cosh \rho_0}{\sinh \rho_0} \right)^2 - 1 \right) = \frac{s(S)^2}{|S|^2}$.

We now sketch briefly the proof of the above theorem. It uses topological methods as in [O-R]. First we recall that an open subset of a surface S is called *incompressible* if the fundamental group of any of its connected components maps injectively into $\pi_1(S)$. It is plain that simply connected open subsets of S are

incompressible. Let \mathcal{E}_λ denote the eigenspace of the Laplacian on S for the eigenvalue λ . For $\epsilon > 0$, let $\mathcal{E}^{\frac{1}{4}+\epsilon}$ be the direct sum of eigenspaces \mathcal{E}_λ with $\lambda \leq \frac{1}{4} + \epsilon$. For $f \neq 0 \in \mathcal{E}^{\frac{1}{4}+\epsilon}$, define the *nodal set* $\mathcal{Z}(f)$ as $f^{-1}(0)$. It is proved in [O-R], using the analyticity of eigenfunctions on \mathbb{H} , that $\mathcal{Z}(f)$ is the union of a finite graph and a discrete set. Let $\mathcal{G}(f)$ be the subgraph of $\mathcal{Z}(f)$ obtained by suppressing those connected components which are homotopic to a point on S (equivalently, those which are contained in a topological disc). Due to this modification, each component of $S \setminus \mathcal{G}(f)$ is incompressible. One of the main observations in [O-R] was that for any $f \neq 0 \in \mathcal{E}^{\frac{1}{4}}$, the Euler characteristic of at least one component of $S \setminus \mathcal{G}(f)$ is strictly negative. For $\epsilon > 0$ there is no reason, in general, to believe such a result for $f \neq 0 \in \mathcal{E}^{\frac{1}{4}+\epsilon}$. However, we will prove the following

Lemma 1.3. *Let S be a closed hyperbolic surface of genus g . Then there exists an explicit constant $\epsilon_0(S) > 0$ depending only on the genus g and the systole of S , such that for any $f \neq 0 \in \mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}$, the Euler characteristic of at least one component of $S \setminus \mathcal{G}(f)$ is strictly negative.*

Let Σ be a Riemannian surface. Let $\Omega \subseteq \Sigma$ be an open set such that the closure $\overline{\Omega}$ is a submanifold with piecewise smooth boundary. Then denote by Δ the Laplace operator of Σ restricted to Ω . *Dirichlet eigenvalues* of Ω are the λ 's such that the problem

$$\begin{cases} \Delta u = \lambda u & \text{on } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

admits a non-zero solution u , continuous on $\overline{\Omega}$ and smooth on Ω . The smallest λ for which such a solution exists is denoted by $\lambda_0(\Omega)$ and is called *the first Dirichlet eigenvalue* of Ω . This number can be defined in terms of Rayleigh quotients in a similar way as the bottom of the spectrum of Ω :

$$\lambda_0(\Omega) = \inf \frac{\int_{\Omega} \|\nabla \phi\|^2 dv}{\int_{\Omega} \phi^2 dv},$$

where the infimum is taken over all non-zero smooth functions ϕ with compact support in Ω . From this characterization it is evident that for any two submanifolds Ω_1 and Ω_2 as above with compact closure, we have $\lambda_0(\Omega_1) \geq \lambda_0(\Omega_2)$ when $\Omega_1 \subsetneq \Omega_2$. The above lemma will be deduced from the following

Proposition 1.4. *Let S be a closed hyperbolic surface of genus g . Let $\Omega \subset S$ be a surface with smooth boundary which is homeomorphic either to a disc or to an annulus. Then there exists a constant $\epsilon(\Omega) > 0$, depending on the length l_Ω of the geodesic in S homotopic to a generator of $\pi_1(\Omega)$ and on the area of Ω , such that the first Dirichlet eigenvalue of Ω satisfies $\lambda_0(\Omega) > 1/4 + \epsilon(\Omega)$. Furthermore there exists an explicit constant $\epsilon_0(S) > 0$ depending only on the systole of S such that $\epsilon(\Omega) > \epsilon_0(S)$.*

Notation 1.5. For any surface $\Omega \subseteq S$ with smooth boundary, $|\Omega|$ denotes the area of Ω for the area measure on S , and $L(\partial\Omega)$ denotes the length of the boundary of Ω .

We shall see in the proof that $\epsilon(\Omega)$ is a strictly decreasing function of $|\Omega|$ when l_Ω is kept fixed and a strictly increasing function of l_Ω when $|\Omega|$ is kept fixed. The statement in the proposition then follows from the observation that both parameters, i.e. $|\Omega|$ and l_Ω , are bounded: the former being bounded above by $4\pi(g-1)$ and the latter being bounded below by $s(S)$.

The proof of the above proposition depends mainly on two geometric inequalities: the Faber–Krahn isoperimetric inequality and Cheeger’s inequality. The scheme of the proof of Theorem A then follows the same lines as that of Theorem 1 in [O-R].

Existence of surfaces with small eigenvalues was proved originally by Randol [R1]. We shall recall a construction of P. Buser [Bu] for the construction of such surfaces. The construction is carried out by first considering a genus g hyperbolic surface admitting a pair of pants decomposition with very short boundary geodesics, then constructing an orthogonal family of functions with small Rayleigh quotient. The number of functions in that family is exactly $(2g-2)$. This gives the existence of at least $(2g-3)$ small eigenvalues (which is the maximum possible number by [O-R]).

After proving Theorem A in §2, in the subsequent parts of the paper we study the behavior of λ_i as a function on the moduli space \mathcal{M}_g . We recall that the moduli space \mathcal{M}_g is the space of all closed hyperbolic surfaces of genus g up to isometry. We focus our interest on the first $2g-2$ non-zero eigenvalues. Theorem A (or even a continuity argument on \mathcal{M}_g) implies one direction of the following

Claim 1.6. *For a family S_n of closed hyperbolic surfaces in \mathcal{M}_g , $\lambda_{2g-2}(S_n)$ tends to $1/4$ if and only if the systole $s(S_n)$ tends to zero.*

The other direction follows from a construction due to P. Buser [Bu], as we shall see in §3.

The above proposition can be compared with the following result of Schoen, Wolpert and Yau [S-W-Y]. Let M be a closed oriented surface of genus g

with a metric of (possibly variable) Gaussian curvature K . For an integer $n \geq 1$ consider the family \tilde{C}_n of curves on M which are a disjoint union of simple closed geodesics and which divide M into $n+1$ components (necessarily $n \leq 2g-3$). Define a number l_n by

$$l_n = \min\{L(C) : C \in \tilde{C}_n\},$$

where $L(C)$ denotes the length of C .

Theorem 1.7 (Schoen–Wolpert–Yau). *Suppose that for some constant $k > 0$ we have $-1 \leq K \leq -k$. Then there exist positive constants α_1, α_2 depending only on g such that for $1 \leq n \leq 2g-3$, we have $\alpha_1 k^{3/2} l_n \leq \lambda_n \leq \alpha_2 l_n$ and $\alpha_1 k \leq \lambda_{2g-2} \leq \alpha_2$.*

Recall that the *Bers constant* β , see [B], which depends only on g , has the property that $l_{2g-2} < \beta$. So this theorem implies that λ_{2g-2} is bounded above by a constant depending only on g . Observe also that the construction of Buser ([Bu], Theorem 8.1.3) leads to the same conclusion. Namely by Buser's construction for any $\delta > 0$ there exists a constant $\epsilon > 0$ such that $\lambda_{2g-2} < \frac{1}{4} + \delta$ for any $S \in \mathcal{M}_g$ with $s(S) < \epsilon$. Since λ_{2g-2} is a continuous function on \mathcal{M}_g and $\mathcal{I}_\epsilon = \{S \in \mathcal{M}_g : s(S) \geq \epsilon\}$ is compact the existence of an upper bound is clear. In this context we would like to mention a paper due to Dodziuk, Pignataro, Randol and Sullivan [D-P-R-S] where the authors obtained results similar to that of [S-W-Y] in the context of arbitrary (non-compact) hyperbolic surfaces.

In §3 we will study the behavior of $\lambda_i(S)$ as $s(S)$ tends to zero. More precisely, let $\overline{\mathcal{M}}_g$ denote the compactification of \mathcal{M}_g obtained by adding the moduli spaces of (not necessarily connected) non-compact finite area hyperbolic surfaces with area equal to $4\pi(g-1)$. Let $\partial\mathcal{M}_g = \overline{\mathcal{M}}_g \setminus \mathcal{M}_g$ be the corresponding boundary of \mathcal{M}_g . We study the behavior of $\lambda_i(S_n)$ when $S_n \in \mathcal{M}_g$ tends to a point in $\partial\mathcal{M}_g$. By Theorem 1.7 and the above discussion, it is clear that $\lambda_i(S)$ is bounded above for all $S \in \mathcal{M}_g$ and for $1 \leq i \leq 2g-2$. Indeed the method using Buser's construction works for any i , showing that λ_i is bounded by a constant depending only on g and i . So for any i we can consider the set

$$V_i = \left\{ \lim_{n \rightarrow \infty} \lambda_i(S_n) : (S_n) \text{ is a sequence in } \mathcal{M}_g \text{ converging to a point in } \partial\mathcal{M}_g \text{ such that } \lim_{n \rightarrow \infty} \lambda_i(S_n) \text{ exists} \right\}.$$

With this notation, the above claim says that $V_{2g-2} = \{\frac{1}{4}\}$. Using a result of Courtois and Colbois [C-C], we will also prove the following assertion:

Claim 1.8. *For any $1 \leq i \leq 2g-3$, there exists a $\Lambda_i(g) \in (0, \frac{1}{4}]$ such that V_i contains the interval $[0, \Lambda_i(g)]$.*

In §4 we study non-compact hyperbolic surfaces of finite area. Recall that for a non-compact hyperbolic surface S of finite area, the spectrum of the Laplace operator is composed of two parts: the discrete part and the continuous part. The continuous part covers the interval $[1/4, \infty)$ and is spanned by Eisenstein series. The discrete part is the union of the *residual spectrum* and *cuspidal spectrum*. The residual spectrum is a finite set contained in the interval $(0, 1/4)$ and it corresponds to poles of the analytic continuation of Eisenstein series. The cuspidal spectrum consists of those eigenvalues whose associated eigenfunctions tend to zero uniformly near any cusp. The number of cuspidal eigenvalues is known to be infinite for arithmetic groups [I]. The cuspidal eigenvalues can possibly appear anywhere in the interval $(0, \infty)$. Denote by $\lambda^c_i(S)$ the i -th *cuspidal eigenvalue* of S .

In analogy to Theorem A, one can investigate the following conjecture.

Conjecture 1. *Let S be a finite area hyperbolic surface of type (g, n) , i.e. topologically equivalent to a genus g surface with n punctures. Then there exists an explicit constant $\epsilon_0(S) > 0$, depending only on the systole of S , such that $\lambda^c_{2g-2+n}(S) > 1/4 + \epsilon_0(S)$.*

This would be an extension of a result of Jean-Pierre Otal and Eulalio Rosas (Theorem 2 in [O-R]). However our methods do not suffice to settle this conjecture. In this connection we state the following conjecture of Otal and Rosas in [O-R] which is motivated by [O], Propositions 2 and 3.

Conjecture 2. *Let S be a finite area hyperbolic surface of type (g, n) . Then $\lambda^c_{2g-2}(S) > 1/4$.*

Now we consider a finite area hyperbolic surface S of type (g, n) . Denote by $\mathcal{T}_{g,n}$ the *Teichmüller space* of all marked hyperbolic surfaces of type (g, n) . For any choice of pair of pants decomposition of S one can define a system of coordinates on $\mathcal{T}_{g,n}$, the *Fenchel–Nielsen coordinates* which consist, for each curve in the pants decomposition, of the length of that curve and a *twist* parameter along that curve ([Bu], Chapter 6). Now we consider the set $\mathcal{T}_{g,n}^0$ of all hyperbolic surfaces in $\mathcal{T}_{g,n}$ for which all twist parameters are equal to zero. Each surface in $\mathcal{T}_{g,n}^0$ carries an involution ι which when restricted to each pair of pants is the orientation reversing involution that fixes the boundary components. This involution induces an involution on each eigenspace of the Laplacian. The eigenfunctions corresponding to the eigenvalue -1 are called *antisymmetric* and the corresponding eigenvalue is called an *antisymmetric eigenvalue*. We denote the i -th antisymmetric cuspidal eigenvalue of $S \in \mathcal{T}_{g,n}^0$ by $\lambda^{o,c}_i(S)$.

Theorem B. *For any $S_0 \in \mathcal{T}_{g,n}^0$ there exists an explicit constant $\epsilon_0(S_0) > 0$, depending only on the systole of the surface S_0 , such that $\lambda^{o,c}_g(S_0) > \frac{1}{4} + \epsilon_0(S_0)$.*

As a matter of fact, the constant $\epsilon_0(S)$ can be taken equal to any number below

$$\min \left\{ \frac{1}{2(2g-2+n)}, \frac{1}{4} \left(\left(\frac{\cosh \rho_0}{\sinh \rho_0} \right)^2 - 1 \right) \right\},$$

where $2s(S) \sinh \rho_0 = |S|$.

Acknowledgements. I would like to express my sincere gratitude to my advisor Jean-Pierre Otal whose encouragements, kindness and patience were significant ingredients in the work.

2. Geometric lower bound on $\lambda_{2g-2}(S)$

2.1. Proof of Proposition 1.4. Suppose first that $\Omega \subseteq S$ is a disc or more generally a domain such that $\pi_1(\Omega)$ maps to zero in $\pi_1(S)$. Then choose an isometric lift of Ω to \mathbb{H} , still denoted by Ω . We will use the Faber–Krahn inequality ([Cha], p. 87) in the following form.

Theorem 2.1 (Faber–Krahn inequality). *Let $\Omega \subseteq \mathbb{H}$ be a domain such that $\partial\Omega$ is smooth. Let D be a geodesic disc in \mathbb{H} with same area as Ω , i.e. $|\Omega| = |D|$. Then*

$$\lambda_0(\Omega) \geq \lambda_0(D),$$

with equality if and only if Ω is isometric to D .

Let $B(t)$ be the geodesic disc in \mathbb{H} with radius t . The geodesic disc with same area as Ω has radius $t(\Omega) = 2 \sinh^{-1}(\frac{|\Omega|}{4\pi})$. By the Faber–Krahn inequality $\lambda_0(B(t(\Omega))) \leq \lambda_0(\Omega)$.

Since Ω is contained in S whose area equals $2\pi(2g-2)$, by the Gauss–Bonnet theorem, $|\Omega| < 2\pi(2g-2)$. Therefore, $B(t(\Omega))$ is contained in the disc with radius $t_0 = 2 \sinh^{-1}(g-1)$. Recall that for two subsurfaces D_1 and D_2 in \mathbb{H} with compact closure, $\lambda_0(D_1) > \lambda_0(D_2)$ when $D_1 \subsetneq D_2$. Thus $\lambda_0(B(t))$ is a strictly decreasing function of t . Hence $\lambda_0(B(t(\Omega))) > \lambda_0(B(t_0))$. Now by Theorem 5 in [Cha], we have

$$\lambda_0(B(t)) > \lim_{s \rightarrow \infty} \lambda_0(B(s)) = \frac{1}{4}.$$

Hence we finally have a strictly positive $\epsilon_1(|\Omega|)$ which depends only on the area $|\Omega|$ of Ω such that $\lambda_0(B(t(\Omega))) = \frac{1}{4} + \epsilon_1(|\Omega|)$. Since $\lambda_0(B(t))$ is a strictly decreasing function of t , $\epsilon_1(|\Omega|)$ is a strictly decreasing function of $|\Omega|$ which is bounded below by the constant $\epsilon_1(S) = \lambda_0(B(t_0)) - \frac{1}{4}$.

Suppose now that Ω is an annulus and that the image of $\pi_1(\Omega)$ in $\pi_1(S)$ is a non-trivial cyclic subgroup $\langle \tau \rangle$. Let \mathbb{T} denote the cylinder $\mathbb{H}/\langle \tau \rangle$. Let γ denote the core geodesic of \mathbb{T} and l the length of γ . Then l is the length of the shortest geodesic of S homotopic to a generator of $\pi_1(\Omega)$. Consider an isometric lift of the annulus Ω to $\mathbb{H}/\langle \tau \rangle$, still denoted by Ω . We need to prove that $\lambda_0(\Omega) > \frac{1}{4} + \epsilon_0(S)$ where $\epsilon_0(S)$ depends only on l and $|\Omega|$. We will use Cheeger's inequality ([Cha], p. 95) in the following form:

Theorem 2.2 (Cheeger's inequality). *Let $\Omega \subsetneq \mathbb{T}$ be a submanifold with piecewise smooth boundary. Let $h(\Omega)$ be the Cheeger constant of Ω . Then*

$$\lambda_0(\Omega) \geq \frac{h^2(\Omega)}{4}.$$

Recall that the Cheeger constant of Ω is equal to $\inf\{\frac{L(\partial V)}{|V|}\}$, where V ranges over all compact submanifolds of Ω with smooth boundary.

The proof of Proposition 1.4 in the case of an annulus follows from Cheeger's inequality and the next lemma.

Lemma 2.3. *Let $\Omega \subsetneq \mathbb{T}$ be a submanifold with piecewise smooth boundary and let $h(\Omega)$ be the Cheeger constant of Ω . Then we have*

$$h(\Omega) > 1 + \epsilon_2(|\Omega|, l),$$

for some $\epsilon_2(|\Omega|, l) > 0$ which depends only on the area of Ω and on the length l of the core geodesic of \mathbb{T} .

Proof. First we observe that the Cheeger constant is bounded below by the quantity $\inf\{\frac{L(\partial V)}{|V|}\}$ where V ranges over connected submanifolds of Ω . Secondly, this infimum is the same when V ranges over all discs or *essential* annuli contained in Ω . Recall that an annulus is *essential* when it is not homotopically trivial in \mathbb{T} . This is because any connected, compact submanifold $V \subseteq \Omega$ is diffeomorphic either to a disc with some discs removed or to an essential annulus with some discs removed. In both cases, taking the union of V with those removed discs, one obtains a submanifold V' which is either a disc or an essential annulus which satisfies $L(\partial V') \leq L(\partial V)$ and $|V'| \geq |V|$. Therefore $\frac{L(\partial V')}{|V'|} \leq \frac{L(\partial V)}{|V|}$.

Suppose now that $V \subseteq \Omega$ is diffeomorphic to a disc. By the isoperimetric inequality ([B-Z], p. 11), one has $(\frac{L(\partial V)}{|V|})^2 \geq 1 + \frac{4\pi}{|V|}$. Therefore if $V \subseteq \Omega$ then $(\frac{L(\partial V)}{|V|})^2 \geq 1 + \frac{4\pi}{|\Omega|}$. Since $|V| < 2\pi(2g - 2)$, we get $(\frac{L(\partial V)}{|V|})^2 > 1 + \frac{1}{g-1}$.

Now we suppose that $V \subseteq \Omega$ is an essential annulus. In order to prove the claim in this case we will need the following notion of *symmetrization*, which is close to the notion of *Steiner symmetrization* ([H], p. 18).

Definition 2.4. Let $V \subseteq \mathbb{T}$ be an essential annulus. The *symmetrization* of V is the annulus $V_0 \subseteq \mathbb{T}$ symmetric with respect to γ with constant width and which has the same area as V .

Recall that the *Fermi coordinates* on \mathbb{T} assign to each point p the pair $(s, r) \in \{\gamma\} \times \mathbb{R}$, where r is the signed distance of p from γ , and s is the point of γ nearest to p . After parametrizing the geodesic γ by arc-length, these coordinates provide a diffeomorphism between \mathbb{T} and $\mathbb{R}/l\mathbb{Z} \times \mathbb{R}$. The hyperbolic metric in these coordinates equals $dr^2 + \cosh r^2 ds^2$.

Lemma 2.5. Let $V \subseteq \mathbb{T}$ be an essential annulus with piecewise smooth boundary and V_0 be the symmetrization of V . Then $L(\partial V) \geq L(\partial V_0)$.




FIGURE 1

Proof. First we consider the case when each component of ∂V is a graph over γ . By that we mean that there exist two functions r_1 and $r_2: [0, l] \rightarrow \mathbb{R}$ such that r_i is a piecewise smooth map (there is a partition $0 = s_1 < s_2 < \dots < s_m = l$ such that each restriction $r_i|_{[s_j, s_{j+1}]}$ is smooth) with $r_i(0) = r_i(l)$ and the components of ∂V are parametrized in Fermi coordinates as $\{(s, r_i(s)), s \in [0, l]\}$ for $i = 1, 2$. Then the components of the symmetrization V_0 of V are the graphs of the constant functions $r_3 = \rho$ and $r_4 = -\rho$ with $\rho = \sinh^{-1}(\frac{|V|}{2l})$. Up to exchanging r_1 and r_2 , we may suppose that $r_1(s) > r_2(s)$ for all $0 \leq s \leq l$. Then we calculate the areas of V and V_0 :

$$|V| = \int_0^l \int_{r_1(s)}^{r_2(s)} \cosh r \, dr \, ds = \int_0^l \{\sinh r_2(s) - \sinh r_1(s)\} \, ds$$

and

$$|V_0| = \int_0^l \int_{-\rho}^{\rho} \cosh r \, dr \, ds = \int_0^l 2 \sinh \rho \, ds = 2l \sinh \rho.$$

The length of ∂V_0 is

$$L(\partial V_0) = 2l \cosh \rho$$

and the length of ∂V satisfies

$$\begin{aligned} L(\partial V) &= \int_0^l \{\dot{r}_1(s)^2 + 1\}^{1/2} \cosh r_1(s) ds + \int_0^l \{\dot{r}_2(s)^2 + 1\}^{1/2} \cosh r_2(s) ds \\ &\geq \int_0^l \{\cosh r_1(s) + \cosh r_2(s)\} ds. \end{aligned}$$

Call L_0 the constant equal to the last expression. Observe that $L(\partial V) = L_0$ if and only if $\dot{r}_1(\theta) = 0 = \dot{r}_2(\theta)$. This implies that r_1 and r_2 are constants.

One has

$$L(\partial V)^2 - |V|^2 \geq (L_0 + |V|)(L_0 - |V|).$$

Now,

$$\begin{aligned} L_0 + |V| &= \int_0^l ((\cosh r_2(s) + \sinh r_2(s)) + (\cosh r_1(s) - \sinh r_1(s))) ds \\ &= \int_0^l (\exp(r_2(s)) + \exp(-r_1(s))) ds \end{aligned}$$

and similarly

$$L_0 - |V| = \int_0^l (\exp(-r_2(s)) + \exp(r_1(s))) ds.$$

Thus we have

$$\begin{aligned} &(L_0 + |V|)(L_0 - |V|) \\ &= \left(\int_0^l (\exp(r_2(s)) + \exp(-r_1(s))) ds \right) \\ &\quad \times \left(\int_0^l (\exp(-r_2(s)) + \exp(r_1(s))) ds \right) \\ &\geq \left(\int_0^l (\exp(r_2(s)) + \exp(-r_1(s)))^{\frac{1}{2}} (\exp(-r_2(s)) + \exp(r_1(s)))^{\frac{1}{2}} ds \right)^2 \\ &\quad \text{(by the Cauchy-Schwarz inequality)} \\ &= \left(\int_0^l (2 + 2 \cosh(r_1(s) + r_2(s)))^{\frac{1}{2}} ds \right)^2. \end{aligned}$$

Since $\cosh x \geq 1 \quad \forall x$, we get $(L_0 + |V|)(L_0 - |V|) \geq 4l^2 = L(\partial V_0)^2 - A(V_0)^2$. Equality holds if and only if r_1, r_2 are independent of s and if $r_1 = -r_2$.

Since by construction $|V| = |V_0|$, the lemma is proven when V is an annulus whose boundary components are graphs over γ .

Now we consider the case of an arbitrary annulus with piecewise smooth boundary. By approximation, it suffices to prove Lemma 2.5 for those V which satisfy the following property: there exists a partition of γ :

$$0 = s_1 < s_2 < \cdots < s_k = l = 0$$

such that over each interval $[s_i, s_{i+1}]$, ∂V is a union of the graphs of finitely many functions. We consider now such an annulus. We consider the strip over $[s_i, s_{i+1}]$ in \mathbb{T} which is diffeomorphic to $[s_i, s_{i+1}] \times \mathbb{R}$ in Fermi coordinates. Denote by V^i the intersection of V with this strip. For $1 \leq i \leq k$, we denote by f_j , $j = 0, 1, 2, \dots, l(i)$ the boundary curves of V^i i.e. in Fermi coordinates the components of ∂V^i are parametrized as $\{(s, f_j(s)): s \in [s_i, s_{i+1}]\}$ for $j = 0, 1, 2, \dots, l(i)$ and for any $s \in [s_i, s_{i+1}]$, $r(f_0(s)) > r(f_1(s)) > \cdots > r(f_{l(i)}(s))$. Now we calculate the area of V^i :

$$\begin{aligned} |V^i| &= \sum_{j=l(i)-1, l(i)-3, \dots, 1} \int_{s_i}^{s_{i+1}} \int_{f_j(s)}^{f_{j+1}(s)} \cosh r \, dr \, ds \\ &= \sum_{j=1}^{l(i)} \int_{s_i}^{s_{i+1}} (-1)^{j+1} \sinh f_j(s) \, ds. \end{aligned}$$

The length of ∂V^i is given by

$$L(\partial V^i) = \sum_{j=1}^{l(i)} \int_{s_i}^{s_{i+1}} \{\dot{f}_j(s)^2 + 1\}^{1/2} \cosh f_j(s) \, ds \geq \int_{s_i}^{s_{i+1}} \sum_{j=1}^{l(i)} \cosh f_j(s) \, ds.$$

Call $L_0(i)$ the constant equal to the last expression and calculate

$$\begin{aligned} L(\partial V)^2 - |V|^2 &= \left(\sum_i L(\partial V^i) \right)^2 - \left(\sum_i |V^i| \right)^2 \\ &\geq \left(\sum_i L_0(i) \right)^2 - \left(\sum_i |V^i| \right)^2 \\ &= \left(\sum_i \int_{s_i}^{s_{i+1}} \sum_{j=1}^{l(i)} \exp[(-1)^{j+1} f_j(s)] \, ds \right) \\ &\quad \times \left(\sum_i \int_{s_i}^{s_{i+1}} \sum_{j=1}^{l(i)} \exp[(-1)^j f_j(s)] \, ds \right) \end{aligned}$$

$$\begin{aligned}
&\geq \left(\sum_i \int_{s_i}^{s_{i+1}} (\exp[(-1)^{0+1} f_0(s)] + \exp[(-1)^{1+1} f_1(s)]) ds \right) \\
&\quad \times \left(\sum_i \int_{s_i}^{s_{i+1}} (\exp[(-1)^0 f_0(s)] + \exp[(-1)^1 f_1(s)]) ds \right) \\
&\geq \left(\int_0^l (2 + 2 \cosh(f_1(s) - f_0(s))^{\frac{1}{2}}) ds \right)^2 \\
&\quad \text{(by Cauchy-Schwarz)} \\
&\geq 4l^2 = L(\partial V_0)^2 - |V_0|^2.
\end{aligned}$$

Hence using the same argument as before we finally prove Lemma 2.5. \square

So now we have $(\frac{L(\partial V)}{|V|}) \geq (\frac{L(\partial V_0)}{|V_0|}) = \frac{\cosh \rho}{\sinh \rho}$ where $|V| = 2l \sinh \rho$. Thus we conclude the proof of Lemma 2.3 by taking

$$\epsilon_2(\Omega, l) = \frac{1}{2} \min \left\{ \frac{\cosh \theta}{\sinh \theta} - 1, \left(1 + \frac{4\pi}{|\Omega|} \right)^{\frac{1}{2}} - 1 \right\}$$

where $|\Omega| = 2l \sinh \theta$. \square

Since $\frac{\cosh \rho}{\sinh \rho}$ is a strictly decreasing function of ρ we have

$$\left(\frac{L(\partial V)}{|V|} \right) \geq \frac{\cosh \rho_1}{\sinh \rho_1} > \frac{\cosh \rho_0}{\sinh \rho_0}$$

where $2l \sinh \rho_1 = |\Omega|$ and $2s(S) \sinh \rho_0 = |S| = 4\pi(g-1)$ (since $V \subseteq \Omega \subsetneq S$). To conclude the proof of Proposition 1.4 we take

$$\epsilon_0(S) = \frac{1}{2} \min \left\{ \epsilon_1(S), \frac{1}{4(g-1)}, \frac{1}{4} \left(\left(\frac{\cosh \rho_0}{\sinh \rho_0} \right)^2 - 1 \right) \right\}. \quad \square$$

Remark 2.6. From the expression of $\epsilon_0(S)$ we observe that if (S_n) is a sequence in \mathcal{M}_g , then $\epsilon_0(S_n)$ tends to zero only if $s(S_n)$ tends to zero. The computations in the proposition also show that for any $\Omega \subseteq S$ diffeomorphic to a disc or to an annulus one has

$$\lambda_0(\Omega) \geq \frac{1}{4} + 2\epsilon_0(S).$$

2.2. Proof of Theorem A. The proof at this point follows the same lines as that of $\lambda_{2g-2}(S) > \frac{1}{4}$ in [O-R] and we refer to [O-R] for the details. We take $\epsilon_0(S)$ as in Proposition 1.4. Consider the space $\mathcal{E}^{\frac{1}{4} + \epsilon_0(S)}$. Recall that \mathcal{E}^λ is the direct sum of the eigenspaces of the Laplacian with eigenvalues less than or equal to λ . Let $f \neq 0 \in \mathcal{E}^{\frac{1}{4} + \epsilon_0(S)}$. The *nodal set* $\mathcal{Z}(f)$ of f is defined as $f^{-1}(0)$. Recall

that $\mathcal{G}(f)$ is the subgraph of $\mathcal{Z}(f)$ obtained by suppressing those connected components which are zero homotopic on S . Each component of $S \setminus \mathcal{G}(f)$ is an open surface, possibly equal to S when $\mathcal{G}(f)$ is empty. The sign of f on a component of $S \setminus \mathcal{G}(f)$ can be defined as follows. There is a finite collection of disjoint closed topological discs (D_j) with $\partial D_j \cap \mathcal{Z}(f) = \emptyset$ such that each component of $\mathcal{Z}(f)$ which is zero homotopic is contained in one of the D_j 's. Therefore each component of $S \setminus \mathcal{G}(f)$ is a union of a component of $S \setminus \mathcal{Z}(f)$ with a finite number of those D_j 's. Define the sign of f on such a component to be the one of f on the corresponding component of $S \setminus \mathcal{Z}(f)$. Now we denote the union of all components with positive, resp. negative, sign as $C^+(f)$, resp. $C^-(f)$. As a consequence of the construction, the surfaces $C^+(f)$ and $C^-(f)$ are incompressible. As recalled earlier, an open subset of a surface S is called *incompressible* if the fundamental group of any of its connected components maps injectively into $\pi_1(S)$. The union of the connected components of $C^+(f)$, resp. $C^-(f)$, which are neither discs nor rings is denoted by $S^+(f)$, resp. $S^-(f)$. The surfaces $S^\pm(f)$ may be empty or disconnected but by construction when they are nonempty, they are incompressible.

Denote the Euler characteristic of $S^+(f)$, resp. $S^-(f)$, by $\chi^+(f)$, resp. $\chi^-(f)$. (We use the convention that the Euler characteristic of the empty set is zero.) The incompressibility property of $S^+(f)$ and $S^-(f)$ gives that $\chi^+(f) + \chi^-(f)$ is greater than $\chi(S)$. By definition, we have $\chi^\pm(f) \leq 0$ with equality only if $S^\pm(f)$ is empty.

Lemma 2.7. *The Euler characteristic of at least one component of $S \setminus \mathcal{G}(f)$ is negative.*

Proof. Let us suppose by contradiction that for some $f \neq 0 \in \mathcal{E}^{\frac{1}{4} + \epsilon_0(S)}$, each component S_i , $1 \leq i \leq m$ of $S \setminus \mathcal{G}(f)$ has non-negative Euler characteristic. So, each such component is homeomorphic either to an open disc or to an open annulus. Since $f \in \mathcal{E}^{\frac{1}{4} + \epsilon_0(S)}$ the Rayleigh quotient of f , $R(f)$ is $\leq \frac{1}{4} + \epsilon_0(S)$. Therefore, since $\mathcal{G}(f)$ has measure zero, for at least one component, say S_1 , one has

$$R(f|_{S_1}) = \frac{\int_{S_1} \|\nabla f\|^2}{\int_{S_1} f^2} \leq \frac{1}{4} + \epsilon_0(S).$$

Now we shall calculate the Rayleigh quotient $R(f|_{S_1})$ and show that our choice of $\epsilon_0(S)$ leads to a contradiction.

Let us assume that S_1 is homeomorphic to an open disc. The case when S_1 is an annulus can be dealt with similarly. Since f is smooth and $f|_{\partial S_1} = 0$ we can choose, by Sard's theorem, a sequence (ϵ_n) of regular values of f converging to 0. Then the level set

$$\{x \in S_1: f(x) = \epsilon_n\}$$

is a smooth submanifold of S for n large enough. Furthermore one of the components of this level set confines a domain $D_n \subsetneq S_1$ homeomorphic to a closed disc with smooth boundary such that $S_1 \setminus D_n$ has arbitrarily small area. Now we consider the Rayleigh quotient, $R(f_n|_{D_n})$ of the function $f_n = f - \epsilon_n$ restricted to the region D_n . This function vanishes on ∂D_n . As ϵ_n converges to 0, $R(f_n|_{D_n})$ converges to $R(f|_{S_1})$. Thus for any $\delta > 0$, in particular for $\epsilon_0(S)/2$, we can find ϵ_n small enough such that

$$R(f_n|_{D_n}) \leq \frac{1}{4} + \epsilon_0(S) + \frac{\epsilon_0(S)}{2} < \frac{1}{4} + 2\epsilon_0(S).$$

Now since D_n is a closed disc with smooth boundary which is contained in $S_1 \subseteq S$, it follows from the Rayleigh quotient characterization of the first Dirichlet eigenvalue of D_n that $R(f_n|_{D_n}) \geq \lambda_0(D_n)$. By Remark 2.6 we have

$$\lambda_0(D_n) \geq \frac{1}{4} + 2\epsilon_0(S).$$

This is a contradiction when n is sufficiently large. \square

So some component of $S \setminus \mathcal{G}(f)$ has negative Euler characteristic. This component is a component of $S^\pm(f)$. Thus we obtain

$$\chi^+(f) + \chi^-(f) < 0.$$

Now we start with some definitions and complete the proof.

Definition 2.8. According to the sign of f on S_i , we denote this component as $S_i^+(f)$ or $S_i^-(f)$. For each such surface with negative Euler characteristic, we consider a *compact core*, i.e. a compact surface $K_i^\pm(f) \subset S_i^\pm(f)$ such that the inclusion is a homotopy equivalence. We then define the surface $\Sigma^+(f)$, resp. $\Sigma^-(f)$, as the union of the compact cores $K_i^+(f)$, resp. $K_i^-(f)$, and of those components (if any) of the complement $S \setminus \bigcup K_i^+(f)$, resp. $S \setminus \bigcup K_i^-(f)$, which are annuli. Therefore, $\Sigma^+(f)$, resp. $\Sigma^-(f)$, is obtained from $\bigcup K_i^+(f)$, resp. $\bigcup K_i^-(f)$, by adding (if any) the annuli between the components of $\bigcup K_i^+(f)$, resp. $\bigcup K_i^-(f)$. We call $\Sigma(f) = \Sigma^+(f) \cup \Sigma^-(f)$ the *characteristic surface* of f , while $\Sigma^+(f)$, resp. $\Sigma^-(f)$, is called the positive, resp. negative, characteristic surface of f . The definition of these surfaces depends uniquely on the choice of compact cores and those are well defined up to isotopy. By construction the Euler characteristic of $\Sigma^+(f)$, resp. $\Sigma^-(f)$, is $\chi^+(f)$, resp. $\chi^-(f)$. It is clear that $\Sigma^+(-f) = \Sigma^-(f)$ and $\Sigma^-(-f) = \Sigma^+(f)$.

2.3. Proof of Theorem A (continued). Let m denote the dimension of the space $\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}$. Theorem A will follow from the inequality $m \leq (2g-2)$. Let $\mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ denote the unit sphere of $\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}$ (for some arbitrary norm)

and let $\mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ be the projective space of $\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}$ i.e. the quotient of $\mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ by the involution $f \rightarrow -f$.

For each integer i with $2 - 2g \leq i \leq -1$, we denote

$$C_i = \{f \in \mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}) : \chi^+(f) + \chi^-(f) = i\}.$$

According to the lemma and its consequence above, $\mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}) = \bigcup_{2-2g}^{-1} C_i$. On the other hand, each C_i is invariant under the antipodal involution. Let P_i be the quotient of C_i under the antipodal involution. The projective space $\mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ is the union of the sets P_i .

Lemma 2.9. *For any integer i such that $2 - 2g \leq i \leq -1$, the covering map $C_i \rightarrow P_i$ is trivial.*

Proof. Let $f \in C_i$. We use the notation introduced in the definition of the characteristic surface of f : $S_i^\pm(f)$ is a connected component of negative Euler characteristic of $S^\pm(f)$ and $K_i^\pm(f)$ is a compact core of $S_i^\pm(f)$. We may assume that the compact core has been chosen in such a way that any connected component of $\mathcal{Z}(f)$ that is contained in some $S_i^\pm(f)$ is indeed contained in the interior of the corresponding $K_i^\pm(f)$.

For any function $g \in \mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}$ close enough to f , and for each i , $K_i^\pm(f)$ is contained in a component $S_i^\pm(g)$ of $S^\pm(g)$. Fix a neighborhood $V(f)$ of f in $\mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ such that these inclusions occur on each surface $K_i^\pm(f)$.

We will show that for any $g \in C_i \cap V(f)$, the characteristic surfaces $\Sigma^+(f)$ and $\Sigma^+(g)$, resp. $\Sigma^-(f)$ and $\Sigma^-(g)$, are isotopic. Choose the compact cores $K_i^\pm(g)$ of surfaces $S^\pm(g)$ so that when $K_i^\pm(f)$ is contained in $S_j^\pm(g)$, it is also contained in the interior of $K_j^\pm(g)$. Now observe that if two components of the boundaries of surfaces $K_j^+(f)$ are homotopic in S then the homotopy between them is achieved by an annulus contained in $\Sigma^+(f)$, by the definition of the characteristic surface. Since this annulus joins two curves of $K_j^+(g)$ by definition of the characteristic surface again, it is contained in one of the connected components $\Sigma^+(g)$ too.

We deduce from this that each connected component of $\Sigma^\pm(f)$ is contained in a connected component of $\Sigma^\pm(g)$ (of the same sign). Since $\Sigma^+(f)$ and $\Sigma^-(f)$ are incompressible in S , they are incompressible in $\Sigma^+(g)$ and $\Sigma^-(g)$ respectively. In particular, their Euler characteristics satisfy

$$\chi^+(f) \leq \chi^+(g) \quad \text{and} \quad \chi^-(f) \leq \chi^-(g);$$

these inequalities can be equalities if and only if the surfaces $\Sigma^+(f)$ and $\Sigma^+(g)$, resp. $\Sigma^-(f)$ and $\Sigma^-(g)$, are isotopic. But since $g \in C_i$, we have

$$\chi^+(f) + \chi^-(f) = i = \chi^+(g) + \chi^-(g).$$

Thus $\Sigma^+(f)$ and $\Sigma^+(g)$ are isotopic. The same holds for $\Sigma^-(f)$ and $\Sigma^-(g)$.

Since the *isotopy class* of $\Sigma^+(f)$ and the *isotopy class* of $\Sigma^-(f)$ are locally constant on C_i , they are constant on each connected component of C_i . Finally we observe that the functions f and $-f$ cannot be in the same connected component of C_i , for otherwise $\Sigma^+(f)$ and $\Sigma^-(f)$ would be isotopic. But two disjoint and incompressible surfaces of negative Euler characteristic contained in S cannot be isotopic. Thus the covering map in Lemma 2.9 is trivial. \square

2.4. Completion of the proof of Theorem A. We conclude the proof of Theorem A following a method of B. Sévenec [Se]. The double covering $\mathbb{S}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}) \rightarrow \mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ is associated to a cohomology class $\beta \in H^1(\mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)}), \mathbb{Z}/2\mathbb{Z})$. Each covering $C_i \rightarrow P_i$ is described by the Čech cohomology class, $\beta|_{P_i}$. Since each of these coverings is trivial, we have $\beta|_{P_i} = 0$. Since $\mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$ is the union of P_i and since there are at most $2g-2$ of them, we have $\beta^{2g-2} = 0$; see [Se], Lemma 8. Since β has order m in the $\mathbb{Z}/2\mathbb{Z}$ -cohomology ring of $\mathbb{P}(\mathcal{E}^{\frac{1}{4}+\epsilon_0(S)})$, we have $m \leq 2g-2$. \square

3. Systole and the Laplace spectrum

In this section we study the eigenvalues of the Laplacian as functions on the moduli space. Recall that the *moduli space* \mathcal{M}_g is the space of all closed hyperbolic surfaces of genus g up to isometry. \mathcal{M}_g can be compactified to a space $\overline{\mathcal{M}}_g$ by adding the moduli spaces of (not necessarily connected) non-compact finite area hyperbolic surfaces with area equal to $4\pi(g-1)$. In this compactification a sequence (S_n) in \mathcal{M}_g , with $s(S_n) \rightarrow 0$, converges to $S_\infty \in \mathcal{M}_{g_0, n_0}$ (with $2g_0-2+n_0=2g-2$) if and only if for any given $\epsilon > 0$ the ϵ -thick part $(S_n^{[\epsilon, \infty)})$ converges to $S_\infty^{[\epsilon, \infty)}$ in the Gromov–Hausdorff topology. Recall that the ϵ -thick part of a surface S is the subset of those points of S where the injectivity radius is at least ϵ . Recall also that the *injectivity radius* of a point $p \in S$ is the radius of the largest geodesic disc that can be embedded in S with center p .

It is a classical result that for any i , λ_i is a continuous function on \mathcal{M}_g (see for instance [C-C]). It is also shown in [C-C] that eigenvalues less than $1/4$ are continuous up to $\partial\mathcal{M}_g$. Let (S_n) be a sequence of surfaces in \mathcal{M}_g which tends to $S_\infty \in \partial\mathcal{M}_g = \overline{\mathcal{M}}_g \setminus \mathcal{M}_g$.

Theorem 3.1 ([C-C] and [He]). *Let $\lambda(S_n)$ be a sequence of eigenvalues of S_n which converges to $\lambda < 1/4$. Then λ is an eigenvalue of S_∞ and up to extracting a subsequence and possibly multiplying by a scaling constant the corresponding eigenfunctions on S_n converge to an eigenfunction on S_∞ uniformly over compact subsets.*

Our situation is a bit different. For a fixed i we shall study the behavior of $\lambda_i(S_n)$ when $S_n \in \mathcal{M}_g$ tends to a point in $\partial\mathcal{M}_g$. Recall

$$V_i = \left\{ \lim_{n \rightarrow \infty} \lambda_i(S_n) : (S_n) \text{ is a sequence in } \mathcal{M}_g \text{ converging to a point in } \partial\mathcal{M}_g \text{ such that the limit exists} \right\}.$$

In [R3] Randol showed a limiting behavior of λ_{2g-2} over some special family. Now we apply Theorem A to prove Claim 1.6 in the following form:

Claim 3.2. $\lambda_{2g-2}(S_n)$ tends to $\frac{1}{4}$ if and only if $s(S_n)$ tends to zero. In particular $V_{2g-2} = \{\frac{1}{4}\}$.

Proof. By Theorem A, if $\lambda_{2g-2}(S_n)$ tends to $\frac{1}{4}$ then $\epsilon_0(S_n)$ tends to zero. For the other direction we use Buser's construction. By the definition of the systole, there is a closed geodesic τ on S such that the length of τ is equal to $s(S)$. Now from the *Collar Theorem* (cf. [Bu]) of L. Keen [K] (see also [R2]) and the explicit computations in [Bu], p. 219, we see that for any $\epsilon > 0$ and any $i \geq 1$ we have $\delta > 0$ such that whenever $s(S) < \delta$, we can find at least i disjoint annuli in the collar neighborhood of τ such that the first Dirichlet eigenvalue of each of the annuli is $\leq \frac{1}{4} + \epsilon$. The corresponding eigenfunctions are orthogonal. Hence we have $\lambda_{i-1}(S) \leq \frac{1}{4} + \epsilon$. Therefore using Theorem A for an $i \geq 2g-1$ we obtain the convergence $\lambda_{i-1}(S_n) \rightarrow \frac{1}{4}$. \square

Now we show that such a limiting behavior is not true in general for $i \leq 2g-3$. More precisely, we shall prove Claim 1.8, rephrased as follows:

Claim 3.3. For any $1 \leq i \leq 2g-3$, there exists $\Lambda_i(g)$, $0 < \Lambda_i(g) \leq \frac{1}{4}$ such that $V_i = [0, \Lambda_i(g)]$.

Before starting the proof we recall the definition of the *Teichmüller space*, \mathcal{T}_g . It is the space of all marked closed hyperbolic surfaces of genus g . Let $S \in \mathcal{T}_g$. Given a pair of pants decomposition of S , we have a coordinate system on \mathcal{T}_g , the Fenchel–Nielsen coordinates. \mathcal{M}_g is the quotient of \mathcal{T}_g by the action of Mod_g , the *Teichmüller modular group*. Since Mod_g acts properly discontinuously on \mathcal{T}_g , $\mathcal{T}_g \rightarrow \mathcal{M}_g$ is a ramified topological covering. Thus the pre-composition of this covering map with λ_i yields a map, also denoted by λ_i , from \mathcal{T}_g to \mathbb{R} . We shall use the same notation for a point in \mathcal{T}_g and its image in \mathcal{M}_g .

Proof of Claim 3.3. We shall prove the claim for $i = 1$. The proof for $1 \leq i \leq 2g-3$ is similar. We choose a pair of pants decomposition \mathcal{P} of an $S \in \mathcal{T}_g$ and consider the corresponding Fenchel–Nielsen coordinates $(l^{\mathcal{P}}_j, \theta^{\mathcal{P}}_j)$ on \mathcal{T}_g . Here $l^{\mathcal{P}}_j$'s denote the *length* coordinates and $\theta^{\mathcal{P}}_j$'s denote the *twist* coordinates (cf. [Bu]). We fix two geodesics γ and γ' among the boundary geodesics of the pants decomposition \mathcal{P} . Thus the length functions l_γ and $l_{\gamma'}$

respectively of γ and γ' are among $l^{\mathcal{P}}_j$'s. Suppose that the pants decomposition is chosen in such a way that γ is non-separating and γ' is separating.

First we prove that V_i is not empty. From a construction of P. Buser [Bu], Theorem 8.1.3, it follows that if $0 < \delta < \frac{1}{24}$ then $\lambda_{2g-3}(S) < \frac{1}{4}$ for any $S \in \mathcal{T}_g$ with $l^{\mathcal{P}}_j(S) < \delta$ for all j (the number $\frac{1}{24}$ has no particular significance other than ensuring this condition). We fix one such δ and consider one $M \in \mathcal{T}_g$ such that $l^{\mathcal{P}}_j(M) < \delta$ for all j . Now consider a sequence of surfaces $(S_n) \in \mathcal{T}_g$ such that $(l^{\mathcal{P}}_j, \theta^{\mathcal{P}}_j)(S_n) = (l^{\mathcal{P}}_j, \theta^{\mathcal{P}}_j)(M)$ for all $(l^{\mathcal{P}}_j, \theta^{\mathcal{P}}_j)$ except l_γ and the $l_\gamma(S_n)$ coordinate decreases to zero as n goes to infinity. Then (S_n) converges to a point $S_\infty \in \partial\mathcal{M}_g$. By our choice of δ (for M) and since the number of components of S_∞ is exactly one, it follows from [C-C], Theorem 0.1, that $0 < \lim_{n \rightarrow \infty} \lambda_1(S_n) = \lambda_1(S_\infty) = p < \frac{1}{4}$. Now consider another sequence (S'_n) , constructed in the same way as (S_n) except that we vary the coordinate $l_{\gamma'}$ instead of l_γ . In this case the limiting surface of the sequence (S'_n) has two components. So using [C-C] again $\lim_{n \rightarrow \infty} \lambda_1(S'_n) = 0$. Thus we see that 0 and p are in V_1 , proving that V_1 is not empty.

Next we prove that whenever some $0 < c \leq \frac{1}{4}$ is in V_1 , the whole interval $(0, c]$ is contained in V_1 . Since c is in V_1 we have a sequence (P_n) in \mathcal{M}_g such that $\lim_{n \rightarrow \infty} \lambda_1(P_n) = c$. Up to extracting a subsequence, we may assume that (P_n) converges to $P_\infty \in \partial\mathcal{M}_g$. Then P_∞ is a finite area connected (since $c > 0$) non-compact hyperbolic surface of type (g', m) (where $g' + \frac{m}{2} = g$). For some marking of P_n , there is a pants decomposition of S , $\gamma_1, \dots, \gamma_k, \dots, \gamma_{3g-3}$ such that $\gamma_1, \dots, \gamma_k$ are exactly those curves on P_n whose lengths tend to zero. Consider the corresponding Fenchel–Nielsen coordinates $(l_i, \theta_i)_{i=1,2,\dots,3g-3}$ on \mathcal{T}_g . These coordinates induce coordinates on $\mathcal{T}_{g',m}$ which will be denoted by the same notation. In these coordinates we can choose representatives of P_n in \mathcal{T}_g such that $(l_i^n, \theta_i^n)(P_n)$ converges to $(l_i^\infty, \theta_i^\infty)$ for $i > k$ and, for $i \leq k$, l_i^n converges to zero. Next, using the Buser construction ([Bu], Theorem 8.1.3), we choose an $N_\infty \in \mathcal{T}_{g',m}$ such that $\lambda_1(N_\infty) = \epsilon < c$. Then by [C-C], Theorem 0.1, for any sequence (N_n) in \mathcal{M}_g converging to N_∞ in $\partial\mathcal{M}_g$, one has $\lim_{n \rightarrow \infty} \lambda_1(N_n) = \epsilon$. In particular we consider the sequence (N_n) such that $(l_i, \theta_i)(N_n) = (l_i, \theta_i)(N_\infty)$ for $i > k$ and $(l_i, \theta_i)(N_n) = (l_i, \theta_i)(P_n)$ for $i \leq k$. Then $\lim_{n \rightarrow \infty} \lambda_1(N_n) = \epsilon$.

At this point we construct a path σ_n in \mathcal{M}_g joining P_n and N_n for each n . Let us consider the path given by the coordinate axes i.e. the path first goes along the l_i axes from $l_i(P_n)$ to $l_i(N_n)$ for each $i = k+1, k+2, \dots, 3g-3$ in the increasing order and then the same for θ_i 's. Finally for any $t \in [\epsilon, c]$ we apply the continuity property of λ_1 on \mathcal{M}_g to get a surface Q_n on σ_n such that $\lambda_1(Q_n) = t$. By construction each point on σ , in particular Q_n , has $(l_i, \theta_i)(Q_n) = (l_i, \theta_i)(P_n)$ for $i \leq k$ and all other $(l_i, \theta_i)(Q_n)$ are bounded by the corresponding coordinates of P_∞ and N_∞ . Hence Q_n converges to a point $Q_\infty \in \partial\mathcal{M}_g$ and since $\lambda_1(Q_n) = t$ for each n we have $\lim_{n \rightarrow \infty} \lambda_1(Q_n) = t$.

Therefore V_1 contains $[\epsilon, c]$ and ϵ being arbitrary, the whole of $(0, c]$. That $\Lambda_i(g) \leq \frac{1}{4}$ follows from the last claim. \square

Remark 3.4. A relevant question to ask here is whether $\frac{1}{4}$ belongs to V_k or not. It is not hard to show, using [C-C], Theorem 0.1, that the existence of a surface $N \in \partial\mathcal{M}_g$ which has k components and $\lambda_1(N) \geq \frac{1}{4}$ would guarantee that $\frac{1}{4} \in V_k$. In this connection we would like to mention that if g , the genus of the surface, is large then a result of Brooks and Makover in [B-M] (see also [B-B-D]) shows that for any given ϵ there exists a surface $S \in \mathcal{M}_g$ with $\lambda_1(S) > \frac{3}{16} - \epsilon$ (and $> \frac{1}{4} - \epsilon$ if one assumes the Selberg conjecture).

4. Non-compact finite area hyperbolic surfaces

In this section we study non-compact finite area hyperbolic surfaces. Recall that $\mathcal{T}_{g,n}$ denotes the Teichmüller space of all marked hyperbolic surfaces with finite area and of geometric type (g, n) . Given any pair of pants decomposition of any $S' \in \mathcal{T}_{g,n}$ one can consider the Fenchel–Nielsen coordinates on $\mathcal{T}_{g,n}$. Fix one such coordinate system on $\mathcal{T}_{g,n}$. Denote by $\mathcal{T}_{g,n}^0$ the set of all surfaces in $\mathcal{T}_{g,n}$ all of whose twist parameters are equal to zero. Recall that each surface in $\mathcal{T}_{g,n}^0$ carries an involution ι which when restricted to each pair of pants is the orientation reversing involution that fixes the boundary components. This involution induces an involution on each eigenspace of the Laplacian. The eigenfunctions corresponding to the eigenvalue -1 are called *antisymmetric* and the corresponding eigenvalue is called an *antisymmetric eigenvalue*. We denote the i -th antisymmetric cuspidal eigenvalue of $S \in \mathcal{T}_{g,n}^0$ by $\lambda^{o,c}_i(S)$.

We observe that in Proposition 1.4 we have considered domains in S which are diffeomorphic either to discs or to annuli. Since S is compact, the domains have compact closures. Now for $S_0 \in \mathcal{T}_{g,n}$, we may have nodal domains whose closure is not compact. To tackle this problem we consider only those domains which are diffeomorphic either to discs or to annuli and where respective boundary curves are not homotopic to a puncture. For any such disc or annulus, Cheeger's inequality is still true (cf. [Cha]). The computations in Lemma 2.9 then apply. Therefore for any $\Omega \subseteq S_0$ diffeomorphic either to a disc or to an annulus whose boundary curves are not homotopic to a puncture, we have an explicit constant $\epsilon_0(S_0) > 0$ such that

$$\lambda_0(\Omega) \geq \frac{1}{4} + 2\epsilon_0(S_0) > \frac{1}{4} + \epsilon_0(S_0).$$

Proof of Theorem B. The proof proceeds along the same lines as that of Theorem A. We choose $\epsilon_0(S_0)$ as above and consider $\mathcal{E}_o^{\frac{1}{4} + \epsilon_0(S_0)}$, the subspace of $C^\infty(S_0)$, spanned by the anti-symmetric cuspidal eigenfunctions with eigenvalue

$\leq \frac{1}{4} + \epsilon_0(S_0)$. Then we use the same arguments as in Theorem A to prove that the dimension of $\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$ is less than g . First for $f \neq 0 \in \mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$ we consider the subgraph $\mathcal{G}(f)$ of $\mathcal{Z}(f)$ obtained by suppressing those components of $\mathcal{Z}(f)$ which are bounded and homotopic to a point in S_0 (equivalently, those which are contained in a bounded disc in S_0). Next we consider the components of $S_0 \setminus \mathcal{G}(f)$ with their signs attached as defined in Section 2.2. Denote by $\mathcal{F}(\iota)$ the fixed point set of the isometry ι . The set $\mathcal{F}(\iota)$ divides S_0 into two isometric components \mathcal{S}_1 and \mathcal{S}_2 . Each \mathcal{S}_i is a non-compact finite area hyperbolic surface with geodesic boundary and genus 0. Each puncture of S_0 gives rise to two ideal points, one on $\partial\mathcal{S}_1$ and another on $\partial\mathcal{S}_2$.

Claim 4.1. *For any $f \neq 0 \in \mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$ each component of $S_0 \setminus \mathcal{G}(f)$ is contained in one of the \mathcal{S}_i 's and is incompressible there.*

Proof. By antisymmetry of f with respect to ι we have $\mathcal{F}(\iota) \subseteq \mathcal{Z}(f)$. Since each bounded component of $\mathcal{F}(\iota)$ is incompressible, $\mathcal{F}(\iota) \subseteq \mathcal{G}(f)$. \square

Now we can argue as in the proof of Lemma 2.7 to conclude that the Euler characteristic of at least one component of $S_0 \setminus \mathcal{G}(f)$ is negative. In fact using the symmetry of $\mathcal{G}(f)$ with respect to ι , the Euler characteristic of at least one component of $\mathcal{S}_j \setminus \mathcal{G}(f)$ is negative for each $j = 1, 2$. Next we consider the unit sphere $\mathbb{S}(\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)})$ and the projective space $\mathbb{P}(\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)})$ over $\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$. Define $\chi^+(f)$, resp. $\chi^-(f)$, as the sum of the Euler characteristics of the components of $\mathcal{S}_1 \setminus \mathcal{G}(f)$ with positive sign, resp. negative. Consider the decomposition of $\mathbb{S}(\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)})$ into sets

$$\mathcal{C}_i = \{f \in \mathbb{S}(\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}): \chi^+(f) + \chi^-(f) = i\}.$$

The arguments in Lemma 2.9 can be applied. Using the incompressibility of the components of $\mathcal{S}_1 \setminus \mathcal{G}(f)$ the possible values of $\chi^+(f) + \chi^-(f)$ are at most $(g-1)$ (since $\chi(\mathcal{S}_i) = 1-g$) for any $f \in \mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$. Exactly the same arguments as in Theorem A work to prove that for any integer i , the covering map $\mathcal{C}_i \rightarrow P_i$ is trivial. We conclude that the dimension of $\mathcal{E}_o^{\frac{1}{4}+\epsilon_0(S_0)}$ is $\leq g$. \square

References

- [B] L. BERS, Spaces of degenerating Riemann surfaces. In L. Greenberg (ed.), *Discontinuous Groups and Riemann Surfaces*. Annals of Mathematics Studies 79. Princeton University Press, Princeton, N.J., and University of Tokyo Press, Tokyo, 1974, 43–55. [MR 0361051](#) [Zbl 0294.32016](#)

- [B-Z] Yu. D. BURAGO and V. A. ZALGALLER, *Geometric Inequalities*. Akademiya Nauk SSSR, Leningradskoe Otdelenie Matematicheskogo Instituta im. V. A. Steklova, "Nauka", Leningrad, 1980. English transl., Grundlehren der mathematischen Wissenschaften 285. Springer Series in Soviet Mathematics. Springer Verlag, Berlin, 1988. Translated from the Russian by A. B. Sosinskiĭ. [MR 0602952](#) [Zbl 0436.52009](#)
- [Bu] P. BUSER, *Geometry and Spectra of Compact Riemann Surfaces*. Progress in Mathematics 106. Birkhäuser Boston, Boston, MA, 1992. [MR 1183224](#) [Zbl 0770.53001](#)
- [B-B-D] P. BUSER, M. BURGER and J. DODZIUK, Riemann surfaces of large genus and large λ_1 . In T. SUNADA (ed.), *Geometry and Analysis on Manifolds*. Lecture Notes in Mathematics 1339. Springer Verlag, Berlin etc., 1988, 54–63. [MR 0961472](#) [Zbl 0646.53040](#)
- [B-C] P. BUSER and G. COURTOIS, Finite parts of the spectrum of a Riemann surface. *Math. Ann.* **287** (1990), 523–530. [MR 1060691](#) [Zbl 0711.58033](#)
- [B-M] R. BROOKS and E. MAKOVER, Riemann surfaces with large first eigenvalue. *J. Anal. Math.* **83** (2001), 243–258. [MR 1828493](#) [Zbl 0981.30031](#)
- [Cha] I. CHAVEL, *Eigenvalues in Riemannian Geometry*. Including a chapter by B. Randol. With an appendix by J. Dodziuk. Pure and Applied Mathematics 115. Academic Press, Orlando, FL, 1984. [MR 0768584](#) [Zbl 0551.53001](#)
- [C-C] B. COLBOIS and G. COURTOIS, Les valeurs propres inférieures à $1/4$ des surfaces de Riemann de petit rayon d'injectivité. *Comment. Math. Helv.* **64** (1989), 349–362. [MR 0998853](#) [Zbl 0684.53040](#)
- [D-P-R-S] J. DODZIUK, T. PIGNATARO, B. RANDOL and D. SULLIVAN, Estimating small eigenvalues of Riemann surfaces. In L. KEEN (ed.), *The Legacy of Sonya Kovalevskaya*. Contemporary Mathematics 64. American Mathematical Society, Providence, RI, 1987, 93–121. [MR 0881458](#) [Zbl 0607.58044](#)
- [H] A. HENROT, *Extremum Problems for Eigenvalues of Elliptic Operators*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006. [MR 2251558](#) [Zbl 1109.35081](#)
- [He] D. A. HEJHAL, Regular b -Groups, degenerating Riemann Surfaces, and spectral theory. *Mem. Amer. Math. Soc.* **88** (1990), no. 437. [MR 1052555](#) [Zbl 0718.11024](#)
- [I] H. IWANIEC, *Introduction to the Spectral Theory of Automorphic Forms*. Biblioteca de la Revista Matemática Iberoamericana. Revista Matemática Iberoamericana, Madrid, 1995. [MR 1325466](#) [Zbl 0847.11028](#)
- [K] L. KEEN, Collars on Riemann surfaces. In L. GREENBERG (ed.), *Discontinuous Groups and Riemann Surfaces*. Annals of Mathematics Studies 79. Princeton University Press, Princeton, N.J., and University of Tokyo Press, Tokyo, 1974, 263–268. [MR 0379833](#) [Zbl 0294.32016](#)
- [O] J.-P. OTAL, Three topological properties of small eigenfunctions on hyperbolic surfaces. In M. KAPRANOV, S. KOLYADA, Yu. I. MANIN, P. MOREE and L. POTYAGAILO (eds.), *Geometry and Dynamics of Groups and Spaces*. Progress in Mathematics 265. Birkhäuser Verlag, Basel, 2008, 685–695. [MR 2402419](#) [Zbl 1187.35145](#)

- [O-R] J.-P. OTAL and E. ROSAS, Pour toute surface hyperbolique de genre g , $\lambda_{2g-2} > 1/4$. *Duke Math. J.* **150** (2009), 101–115. [MR 2560109](#)
[Zbl 1179.30041](#)
- [R1] B. RANDOL, Small eigenvalues of the Laplace operator on compact Riemann surfaces. *Bull. Amer. Math. Soc.* **80** (1974), 996–1000. [MR 0400316](#)
[Zbl 0317.30017](#)
- [R2] —, Cylinders in Riemann surfaces. *Comment. Math. Helv.* **54** (1979), 1–5. [MR 0522028](#) [Zbl 0401.30036](#)
- [R3] —, A remark on λ_{2g-2} . *Proc. Amer. Math. Soc.* **108** (1990), 1081–1083. [MR 1000167](#) [Zbl 1000167](#)
- [Se] B. SÉVENNEC, Multiplicity of the second Schrödinger eigenvalue on closed surfaces. *Math. Ann.* **324** (2002), 195–211. [MR 1931764](#)
[Zbl 1053.58014](#)
- [S-W-Y] R. SCHOEN, S. WOLPERT, and S.-T. YAU, Geometric bounds on the low eigenvalues of a compact surface. In R. OSSERMAN and A. WEINSTEIN (eds.), *Geometry of the Laplace Operator*. Proceedings of Symposia in Pure Mathematics XXXVI. American Mathematical Society, Providence, R.I., 1980, 279–285. [MR 0573440](#) [Zbl 0446.58018](#)

(Reçu le 27 novembre 2012)

Sugata MONDAL, Institut de Mathématiques de Toulouse, Université Paul Sabatier,
118, route de Narbonne 31062 Toulouse Cedex 9, France
e-mail: sugmondal2000@gmail.com