

# **The zonal position of "Nothofagus" forests = Die bioklimatische Vegetationszonen der "Nothofagus"-Wälder = Las zonas de vegetación bioclimáticas de los bosques de "Nothofagus"**

Autor(en): **Haemet-Ahti, Leena**

Objekttyp: **Article**

Zeitschrift: **Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich**

Band (Jahr): **91 (1986)**

PDF erstellt am: **27.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-308845>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## The zonal position of *Nothofagus* forests

Die bioklimatischen Vegetationszonen der *Nothofagus*-Wälder

Las zonas de vegetación bioclimáticas de los bosques de  
*Nothofagus*

by

Leena HAEMET-AHTI

### CONTENTS

1. Introduction	218
2. Vegetation zones in the northern and southern hemisphere	218
3. Distribution of <u>Nothofagus</u> forests	219
4. Comparison of the <u>Nothofagus</u> forests to Eurasian forests	222
Summary - Zusammenfassung - Resumen	223
References	224

## 1. INTRODUCTION

The genus Nothofagus is biogeographically one of the most interesting plant genera, and there are several papers dealing with its distributional history (e.g. HUMPHRIES 1981 and papers cited by him, ASH 1982). Geobotanically, Nothofagus is not less exciting, and a North European botanist is especially interested in the bioclimatic zonal position of the Nothofagus forests because they represent the southernmost forests in the world, forming the main portion of the southernmost horizontal and vertical timberlines on the globe. My purpose is to try to interpret the zonal position of the Nothofagus forests in the terms used in the northern hemisphere by my colleagues and myself (AHTI et al. 1968, HAE-MET-AHTI 1965, 1976, 1979, 1981, HAE-MET-AHTI et al. 1974, TUHKANEN 1980, 1984) and to accomodate this terminology to the southern hemisphere. This comparison was already started by TUHKANEN (1984), who (1980, 1984) presented the climatic indices for the boreal zone and its sectors.

## ACKNOWLEDGEMENTS

I should like to extend my warmest thanks to Dr. Ulrich Eskuche, Corrientes, Argentina, Dr. David Galloway, London, U.K. (formerly Christchurch, New Zealand), and Mr. Rex Filson, Melbourne, Australia, who acted as the leaders of the excellent field trips that allowed me to become acquainted with various Nothofagus forests. Dr. Timo Koponen, Helsinki, Finland, told me about his observations in New Guinea, Dr. Sakari Tuhkanen and Mr. Ilpo Kuokka, Helsinki, gave me valuable information on the conditions in Tierra del Fuego.

## 2. VEGETATION ZONES IN THE NORTHERN AND SOUTHERN HEMISPHERE

The asymmetry of the world vegetation (TROLL 1948) has had a strong influence on the concepts of vegetation geography: often the southern part of the globe is - even to-day - supposed to have fewer vegetation zones than the northern part. For instance, SCHROEDER (1983) indicates that his austral zone has three counterparts in the northern hemisphere, viz. the meridional, nemoral, and boreal zones. The concept of the austral

zone, however, seems to be too wide for many geobotanists and they have accepted more detailed concepts in their global (e.g. WALTER and BOX 1976) or local (e.g. MEURK 1984 for New Zealand) classifications. The recognition of ecogeographically corresponding units in the southern and northern amphitropical vegetation is desirable, but not easy, because of the different flora and in part the different physiognomy as emphasized, e.g. by BLISS (1979) in his comparision of arctic and antarctic vegetation. However there are some promising efforts which show that although the vegetation zones in the northern and southern hemisphere outside the tropics are not identical they are nevertheless comparable. For instance, CZAJKA's (1968) detailed descriptive profile from Alaska to Tierra del Fuego is instructive. SCHWAAR (1981) pointed out the close phytosociological relationship of some peatland communities of Tierra del Fuego with certain holarctic units, e.g. with Oxycocco-Sphagnetea. TUHKANEN (1984) even published a map on the southern counterparts of the boreal zone primarily based on the biotemperature values.

### 3. DISTRIBUTION OF NOTHOFAGUS FORESTS

In South America the forests dominated by the deciduous trees Nothofagus alpina. N. obliqua and sometimes N. antarctica as well as by the evergreens. N. dombeyi and N. nitida (e.g. SCHMITHUESEN 1956, OBERDORFER 1960, HUECK 1966, ESKUCHE 1968, 1969, WEINBERGER 1973, McQUEEN 1977, QUINTANILLA 1977, VEBLEN et al. 1980, HUECK and SEIBERT 1981) are obviously chiefly antitemperate but show some geographic variation due to the existence of different bioclimatic sectors or partly different sub-zones within the antitemperate main zone. The southern horizontal limit of the antitemperate vegetation zone lies close to 44°S (TUHKANEN 1984, p. 42) where also the southern limit of QUINTANILLA's (1977, p. 22) temperate vegetation zone also lies.

The southern counterpart of the boreal vegetation zone, the antiboreal zone, includes the deciduous forests of N. pumilio and N. antarctica, and the evergreen ones of N. betuloides (cf. the descriptions by, e.g. KALELA 1941, ROIVAINEN 1954, SCHMITHUESEN 1956, OBERDORFER 1960, HOLDGATE 1961, HUECK 1966, PISANO 1970, 1971, 1973, 1977, 1981, 1983, MOORE

1970, 1983, YOUNG 1972, ESKUCHE 1973, McQUEEN 1976, DOLLENZ 1980, 1981, 1982a, 1982b, HUECK and SEIBERT 1981). There are considerable differences in the climatic ecology between the three antiboreal Nothofagus species, as beautifully illustrated by WEINBERGER (1973) and QUINTANILLA (1977). N. antarctica and N. pumilio can be compared with the European Betula pubescens subsp. tortuosa, which forms the vertical timberline in northern Fennoscandia and is similar in physiognomy. However, N. antarctica has a wider geobotanical amplitude: its lowermost occurrences are in the antitemperate zone, often as a tree of second-growth forests. It can also form the arid timberline against the Patagonian steppes. Its uppermost limit in the mountains seems to be located in the middle orointiboreal zone and the southernmost ones are not northern antiboreal (cf. TUHKANEN 1984). N. pumilio is mainly antiboreal; the lowermost and northernmost occurrences seem to be in the antihemiboreal subzone, and the uppermost ones in the middle antiboreal subzone.

The evergreen N. betuloides also seems to be fairly strictly antiboreal ranging up to the middle orointiboreal subzone. The antiboreal forest vegetation in South America apparently contains the counterparts for the hemiboreal, southern boreal and middle boreal subzones only, and the middle boreal one is exclusively present as a vertical subzone. The counterparts of the northern boreal subzone do exist but are obviously completely treeless and only represented as a vertical subzone at higher elevations. The same applies to the hemiarctic subzone.

In New Zealand there are four Nothofagus species - all evergreen - forming forests and often also timberlines in mountains (e.g., P. WARDLE 1964, 1967, J. WARDLE 1970, GODLEY 1975). In fact, the ranges of N. truncata and N. fusca are apparently antimeridional to (oro) antitemperate, while N. menziesii and N. solandri are mainly antitemperate to orointiboreal. According to MEURK's (1984) terminology the range of the first two species is warm to cool temperate and that of the last two species temperate to subantarctic. The antiboreal zone occurs on the South Island of New Zealand only vertically on the slopes of the mountains. The upper orointiboreal subzone seems to be completely treeless, because the horizontal timberline formed by Nothofagus is located in the middle of the lower orointiboreal subzone or sometimes even down in it. If we compare the climatic values of the timberline areas in New Zealand (e.g. MORRIS 1965) with parameters published by TUHKANEN (1984) for the boreal zone they indicate highly oceanic middle boreal climate with a

cool but long growing period. This is also distinctly seen in P. WARDLE's (1965, 1971, 1972, 1977) comparisons of the New Zealand timberlines with some North American and Eurasian ones. His experiments to grow some North American timberline trees above the tree line in New Zealand are very inspiring (cf. P. WARDLE 1971, 1972, 1973), but again distinctly show that the timberline is not always an equivalent line in the different mountains (cf. HAEMET-AHTI 1979). The timberlines in New Zealand also differ physiognomically from most of the timberlines in southern South America because all the timberline trees, including the Nothofagus species, in New Zealand are evergreen rather than deciduous.

In Tasmania there is the only deciduous Nothofagus species of the eastern hemisphere, viz. N. gunnii. It forms stands on exposed sites in high mountains, sometimes close to the timberline (e.g. JACKSON 1965, HOWARD 1981, COSTIN 1981, MINCHIN 1981). N. gunnii seems to occur in the oroantitemperate and perhaps also oroantihemiboreal zone. In SE Australia there are two Nothofagus species: N. gunninghamii, which also occurs in Tasmania, and N. moorei (e.g. HOWARD and HOPE 1970, HOWARD 1973, 1981, HOWARD and ASHTON 1973). They are evergreen and apparently antimeridional to antitemperate.

In New Guinea several species of the genus Nothofagus which all are evergreen occur (ASH 1982). They occur in the lower, middle, and upper montane zone, being frequent only in the middle montane zone (JOHNS 1982). The zones apparently correspond to the horizontal subtropical, meridional, and southern temperate (= northern antitemperate) zones. In New Caledonia there are also many Nothofagus species and geobotanically they seem to be much like the New Guinean ones (cf., DAWSON 1966, VAN STEENIS 1971). In New Guinea as well as in New Caledonia Nothofagus is not a timberline tree and is most abundant in the oroantimeridional zone, far below the timberline (e.g. ASH 1982).

If we compare the northernmost and southernmost timberline forests of the globe, the roles of the needle-leaved and broad-leaved, as well as the evergreen and deciduous, tree species seem to be different on the different continents contrary to some customary generalizations.

#### 4. COMPARISON OF THE NOTHOFAGUS FORESTS TO EURASIAN FORESTS

At the horizontal and vertical timberlines in northern Eurasia there are deciduous broad-leaved trees (Betula pubescens subsp. tortuosa, B. erma-nii, Alnus maximowiczii) in the most oceanic areas (HAEMET-AHTI and AHTI 1969) and the deciduous needle-leaved conifers (Larix sibirica, L. gme-linii) in the most continental areas. Amongst these the evergreen conifers (mainly Picea abies subsp. ovata) predominate. In North America deciduous broad-leaved trees (Alnus sinuata) occur at the timberline only in southeastern Alaska and adjacent Canada. The whole truly polar timberline is formed by evergreen conifers (Picea glauca, P. mariana and in the easternmost part Abies balsamea), and among them a deciduous conifer (Larix laricina) is common, especially in the continental areas (e.g. HUSTICH 1966). In the southernmost part of the southern hemisphere there are no conifers at the timberline. The evergreen broad-leaved trees (N. betuloides) occur in the extremely oceanic part of southernmost South America as well as in New Zealand (N. menziesii, N. solan-dri). The deciduous broad-leaved trees (N. pumilio, N. antarctica) occur at the timberline in southernmost South America only.

Also the forests at the margins of the steppe areas under boreal and antiboreal circumstances are quite different in the different continents. In Eurasia boreal steppes or steppe patches are surrounded by deciduous conifers (e.g. Larix gmelinii in northern boreal Yakutia, L. sibirica in hemiboreal northern Mongolia), and in North America either by evergreen conifers (Picea glauca in the northern and middle boreal Yukon) or by deciduous broad-leaved trees (Populus tremuloides and P. balsamifera in the hemiboreal part of Alberta). In South America deciduous broad-leaved trees (Nothofagus antarctica) occur on the margins of the southern Patagonian antihemiboreal and northern antiboreal steppes. There is hardly any general rule to explain the occurrence of these taxonomically, ecomorphologically, and physiognomically different tree species at the northernmost or southernmost timberlines. These timberlines are geobotanically simply not identical, and they are caused by different minimum factors (e.g. excessive cold, aridity, humidity, wind, or their combinations). It seems unlikely that we should be able to find a general law based only on the present climatic or other ecological circumstances. For instance, the concept of the diurnal climate (TROLL

1948 etc.) as a reason for the lack of the deciduous Nothofagus species in New Zealand was shown to be untenable by P. WARDLE (1963), who found growth rings and other indications produced by a distinctly seasonal climate there.

It is also necessary to consider the history of the flora when we make large-scale geobotanical comparisons over the globe, as stated by many authors in different connections (e.g. by DU RIETZ 1960, HUEBL 1969, HAEMET-AHTI and AHTI 1969, WALTER 1977, HAEMET-AHTI 1978, KLOETZLI 1983). The individual migrational histories of each species are expected to provide a logical explanation for the diversity in the present-day ranges and ecology of the Nothofagus species.

#### SUMMARY

The bioclimatic vegetation zones of the Nothofagus forests in southern South America and New Zealand are antitemperate to middle (oro) antitropical, in Australia antimeridional to antitemperate, in New Guinea and New Caledonia subtropical to antitemperate. Nothofagus species are timberline trees only in South America and New Zealand, and geobotanically the highest Nothofagus timberlines in both areas are obviously in the middle oroantiboreal subzone. In South America there are deciduous and evergreen species at the timberline, while in New Zealand only evergreen species are present. There is hardly any general rule based on the present ecoclimatic conditions to explain the occurrence of the evergreen versus deciduous broad-leaved or evergreen versus deciduous coniferous tree species at different kinds of timberlines in the northernmost and southernmost parts of the globe. The main reason for the present distribution of the different ecophysiognomic elements is apparently the history of the flora.

#### ZUSAMMENFASSUNG

Die bioklimatischen Vegetationszonen der Nothofagus-Wälder in Südamerika und Neuseeland sind antitemperiert zu mittel(oro)antiboreal, in Australien antimeridional zu antitemperiert, in Neu-Guinea und Neukaledonien subtropisch zu antitemperiert. Nothofagus-Arten bilden Waldgrenzen nur in Südamerika und Neuseeland. Die höchste Waldgrenze in den beiden Gebieten ist wahrscheinlich in der mittel-oroantiborealen Zone. In Südamerika gibt es sommergrüne und immergrüne Arten an den Waldgrenzen während in Neuseeland nur immergrüne Arten vorkommen. Es gibt kaum eine allgemeine Regel, die aufgrund der heutigen ökoklimatischen Verhältnisse das Vorkommen der immergrünen und sommergrünen Laubbäume oder der immergrünen und sommergrünen Nadelbäume an den verschiedenen Waldgrenzen in den nördlichsten und südlichsten Teilen der Erdkugel erklären könnte. Der Hauptgrund der heutigen Vorkommen der verschiedenen ökophysiognomischen Elemente ist wahrscheinlich die Geschichte der Flora.

## RESUMEN

Las zonas de vegetación bioclimáticas de los bosques de Nothofagus en Sudamérica y Nueva Zelandia son antitemperadas a medio-(oro)antiboreales; en Australia son antimeridionales a antitemperadas. Solamente en Sudamérica y en Nueva Zelandia, especies de Nothofagus establecen límites de bosque. En las dos regiones, el límite de bosque más elevado se encuentra, probablemente en la zona medio-oroantiboreal. En Sudamérica hay especies caducifolias y siempreverdes en los límites de bosque, solamente siempreverdes en Nueva Zelandia. Acaso existe una regla general para explicar la presencia de los arboles caducifolios o siempreverdes, sean latifoliados o acicularifolios, en los diferentes límites de bosques en los extremos boreales y australes del globo terrestre bajo las condiciones del clima actual. La distribución actual de los distintos elementos ecofisionómicos se debe, probablemente, a la historia de la flora.

## REFERENCES

- AHTI T., HAEMET-AHTI L. and JALAS J., 1968: Vegetation zones and their sections in northwestern Europe. Ann.Bot.Fenn. 5, 169-211.
- ASH J., 1982: The Nothofagus Blume (Fagaceae) of New Guinea. In: GRES-SITT J.L. (ed.), Biography and ecology of New Guinea. Monogr.Biol. 42, 355-380.
- BLISS L.C., 1979: Vascular plant vegetation of the southern circumpolar region in relation to antarctic, alpine and arctic vegetation. Canad. J.Bot. 57, 2167-2178.
- COSTIN A.B., 1981: Vegetation of high mountains in Australia. In: KEAST A. (ed.), Ecological biogeography of Australia. Monogr.Biol. 41, 719-731.
- CZAJKA W., 1968: Los perfiles vegetales de las Cordilleras entre Alaska y Tierra del Fuego. Colloq.Geogr. 9, 117-121.
- DAWSON J.W., 1966: Observations on Nothofagus in New Caledonia. Tuatara 14, 1-7.
- DOLLENZ A.O., 1980: Estudios fitosociológicos en el archipiélago del Cabo de Hornos. An.Inst.Patagonia 11, 225-238.
- DOLLENZ A.O., 1981: Estudios fitosociológicos en el archipiélago del Cabo de Hornos. An.Inst.Patagonia 12, 173-182.
- DOLLENZ A.O., 1982a: Estudios fitosociológicos en el archipiélago del Cabo de Hornos. An.Inst.Patagonia 13, 145-151.
- DOLLENZ A.O., 1982b: Estudios fitosociológicos en las reservas forestales Alacalufes e Isla Riesco. An.Inst.Patagonia 13, 161-170.
- DU RIETZ G.E., 1960: Remarks on the botany of the southern cold temperate zone. Proc.Royal Soc. London (B) 152, 500-507.
- ESKUCHE U., 1968: Fisionomía y sociología de los bosques de Nothofagus dombeyi en la región de Nahuel Huapi. Vegetatio 16, 192-201.
- ESKUCHE U., 1969: Berberitzengebüsche und Nothofagus antarctica-Wälder in Nordwestpatagonien. Vegetatio 19, 264-285.
- ESKUCHE U., 1973: Estudios fitosociológicos en la norte de Patagonia. Phytocoenologia 1, 64-113.
- GODLEY E.J., 1975: Flora and vegetation. In: KUSCHEL G. (ed.), Biogeography and ecology in New Zealand. Monogr.Biol. 27, 177-229.
- HAEMET-AHTI L., 1965: Notes on the vegetation zones of Western Canada, with special reference to the forests of Wells Gray Provincial Park, British Columbia. Ann.Bot.Fenn. 2, 274-300.

- HAEOMET-AHTI L., 1976: Bioticheskie podrazdeleniya borealnoy zony. Geobot.Kartogr. 1976, 51-58.
- HAEOMET-AHTI L., 1978: Timberline meadows in Wells Gray Park, British Columbia, and their comparative geobotanical interpretation. Syesis 11, 187-211.
- HAEOMET-AHTI L., 1979: The dangers of using the timberline as the "zero line" in comparative studies on altitudinal vegetation zones. Phyto-coenologia 6, 49-54.
- HAEOMET-AHTI L., 1981: The boreal zone and its biotic subdivision. Fennia 159, 69-75.
- HAEOMET-AHTI L. and AHTI T., 1969: The homologies of the Fennoscandian mountain and coastal birch forests in Eurasie and North America. Vegetatio 19, 208-219.
- HAEOMET-AHTI L., AHTI T. and KOPONEN T., 1974: A scheme of vegetation zones for Japan and adjacent regions. Ann.Bot.Fenn. 11, 59-88.
- HOLDGATE M.W., 1961: Vegetation and soils in the South Chilean Islands. J.Ecol. 49, 559-580.
- HOWARD T.M., 1973: Studies in the ecology of Nothofagus cunninghamii Oerst. Austral.J.Bot. 21, 67-102.
- HOWARD T.M., 1981: Southern closed-forest. In: GROVES R.H. (ed.), Australian vegetation. Cambridge Univ. Press. 104-120.
- HOWARD T.M. and ASHTON D.H., 1973: The distribution of Nothofagus cunninghamii rainforest. Proc.Royal Soc.Victoria 86, 47-76.
- HOWARD T.M. and HOPE G.S., 1970: The present and past occurrence of beech (Nothofagus cunninghamii Oerst.) at Wilsons Promontory, Victoria, Australia. Proc.Royal Soc.Victoria 83, 199-210.
- HUEBL E., 1969: Gedanken zur Verbreitung von sommergrüner und immergrüner Vegetation. Acta Bot.Croat. 28, 139-148.
- HUECK K., 1966: Die Wälder Südamerikas. Vegetationsmonographien der einzelnen Grossräume. Fischer, Stuttgart. 2, 1-422.
- HUECK K. and SEIBERT P., 1981: Vegetationskarte von Südamerika. (2nd ed.). Vegetationsmonographien der einzelnen Grossräume. Fischer, Stuttgart. 2a, 1-90.
- HUMPHRIES C.J., 1981: Biogeographical methods and the southern beeches (Fagaceae: Nothofagus). In: FUNK V.A. and BROOKS D.R. (eds.), Advances in cladistics. Proc.1st Meet.Hennig Soc., New York Bot.Garden, Bronx, New York, 177-208.
- HUSTICH I., 1966: On the forest-tundra and the northern tree-lines. Ann. Univ.Turkuensis A II 36, 7-47.
- JACKSON, W.D., 1965: Vegetation. In: DAVIES J.L. (ed.), Atlas of Tasmania. Lands and Surveys Department, Hobart. (not seen, cited according to WILLIAMS W.D. (ed.), 1974, Biogeography and ecology in Tasmania. Junk, The Hague. 498 pp.
- JOHNS R.J., 1982: Plant zonation. In: GRESSITT J.L. (ed.), Biogeography and ecology of New Guinea. Monogr.Biol. 42, 309-330.
- KALELA E.K., 1941: Ueber die Holzarten und die durch die klimatischen Verhältnisse verursachten Holzartenwechsel in den Wäldern Ostpatagoniens. Ann.Acad.Sci.Fenn.AIV,Biol. 2, 1-57.
- KLÖTZLI F., 1983: Standörtliche Grenzen von Fagaceen, ein Vergleich in beiden Hemisphären. Tuexenia 3, 47-65.
- MCQUEEN D.R., 1976: The ecology of Nothofagus and associated vegetation in South America. Tuatara 22, 38-68.
- MCQUEEN D.R., 1977: The ecology of Nothofagus and associated vegetation in South America. Tuatara 22, 233-265.
- MEURK C.D., 1984: Bioclimatic zones for the Antipodes - and beyond? New Zealand J.Ecol. 7, 175-181.

- MINCHIN P.R., 1981: The montane vegetation of Tasmania. In: JACKSON W.D. (ed.), The vegetation of Tasmania. XIII Intern.Bot.Congr.Field Trip Tasmania, 28, 85-99.
- MORRIS J.Y., 1965: Climate investigations in the Craigieburn Range, New Zealand. New Zealand J.Sci. 8, 556-582.
- MOORE D.M., 1970: Southern oceanic wet-heathlands (including Magellanic moorland). In: SPECHT R.L. (ed.), Heathlands and related shrublands of the world. Elsevier, Amsterdam. 54-59.
- MOORE D.M., 1983: Flora of Tierra del Fuego. Missouri Bot.Garden, St. Louis. 396 pp.
- OBERDORFER E., 1960: Pflanzensoziologische Studien in Chile. Flora et Veget.Mundi 2, 1-208.
- PISANO V.E., 1970: Vegetación del área de los Fjordos Toro y Condor y Puerto Cutter Cove (Canal Jerónimo, Magallanes). An.Inst. Patagonia 1, 27-40.
- PISANO V.E., 1971: Communidades vegetales del área del Fjordo Parry, Tierra del Fuego. An.Inst.Patagonia 2, 93-133.
- PISANO V.E., 1973: La vida en los Parques Nacionales de Magallanes. Inst.Patagonia Ser.Monogr. 6, 1-64.
- PISANO V.E., 1977: Fitogeografía de Fuego-Patagonia Chilena. I. Comunidades vegetales entre las latitudes 52° y 56° sur. An.Inst.Patagonia 8, 121-250.
- PISANO V.E., 1981: Bosquejo fitogeográfico de Fuego-Patagonia. An.Inst. Patagonia 12, 159-171.
- PISANO V.E., 1983: The Magellanic tundra complex. In: GORE A.J.P. (ed.) Mires: Bog, fen and moor. B. Regional studies. Elsevier, Amsterdam, 295-329.
- QUINTANILLA V., 1977: A contribution to the phytogeographical study of temperate Chile. Biogeographica 8, 31-41.
- ROIVAINEN H., 1954: Studien über die Moore Feuerlands. Ann.Bot.Soc. Vanamo 28 (2), 1-205.
- SCHMITHUESEN J., 1956: Die räumliche Ordnung der chilenischen Vegetation. Bonner Geogr.Abh. 17, 1-86.
- SCHROEDER F.-G., 1983: Die thermischen Vegetationszonen der Erde. Tuexenia 3, 31-46.
- SCHWAAR J., 1981: Amphi-arktische Pflanzengesellschaften in Feuerland. Phytocoenologia 9, 547-572.
- STEENIS van, C.G.G., 1971: Revision of Nothofagus in New Caledonia. Adansonia 11, 615-624.
- TROLL C., 1948: Der asymmetrische Aufbau der Vegetationszonen und Vegetationsstufen auf der Nord- und Südhalbkugel. Ber.Geobot.Forsch.Inst. Rübel 1947, 46-83.
- TUHKANEN S., 1980: Climatic parameters and indices in plant geography. Acta Phytogeogr.Suecica 67, 1-105.
- TUHKANEN S., 1984: A circumboreal system of climatic-phytogeographical regions. Acta Bot.Fenn. 127, 1-50.
- VEBLEN T.T., SCHLEGEL F.M. and ESCOBAR B.S., 1980: Structure and dynamics of old-growth Nothofagus forest in the Valdivian Andes, Chile. J.Ecol. 68, 1-31.
- WALTER H., 1977: Vegetationszonen und Klima. Ulmer, Stuttgart. 309 pp.
- WALTER H. and BOX E., 1976: Global classification of natural terrestrial ecosystems. Vegetatio 32, 75-81.
- WARDLE J.A., 1970: The ecology of Nothofagus solandri. New Zealand J. Bot. 8, 494-646.
- WARDLE P., 1963: Growth habits of New Zealand subalpine shrubs and trees. New Zealand J.Bot. 1, 18-47.

- WARDLE P., 1964: Facets of the distribution of forest vegetation in New Zealand. *New Zealand J.Bot.* **2**, 352-366.
- WARDLE P., 1965: A comparison of alpine timberlines in New Zealand and North America. *New Zealand J.Bot.* **3**, 113-135.
- WARDLE P., 1967: Biological flora of New Zealand. (2). *Nothofagus menziesii*. *New Zealand J.Bot.* **5**, 549-554.
- WARDLE P., 1971: An explanation for alpine timberline. *New Zealand J. Bot.* **9**, 371-402.
- WARDLE P., 1972: New Zealand timberlines. *Tussock Grasslands and Mountain Lands Institute Review* **23**, 31-48.
- WARDLE P., 1973: New Zealand timberlines. *Arctic Alpine Research* **5(3)**, A127-A135.
- WARDLE P., 1977: Japanese timberline and some geographic comparisons. *Arctic Alpine Research* **9(3)**, 249-258.
- WEINBERGER P., 1973: Beziehungen zwischen mikroklimatischen Faktoren und natürlicher Verjüngung araukano-patagonischer Nothofagusarten. *Flora* **162**, 157-179.
- YOUNG S.B. 1972: Subantarctic rain forest of Magellanic Chile: distribution, composition, and age and growth rate studies of common forest trees. *Antarctic Res.Ser. U.S.* **20**, 307-322.

Address of the author: Prof. Dr. Leena Hämet-Ahti  
Botanical Garden  
University of Helsinki  
Unionink. 44,  
SF-00170 Helsinki, Finland