

List of figures

Objekttyp: **Index**

Zeitschrift: **Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich**

Band (Jahr): **94 (1989)**

PDF erstellt am: **01.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

LIST OF FIGURES

	page
1. Location of the study area.	8
2. Aerial view of the study area.	9
3. Climatic diagram of Sackville, New Brunswick.	9
4. Temperature, rainfall and water level at the site in 1982.	10
5. Schematic profile of the floating mat in the undrained basin.	11
6. Experimental design.	13
7. Definitions of assimilation period and duration of senescence.	16
8. General phenological development of <u>Typha glauca</u> during 1982 on unburned and unfertilized plots in the undrained basin.	21
9. Development of the shoot density of <u>Typha glauca</u> on unburned and burned plots in the undrained basin, on which were superimposed five fertilizer treatments.	26
10. Development of the shoot density of <u>Typha glauca</u> on unburned and burned plots in the drained basin, on which were superimposed five fertilizer treatments.	27
11. Shoot density on June 19 and final shoot density of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	28
12. Shoot density of <u>Typha glauca</u> on June 19, in percent of final density, under different draining, burning and fertilizer treatments.	29
13. Shoot density of <u>Typha glauca</u> on June 19, in percent of final density, in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	30
14. Shoot height on June 19 and final shoot height of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	32
15. Development of mean total shoot height and mean height of the green shoot portion of <u>Typha glauca</u> on unburned and burned plots in the undrained basin, on which were superimposed five fertilizer treatments.	34
16. Development of mean total shoot height and mean height of the green shoot portion of <u>Typha glauca</u> on unburned and burned plots in the drained basin, on which were superimposed five fertilizer treatments.	35
17. Basal shoot circumference of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	37
18. Basal shoot circumference of <u>Typha glauca</u> in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	38
19. Number of leaves per shoot of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	39
20. Number of leaves per shoot of <u>Typha glauca</u> in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	40
21. Length of the assimilation periods 50% and 0% of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	42
22. Length of the assimilation periods 50% and 0% of <u>Typha glauca</u> in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	43

23. Start of the assimilation periods 50% and 0% of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	44
24. Start of the assimilation periods 50% and 0% of <u>Typha glauca</u> in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	46
25. Day after May 15, on which mean shoot height of <u>Typha glauca</u> reached 71.1 cm, that is 50% of the final height in the undrained, unburned and unfertilized treatment plots, under different draining, burning and fertilizer treatments.	48
26. Day after May 15, on which mean shoot height of <u>Typha glauca</u> reached 71.1 cm, that is 50% of the final height in the undrained, unburned and unfertilized treatment plots, in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	49
27. Duration of senescence 50% and 100% of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	51
28. Duration of senescence 50% and 100% of <u>Typha glauca</u> , in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	52
29. Height of the green shoot portion of <u>Typha glauca</u> on October 3, in percent of total shoot height, under different draining, burning and fertilizer treatments.	53
30. Height of the green shoot portion of <u>Typha glauca</u> on October 3, in percent of total shoot height, in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	54
31. Susceptibility of <u>Typha glauca</u> to drought: Height of green parts, during June and July 1982, in the drained treatment on which were superimposed burning and fertilizer treatments.	57
32. Susceptibility of <u>Typha glauca</u> to drought: Leaf die-back on June 19, under different draining, burning and fertilizer treatments.	58
33. Susceptibility of <u>Typha glauca</u> to drought: Leaf die-back on June 19 in the drained basin, in response to different fertilizer treatments which were superimposed on two burning regimes.	59
34. Insect damage: Percentage of <u>Typha glauca</u> shoots attacked by stem-boring insect larvae, under different draining, burning and fertilizer treatments.	62
35. Insect damage: Percentage of <u>Typha glauca</u> shoots attacked by stem-boring insect larvae, in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	63
36. Shoot standing crop of <u>Typha glauca</u> as predicted from mean shoot height and mean shoot density (contour plot).	64
37. Shoot standing crop of <u>Typha glauca</u> under different draining, burning and fertilizer treatments.	65
38. Shoot standing crop of <u>Typha glauca</u> in response to different fertilizer treatments which were superimposed on four draining x burning regimes.	66
39. Aboveground standing crop of plants other than <u>Typha</u> under two different fertilizer treatments which were superimposed on four draining x burning regimes.	69
40. Litter load under two different fertilizer treatments which were superimposed on four draining x burning regimes.	70