

Liste of figures : limnology

Objekttyp: **Index**

Zeitschrift: **Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich**

Band (Jahr): **105 (1991)**

PDF erstellt am: **04.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

List of figures / Limnology	page
2.1. Map of the sampling area with the sampling stations. Distance between contour lines: 5 m. Mean water level from September 25 to October 27, altitude above sea level: 435.5 m.	80
3.1. Water level at the time of investigations and mean monthly water level during the investigation period from April 1987 to April 1988. Data from the Swiss Hydrological Service (Schweizerische Landeshydrologie und -geologie).	87
3.2. Average vertical attenuation coefficient of photosynthetically available radiation and transparency (Secchi-depth) in the littoral and in the pelagial during the investigation period from April 1987 to April 1988. m.v. = missing value.	88
3.3. Mean temperature in the littoral for the 0-5 m depth interval and in the pelagial for the 0-10 m depth interval during the investigation period from April 1987 to April 1988.	89
3.4. Conductivity in the littoral (mixed water, water column from the 0-4 m depth interval) and in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	91
3.5. pH values in the littoral for the 0-5 m depth interval and in the pelagial for the 0-10 m depth interval during the investigation period from April 1987 to April 1988.	92
3.6. Chloride concentration in the littoral (mixed water, water column from the 0-4 m depth interval) and in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	93
3.7. Oxygen concentration in the littoral for the 0-5 m depth interval and in the pelagial for the 0-10 m depth interval during the investigation period from April 1987 to April 1988.	94
3.8. Oxygen saturation in the littoral for the 0-5 m depth interval and in the pelagial for the 0-10 m depth interval during the investigation period from April 1987 to April 1988.	94
3.9. Nitrate concentration (NO_3^- -N) in the littoral (mixed water, water column from the 0-4 m depth interval) and in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	95
3.10. Ortho-phosphate concentration (PO_4^{3-} -P in the littoral (mixed water, water column from the 0-4 m depth interval) and in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	96
3.11. Total phosphorus concentration in the littoral (mixed water, water column from the 0-4 m depth interval) and in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	97
4.1. Mean density of individuals for phytoplankton at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	114

4.2.	Wet weight of phytoplankton at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	114
4.3.	Relative frequency of the phytoplankton at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	116
4.4.	Relative amounts of several algal groups on the wet weight of the plankton at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	117
4.5.	Chlorophyll a concentration at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	119
4.6.	Dry weight at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	119
4.7.	Ash-free dry weight at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	121
4.8.	Ratio of ash-free dry weight to total dry weight at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	121
4.9.	Density of individuals for zooplankton at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling stations in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depths) during the investigation period from April 1987 to April 1988.	122
5.1.	Rack of glass slides with bar to fix on the ground. L: 48.5 cm, H: 8.5 cm (without bar).	125
5.2.	Exposure depth of the glass slides at 2.5 m and 5 m depths during the investigation period from May 1987 to April 1988.	126
5.3.	Chlorophyll a content on the glass slides at 2.5 m and 5 m depths during the investigation period from May 1987 to April 1988. m.v. = missing value.	130
5.4.	Ash-free dry weight on the glass slides at 2.5 m and 5 m depths during the investigation period from May 1987 to April 1988. m.v. = missing value.	132
5.5.	Percent of chlorophyll a of the ash-free dry weight on the glass slides at 2.5 m and 5 m depths during the investigation period from May 1987 to April 1988.	136
5.6.	Degree of cover of diatoms, cyanophytes and chlorophytes on the glass slides at 2.5 m and 5 m depths (diatoms only) during the investigation period from May 1987 to April 1988. m.v. = missing value, values for the cyanophytes and	

the chlorophytes at 5 m depth always $\leq 10\%$.	139
5.7. Relative frequency of the main diatom species on the glass slides at 2.5 m depth during the investigation period from May 1987 to April 1988. Hatched = major species (relative frequency at each sampling stations 10% at least once), numbers 1-6 = species with relative frequencies at the corresponding sampling station 10% at least once. m.v. = missing value.	148
5.8. Relative frequency of the main diatom species on the glass slides at 5 m depth during the investigation period from May 1987 to April 1988. Hatched = major species (relative frequency at each sampling stations 10% at least once), numbers 1-6 = species with relative frequencies at the corresponding sampling station 10% at least once. m.v. = missing value.	149
5.9. Relative frequency of the diatom differential groups on the glass slides at 2.5 m and 5 m depth during the investigation period from May 1987 to April 1988. rH 1 = 100% relative frequency, \bar{u} s = very sensitive species, s = sensitive species, wt = less tolerant species, t = tolerant species.	159
6.1. Labyrinth trap (20x14x2.5 cm) for macroinvertebrates. Left: as exposed at the lake bottom, right: opened.	162
6.2. Properties of lake bottom at the six sampling stations at 5 m depth. Average estimates of the investigations from January and from April 1988.	163
6.3. Dry weight (TG) and ash-free dry weight (AFTG) of the sedimented particles and ratio of ash-free dry weight to total dry weight at sampling stations 1, 2 and 3 in the littoral during the investigation period from April 1987 to April 1988.	165
6.4. Dry weight (TG) and ash-free dry weight (AFTG) of the sedimented particles and ratio of ash-free dry weight to total dry weight at sampling stations 4, 5 and 6 in the littoral during the investigation period from April 1987 to April 1988.	166
6.5. Mean number of individuals per labyrinth trap for macroinvertebrates at the six sampling stations during the investigation period from May 1987 to April 1988. N = 4-6. m.v. = missing value.	169
6.6. Relative frequency of the main macroinvertebrate groups at the six sampling stations during the investigation period from May 1987 to April 1988. m.v. = missing value.	172
6.7. Number of individuals per labyrinth trap for <i>Dugesia polychroa/lugubris</i> , <i>Policelis nigra/tenuis</i> and <i>Dendrocoelus lacteum</i> at the six sampling stations during the investigation period from May 1987 to April 1988. Note the different scales. N = 4-6. m.v. = missing value.	173
6.8. Number of individuals per labyrinth trap for <i>Herpobdella octoculata</i> , <i>H. stagnalis</i> and <i>Glossiphonia complanata</i> at the six sampling stations during the investigation period from May 1987 to April 1988. Note the different scales. N = 4-6. m.v. = missing value.	174
6.9. Number of individuals per labyrinth trap for <i>Asellus aquaticus</i> at the six sampling stations during the investigation period from May 1987 to April 1988. N = 4-6. m.v. = missing value.	175
6.10. Number of individuals per labyrinth trap for snails at the six sampling stations during the investigation period from May 1987 to April 1988. N = 4-6. m.v. = missing value.	176
7.1. Area of the theoretical (0-20 m depth interval) and the actual (0-5 m) shallow water area and the vegetation area in the Reuss delta region. Data from 1982/83 by LACHAVANNE et al. (1985).	

Sector 1: on the right of the River Reuss, Allmeini and Schützenrütli	
Sector 2: on the left of the River Reuss, Flüelerschachen and Schanz	
Sector 3: on the left of the River Reuss, Schwäb and Seedorferbucht.	188
7.2.-	
7.7. Comparison of the vegetation area (size and flora composition) in the littoral 1982/83 and 1988.	
Sector 1 = Allmeini and Schützenrütli	
Sector 2 = Flüelerschachen and Schanz	
Sector 3 = Schwäb and Seedorferbucht	
----- = 5 m isobath (boundary of the actual shallow water zone with regard to the mean water level of many years, altitude above sea level: 433.6 m.	
Investigation from 1982/83: x = occurrence of the species without declaration of the relative frequency .	
Investigation from 1988: x = occurrence of one or a small number of	191-197
8.1. Density of individuals for <i>Tabellaria fenestrata</i> , <i>Rhodomonas minura</i> and the group of the diverse flagellates in August (before the flood event) and in September (after the flood event) at the sampling stations in the littoral and in the pelagial. Pel li = pelagial on the left of River Reuss mouth, Pel re = pelagial on the right of River Reuss mouth.	205
9.1. Possible development of the phytoplankton community in the littoral of Lake Uri as a result of the new formation of the River Reuss mouth.	211
9.2. Possible development of the macroinvertebrate communities in the littoral of Lake Uri as a result of the new formation of the River Reuss mouth.	214

List of tables / Limnology

2.1. Latitude and longitude of the sampling stations in the pelagial and the littoral. Latitude and longitude refers to the contour map 1:5000, Reusshochwasser, Canton Uri.	79
2.2. Dates of the chemical, physical and biological investigations and exposition time of the glass slides.	81
3.1. Annual mean and standard deviation of the investigated parameters during the period April 1987 to April 1988. M = mean, S = standard deviation.	86
3.2. Results of the statistical comparisons between the six sampling stations in the littoral and between the two in the pelagial as well as between the littoral and the pelagial. Significance level $p \leq 0.05$, * = $p > 0.1$.	98
3.3. Results of the bacteriological investigation from August 5, 1987 (carried out by the Labor der Urkantone) and February 1, 1988 (our investigation). The investigation were carried out in the Giessen (a channel which acts as a receiving body of water for the water treatment plant of Altdorf), within the bay behind the island for the birds and to the left of the island. * = at the mouth of Giessen, ** = about 50 m upstream of the Giessen mouth, *** = on the right to the Giessen mouth, near the shore.	102
3.4. Comparison of the nutrient concentrations of some Swiss lakes. Values of Lake Uri from January to March 1988. * = Mean of the littoral stations	