

# Summary

Objekttyp: **Chapter**

Zeitschrift: **Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich**

Band (Jahr): **120 (1994)**

PDF erstellt am: **27.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

der Fall ist. Aus diesem Grund wird empfohlen, diese Grossseggenbestände zu einem *Magnocaricetum* zusammenzufassen. Falls trotzdem eine Unterteilung vorgenommen werden soll, so ist diese nach der dominierenden *Carex*-Art vorzunehmen.

## SUMMARY

In the present study particular interest is focused on the site conditions of *Magnocaricion* associations which are dominated by *Carex elata*, *Carex paradoxa*, *Carex paniculata* and *Carex riparia*. The associations are: *Caricetum elatae typicum*; *Caricetum elatae comaretosum*, typical variant; *Caricetum elatae comaretosum*, *Carex paradoxa*-variant; *Caricetum paniculatae*; *Caricetum ripariae*; *Caricetum vesicariae*.

Most of the study sites are situated in Kanton Zürich. Further sites are at the border of the lake of the Reuss (Kanton Aargau), at the southern border of the lake of Neuenburg, and in the Jura.

The phytosociological relevés of the study sites were grouped, first, according to the characteristic and differential species and second, by multivariate analysis.

During the vegetation periods from 1986 to 1988 the ground-water table was monthly measured and water samples were chemically analysed (pH, electrical conductivity, ortho-PO<sub>4</sub>, P-tot, NO<sub>3</sub>, NH<sub>4</sub>, Na, K, Ca, Mg). Soil samples were taken in autumn and chemical analyses were carried out for C<sub>org</sub>, pH, ortho-PO<sub>4</sub>, P-tot, NO<sub>3</sub>, NH<sub>4</sub>, Na, K, Ca, Mg. Furthermore, the above- and belowground tissue nutrient concentrations (N-tot, P-tot, C<sub>org</sub>, K, Na, Ca, Mg, Mn, Fe, Cu) of the *Carex* species and their biomass were investigated. The exponential mean temperatures were measured.

A principal component analysis was carried out using the data of the site conditions. Their relation to the vegetation groups was investigated by discriminant and variant analyses.

The study sites show differences in the electrical conductivity and the phosphorus concentrations of the water, the total nitrogen and the organic carbon concentrations and the V-value of the soil. (All studied soils belong to the "Anmoor"-soils.) These differences do not clearly correlate to the vegetation groups.

The vegetation groups formed by multivariate analyses are better differentiated than the vegetation groups formed by characteristic species. In both cases the S-value of the soil is one of the most discriminant factors. Moreover the Mg-concentrations of the soil and the electrical conductivity of the water also play a differential role.

However, the occurrence of these *Magnocaricion* associations at a certain site is not explained by these factors. The uptake of nitrogen and phosphorus by the plants seems to be more important. This is influenced by the periodically high water levels and floodings. The uptake of phosphorus depends on the solubility in the soil, which is related to the water regime. The nitrogen uptake depends on the ability of the plants to use ammonium instead of nitrate. If the plants use only nitrate, the water flow is important. The nitrogen and phosphorus uptake cannot be demonstrated by the water and soil data, but can be seen by the nitrogen and phosphorus concentrations of the plants. *Carex paniculata* has higher average nitrogen and phosphorus concentrations and *Carex riparia* higher average phosphorus concentrations than *Carex elata* and *Carex paradoxa*. For these four *Carex* species the uptake of nitrogen and phosphorus in relation to the water regime is discussed. The present study demonstrates that the nutrient concentrations of the *Carex* species as well as the site conditions of the studied associations do not differ significantly. Therefore it is proposed to categorise these six associations as one - *Magnocaricetum* - with the possible further sub-division of the dominating *Carex* species.