

The European mole

Autor(en): **[s.n.]**

Objekttyp: **Article**

Zeitschrift: **Helvetia : magazine of the Swiss Society of New Zealand**

Band (Jahr): **75 (2009)**

Heft [8]

PDF erstellt am: **30.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-944524>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

Sender:

The Secretary of the Swiss Society of New Zealand Inc
Trudi Brühlmann
401 Marine Drive
Mahina Bay
Eastbourne
Lower Hutt 5013

New Zealand
Permit No. 93966



HELVE⁺IA

Magazine for Members of the Swiss Society of New Zealand Inc

FAUNA IN SWITZERLAND

The European Mole

The European mole has a cylindrical body and is around 12 cm long. Females are typically smaller than males. The eyes are small and hidden behind fur. Its ears are just a small ridge in the skin. The fur is usually dark grey, but the actual range of colors is larger, as due to the subterranean habits there is no disadvantage in off-colored fur.

Moles are found throughout Middle Asia and Europe, but not in Ireland. They are present in most habitats where the soil is deep enough to allow tunnelling but are uncommon in coniferous forests, on moorlands and in sand dunes, probably because their prey is scarce.

Moles spend almost all their lives underground in a system of permanent and semi-permanent tunnels which form a complex network hundreds of metres long at varying depths in the soil. The deepest tunnels are used most in times of drought and low temperatures. Permanent tunnels are used repeatedly for feeding over long periods of time, sometimes by several generations of moles. The fore limbs are used to dig, shearing soil from the sides of the tunnel with alternate strokes. Hind limbs are used to brace the mole's body against the tunnel walls. The mole turns round, scoops up accumulated soil with its fore limbs and pushes it along a previously dug side tunnel leading to the surface. The soil is pushed out above ground to form a molehill.

Within the tunnel system moles construct one or more spherical nest chambers, each lined with a ball of dry plant material. Nests are used for sleeping and for raising young.

Moles sometimes construct very large mounds containing more than 750kg of soil. The mounds have an internal structure with one or more nests

and a network of tunnels and can have food stores. They are built by males and females, and they occur most often in areas with shallow soil on hard substrate and in areas prone to flooding. The fortress acts as a refuge in times of flooding and also helps insulate the nest against low temperatures.



European mole, *Talpa europaea*

Earthworms are the most important component of the mole's diet; an 80g mole needs 50g of earthworms per day. Moles also eat many insect larvae particularly in the summer. Earthworms dominate the winter diet. Moles sometimes collect and store them alive in special chambers. The stored worms are immobilised by a bite to the head segment; 470 worms have been recorded in one chamber. Food is either actively dug out of the soil by the mole or more often collected from the floor of the tunnel. Many soil animals fall through into the tunnels. Moles rarely forage on the surface.

internet