

# A remark on anisotropic superconducting states

Autor(en): **Feldmann, Joel / Knörrer, Horst / Trubowitz, Eugene**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **64 (1991)**

Heft 5

PDF erstellt am: **21.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116318>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A Remark on Anisotropic Superconducting States

Joel Feldman\*  
Department of Mathematics  
University of British Columbia  
Vancouver, B.C. V6T 1Y4  
CANADA

Horst Knörrer  
Eugene Trubowitz  
Mathematik  
ETH-Zentrum  
CH-8092 Zürich  
SWITZERLAND

(23. III. 1991)

Abstract

We show that, in three dimensions, there are no nontrivial, isotropic, unitary solutions of the gap equation for angular momentum greater than one, while in two dimensions they exist in all angular momentum sectors.

---

\* Research supported in part by the Natural Science and Engineering Research Council of Canada

Consider the many Fermion system in three dimensions characterized by the effective potential

$$\mathcal{G}(\psi^e, \bar{\psi}^e) = \log \frac{1}{Z} \int e^{-\lambda \mathcal{V}(\psi + \psi^e, \bar{\psi} + \bar{\psi}^e)} d\mu_C(\psi, \bar{\psi}),$$

$$\begin{aligned} \mathcal{V}(\psi, \bar{\psi}) = \frac{1}{2} \sum_{\mathbf{a}_i \in \{\uparrow, \downarrow\}} \int \prod_{i=1}^4 \frac{d^4 k_i}{(2\pi)^4} (2\pi)^4 \delta(k_1 + k_2 - k_3 - k_4) \delta_{\mathbf{a}_1, \mathbf{a}_3} \delta_{\mathbf{a}_2, \mathbf{a}_4} \\ \langle k_1, k_2 | V | k_3, k_4 \rangle \bar{\psi}(k_1, \mathbf{a}_1) \bar{\psi}(k_2, \mathbf{a}_2) \psi(k_4, \mathbf{a}_4) \psi(k_3, \mathbf{a}_3), \end{aligned}$$

where  $d\mu_C(\psi, \bar{\psi})$  is the fermionic Gaussian measure in the Grassmann variables

$$\{\psi(\xi), \bar{\psi}(\xi) | \xi = (\tau, \mathbf{x}, \sigma), \tau \in \mathbf{R}, \mathbf{x} \in \mathbf{R}^3, \sigma \in \{\uparrow, \downarrow\}\}$$

with covariance

$$\begin{aligned} C(\xi_1, \xi_2) &= \langle \psi(\xi_1) \bar{\psi}(\xi_2) \rangle \\ &= \delta_{\sigma_1, \sigma_2} \int \frac{d^{d+1} k}{(2\pi)^{d+1}} \frac{e^{i\langle k, \xi_1 - \xi_2 \rangle_-}}{ik_0 - e(\mathbf{k})} \\ \langle k, (\tau, \mathbf{x}) \rangle_- &= -k_0 \tau + \mathbf{k} \cdot \mathbf{x}, \quad \mathbf{k} = (k_0, \mathbf{k}) \\ e(\mathbf{k}) &= \frac{\mathbf{k}^2}{2m} - \mu. \end{aligned}$$

and where the two-body interaction  $\langle k_1, k_2 | V | k_3, k_4 \rangle$  is rotation invariant. That is

$$\langle Rk_1, Rk_2 | V | Rk_3, Rk_4 \rangle = \langle k_1, k_2 | V | k_3, k_4 \rangle$$

for any element  $R$  of  $SO(3)$  acting on spatial components. The chemical potential  $\mu$  in  $e(\mathbf{k})$  determines the electron density of the model.

The infrared behaviour of this model is determined (see [FT]) by a running coupling “constant”  $F^{(h)}(t', s')$ ,  $h \leq 0$  where at scale  $h$  the momentum  $k$  is restricted to a shell  $M^h$  away from the Fermi surface  $e(\mathbf{k}) = 0$  and  $t' = \left(0, \frac{t}{|t|} k_F\right)$  projects  $t$  onto the Fermi surface. Initially

$$F^{(0)}(t', s') = -\lambda \langle t', -t' | V | s', -s' \rangle.$$

The kernel  $F^{(h)}(t', s')$  defines an operator on  $L^2(k_F S^2)$ .

By rotation invariance the operator  $F^{(h)}$  commutes with the action of  $SO(3)$ . Therefore the eigenspaces of  $F^{(h)}$  coincide with the  $SO(3)$  irreducible invariant subspaces

of  $L^2(k_F S^2)$ . Recall that the space  $H^n$ , obtained by restricting homogeneous harmonic polynomials of degree  $n$  to  $S^2$ , is a  $2n + 1$  dimensional  $SO(3)$  irreducible invariant subspace of  $L^2(k_F S^2)$  and that

$$L^2(k_F S^2) = \oplus_{n \geq 0} H^n.$$

It follows that

$$F^{(h)}(t', s') = \sum_{n \geq 0} \lambda_n^{(h)} \pi_n(t', s')$$

where  $\pi_n$  is the orthogonal projection onto  $H^n$  and  $\lambda_n, n \geq 0$  is the spectrum of  $F^{(h)}$ . Here,  $\pi_n(t', s') = (2n + 1)k_F^{-2-n} P_n(\langle t', s' \rangle)$  where  $P_n$  is the Legendre polynomial of degree  $n$ .

It is widely believed that any (sufficiently weak) interaction  $\langle k_1, k_2 | V | k_3, k_4 \rangle$  flows, after, say,  $h$  steps, to an effective interaction  $F^{(h)}$  that is dominated by a single attractive angular momentum sector  $\lambda_\ell^{(h)} > 0$  (see [KL]). The infrared behaviour is then likely to be determined by the corresponding BCS model with gap equation

$$\Delta(\mathbf{p}) = \frac{1}{2} \int_{|e(\mathbf{q})| \leq \epsilon} \frac{d^3 \mathbf{q}}{(2\pi)^3} \lambda_\ell^{(h)} \pi_n(\mathbf{p}', \mathbf{q}') \Delta(\mathbf{q}) \frac{1}{E(\mathbf{q})} \tanh \left( \frac{1}{2} \beta E(\mathbf{q}) \right). \tag{1}$$

Here,

$$\Delta(\mathbf{p}) = (\Delta_{\sigma, \sigma'}(\mathbf{p}))_{\sigma, \sigma' \in \{\uparrow, \downarrow\}}$$

is a  $2 \times 2$  matrix satisfying

$$\Delta(\mathbf{p}) = -\Delta(-\mathbf{p})^T$$

and

$$E(\mathbf{q})^2 = e(\mathbf{q})^2 + \Delta(\mathbf{q})^* \Delta(\mathbf{q}).$$

The expression  $\frac{1}{E(\mathbf{q})} \tanh \left( \frac{1}{2} \beta E(\mathbf{q}) \right)$  is unambiguously defined by expanding  $\frac{1}{\sqrt{x}} \tanh \left( \frac{1}{2} \beta \sqrt{x} \right)$  as a power series in  $x$ . For a derivation of (1) see [AB], [BW].

Every solution of (1) is of the form

$$\Delta(\mathbf{p}) = (Y_{\sigma, \sigma'}(\mathbf{p})), \quad Y_{\sigma, \sigma'} \in H_\ell.$$

The simplest solutions are unitary and isotropic. A solution is unitary when

$$\Delta(\mathbf{p})^* \Delta(\mathbf{p}) = |d(\mathbf{p})|^2 I$$

and isotropic when  $d(\mathbf{p})$  is a constant. In this case the quasiparticle dispersion relation  $(\epsilon(\mathbf{q})^2 + |d|^2)^{\frac{1}{2}}$  is isotropic and has a gap  $|d|$  determined by

$$1 = \frac{1}{2} \int_{|\epsilon(\mathbf{q})| \leq \epsilon} \frac{d^3 \mathbf{q}}{(2\pi)^3} \lambda_\ell^{(h)} (\epsilon(\mathbf{q})^2 + |d|^2)^{-\frac{1}{2}} \tanh \left[ \frac{1}{2} \beta (\epsilon(\mathbf{q})^2 + |d|^2)^{\frac{1}{2}} \right] \quad (2)$$

when  $d \neq 0$ . Intuitively, they have the best chance of being stable.

There are two important examples of isotropic, unitary solutions. For  $\ell = 0$  there is the BCS model

$$\Delta = \begin{bmatrix} 0 & d \\ -d & 0 \end{bmatrix}$$

for phononic superconductivity. Balian and Werthamer discovered, in the  $\ell = 1$  sector, the solution

$$\Delta = d \begin{bmatrix} -\mathbf{p}_1 + i\mathbf{p}_2 & \mathbf{p}_3 \\ \mathbf{p}_3 & \mathbf{p}_1 + i\mathbf{p}_2 \end{bmatrix}, \quad \mathbf{p}_1^2 + \mathbf{p}_2^2 + \mathbf{p}_3^2 = k_F^2$$

which describes the B phase of  $\text{He}^3$ .

**Theorem** *There are no nontrivial, isotropic, unitary solutions of (1) for  $\ell \geq 2$ .*

One therefore expects that solutions will have nodes for  $\ell \geq 2$  making the flow harder to control. Such nodes are observed in the A phase of  $\text{He}^3$  and in the  $\ell = 2$  theory of heavy fermionic superconductivity. Nodes also appear in the gap function for systems with cubic symmetry. See, for example, [VG].

The proof of Theorem 1 follows immediately from the

**Lemma** *Let  $f, g \in H_\ell$  satisfy  $f\bar{f} + g\bar{g} = 1$  on  $S^2$ . Then,  $\ell = 0, 1$ .*

**Proof** Let  $P_\ell$ ,  $\ell \geq 0$ , be the homogeneous polynomials of degree  $\ell$  on  $\mathbf{R}^3$  with  $\text{SO}(3)$  invariant inner product

$$\langle f, g \rangle := f \left( \frac{\partial}{\partial k_1}, \frac{\partial}{\partial k_2}, \frac{\partial}{\partial k_3} \right) \bar{g}.$$

As usual  $H_\ell$  is identified with  $H_\ell^*$  by the  $\text{SO}(3)$  equivariant isomorphism

$$f \mapsto \langle \cdot, \bar{f} \rangle.$$

We shall show that under the hypothesis of the lemma

$$U = f \otimes \bar{f} + \bar{f} \otimes f + g \otimes \bar{g} + \bar{g} \otimes g$$

is the (unique up to scalars)  $SO(3)$  invariant element of  $H_\ell \otimes H_\ell$ . It follows that the homomorphism

$$U \in H_\ell \otimes H_\ell \cong H_\ell \otimes H_\ell^* \cong \text{Hom}(H_\ell, H_\ell)$$

commutes with  $SO(3)$  and is of rank at most four. Moreover, by Schur's Lemma,  $U$  is an isomorphism since  $H_\ell$  is irreducible. Consequently,  $2\ell + 1 \leq 4$ .

Consider the  $SO(3)$  equivariant multiplication map

$$H_\ell \otimes_s H_\ell \xrightarrow{M} P_{2\ell}$$

$$\sum c_j \phi_j \otimes \psi_j \mapsto \sum_j c_j \phi_j \psi_j.$$

Observe that

$$\dim H_\ell \otimes_s H_\ell = 2\ell + 1 + \frac{(2\ell + 1)(2\ell)}{2} = \binom{2\ell + 2}{2} = \dim P_{2\ell}$$

and

$$MU = 2|k|^{2\ell}.$$

If  $M$  is surjective it is an isomorphism and  $U$  is invariant.

The projection of

$$M \left( (k_1 + ik_2)^\ell \otimes_s (k_1 - ik_2)^\ell \right) = (k_1^2 + k_2^2)^\ell$$

onto the irreducible subspace  $|k|^{2(\ell-m)} H_{2m}$  of  $P_{2\ell}$  is nonzero because

$$\begin{aligned} & \left\langle |k|^{2(\ell-m)} (k_1 + ik_3)^{2m}, (k_1^2 + k_2^2)^\ell \right\rangle \\ &= \left( \frac{\partial}{\partial k_1} + i \frac{\partial}{\partial k_3} \right)^{2m} \Delta^{\ell-m} (k_1^2 + k_2^2)^\ell \\ &= \prod_{j=0}^{\ell-m-1} 4(\ell-j)^2 \left( \frac{\partial}{\partial k_1} + i \frac{\partial}{\partial k_3} \right)^{2m} (k_1^2 + k_2^2)^m \\ &\neq 0. \end{aligned}$$

Recall that every invariant subspace of  $P_{2\ell}$  is of the form

$$\oplus_{j_i} |k|^{2j_i} H_{2(\ell-j_i)}$$

with  $0 \leq j_1 < j_2 \cdots < j_r \leq \ell$  and in particular

$$P_{2\ell} = \bigoplus_{j=0}^{\ell} |k|^{2j} H_{2(\ell-j)}.$$

Finally the image of  $M$  is invariant and therefore all of  $P_{2\ell}$ . ■

We observe that in two dimensions there are unitary isotropic solutions of the gap equation for every angular momentum. For example,

$$\Delta(\mathbf{p}) = d \begin{bmatrix} \cos \ell\theta & \sin \ell\theta \\ \sin \ell\theta & -\cos \ell\theta \end{bmatrix}$$

when  $\ell$  is odd and

$$\Delta(\mathbf{p}) = d \begin{bmatrix} 0 & e^{i\ell\theta} \\ -e^{i\ell\theta} & 0 \end{bmatrix}$$

when  $\ell$  is even. Here,  $\mathbf{p} = |\mathbf{p}|(\cos \theta, \sin \theta)$ .

#### References

- [AB] P.W. Anderson and W.F. Brinkman, Theory of Anisotropic Superfluidity in "Basic Notions of Condensed Matter Physics" by P.W. Anderson, Benjamin/Cummings, Menlo Park (1984).
- [BW] R. Balian and N.R. Werthamer, Superconductivity with Pairs in a Relative p Wave, Phys. Rev. 131, 1553-1564 (1963).
- [FT] J. Feldman and E. Trubowitz, The Flow of an Electron-Phonon System to the Superconducting State, to appear in Helvetica Physica Acta.
- [KL] W. Kohn and J.M. Luttinger, New Mechanism for Superconductivity, Phys. Rev. Lett. 15, 524-526 (1965).
- [VG] G.E. Volovik and L.P. Gor'kov, An Unusual Superconductivity in  $UBe_{13}$ , JETP Lett. 39, 674-677 (1984).