

Paramagnetic resonance of Fe³⁺ in some feldspars

Autor(en): **Marfunin, A.S. / Bershov, L.V. / Meilman, M.L.**

Objekttyp: **Article**

Zeitschrift: **Schweizerische mineralogische und petrographische Mitteilungen
= Bulletin suisse de minéralogie et pétrographie**

Band (Jahr): **47 (1967)**

Heft 1: **Feldspäte**

PDF erstellt am: **05.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-36927>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Paramagnetic Resonance of Fe³⁺ in some Feldspars

By *A. S. Marfunin, L. V. Bershov, M. L. Meilman* (Moscow)*)
and *J. Michoulier* (Grenoble)**)

With 4 figures and 2 tables in the text

Abstract

The EPR spectrum of Fe³⁺ in oligoclase, microcline and some other feldspars has been investigated at 35000 Mc/s and 9500 Mc/s. The parameters of the Hamiltonian spin were calculated. A comparison of the orientation of the g-factors and of the principal directions of electric field gradients leads to the assignment of Fe³⁺ to the T₁ (o) site.

INTRODUCTION

The electron paramagnetic resonance (EPR) investigation of Fe³⁺ in feldspars is related to many problems.

Firstly, it is the most direct method of obtaining complete data about the position of iron impurities in feldspars. This technique has enabled a great deal of new information concerning this old problem to be obtained (COOMBS, 1960).

Still more interesting is the use of Fe³⁺ ions as a kind of probe giving information about the interaction of the Fe³⁺ ions with their surroundings.

If the iron ions enter feldspars substitutionally for the Al³⁺ ions and if Fe³⁺ follows Al³⁺ as regards the Al \rightleftharpoons Si distribution among tetrahedral sites, it would then be possible to use Fe³⁺ for investigating order-disorder relations, composition changes and twinning in members of the feldspar group.

*) Institute for ore deposits, Geology, Petrology, Mineralogy and Geochemistry, Moscow.

**) Laboratoire de Physique générale et Spectrométrie physique, Institut Fourier, Grenoble.

From the EPR point of view the complicated series of feldspar solid solutions with their order-disorder relations are highly suited for studying the further possibilites of this method. The triclinic local symmetry of Al-O₄ tetrahedra permits an investigation of the interaction of paramagnetic ions with non rhombic crystal fields.

Feldspars are compounds which like corundum, spinel, beryl (G. BURNS and W. J. NICHOLSON, 1963), spondumene (A. MANOOGIAN, F. HOLUJ, and J. W. CARSWELL, 1965) permit the comparision of crystal field by several methods: the values of eQq/h of Al³⁺ host atoms from electric quadrupole resonance spectra (ST. HAFNER and P. HARTMANN, 1964), the values of eQq/h of Fe⁵⁷ impurity atoms from the Mössbauer techniques; and the terms in the EPR Hamiltonian spin of paramagnetic ions Fe³⁺, Cr³⁺, replacing Al³⁺.

The single-crystal spectrum of the Fe³⁺ in Al-O₄ feldspar tetrahedra can be used in connection with the interpretation of tetrahedral Fe³⁺ EPR spectra in glasses (T. G. CASTNER a. o., 1960) and in powders (E. BOESMAN and D. SCHOEMAKER, 1961).

The EPR spectrum of Fe³⁺ in microcline has recently been measured by U. HÖCHLI (1964) at frequencies 8880, 9990 and 11340 Mc/s. However, the spectra at X-band frequencies (about 9500 Mc/s) are very complicated and the analysis of the spectrum in this band is extremely difficult as may be seen from Figures 1, 2 and 3 of this paper.



We have investigated the spectra of Fe³⁺ in oligoclase (A. S. MARFUNIN and J. MICHOULIER, 1966) and microcline at frequencies 35000 Mc/s (Q-band) and 9500 Mc/s (X-band). Preliminary results have been obtained for low sanidine and some other feldspars.

In an earlier paper the applications of EPR to mineralogy were considered (A. S. MARFUNIN, 1965).

EXPERIMENTAL PART

Measurements were made with an Q-band spectrometer (Institut Fourier, Grenoble) and an X-band spectrometer (Institute for Ore Deposits Geology, Petrology, Mineralogy and Geochemistry, Moscow).

Two different techniques were used in the determination of the principal directions of the magnetic axes: 1. by finding the extreme positions of resonance lines displayed on an oscilloscope, and 2. by plotting curves of equal magnetic field on a Fedorov sphere (B. I. ZUKERMAN, M. L. MEILMAN, L. P. SOROKINA, 1962).

Fig. 1. Energy-level diagram for Fe^{3+} in microcline.Fig. 2. Energy-level diagram for Fe^{3+} in oligoclase.

The data obtained from the measurements were treated in an "Strela" computer program.

The investigated specimens were as follows:

1. Oligoclase from pegmatites, North Karelia, Ab_{90} , An_{10} , $2V = +80^\circ$, white with blue schiller; only very thin albite twins.
2. Microcline from nepheline syenites, Khibin massiv (Kola peninsula), $2V = -82^\circ$, $\perp(010)$ $\text{Ng} = 18,5^\circ$, $\perp(001)$ $\text{Nm} = 12^\circ$. Large microcline twins. Colour white to grey.
3. Low sanidine-“Fe-orthoclase” from Madagaskar, $2V = -38^\circ$ in $\perp(010)$. Colourless, semi-transparent.

The iron content is 0,0n% in oligoclase, 0,n% in microcline and 0,5—p,7% in sanidine (spectrographic analysis).




Fig. 3. Angular dependence of the EPR spectrum of Fe^{3+} in oligoclase at 9500 Mc/s in the planes Z-x, x-y, y-z of the magnetic axes.

RESULTS

A. Oligoclase and microcline

The EPR spectra of Fe³⁺ in oligoclase and microcline are described by the following expression for the Spin Hamiltonien (W. LOW and E. L. OFFENBACHER, 1965; M. M. ZARIPOV and L. JA. SHEKUN, 1964).

$$H = \frac{1}{3} b_2^0 O_2^0 + \frac{1}{3} b_2^2 O_2^2 + \frac{1}{60} b_4^0 O_4^0 + \frac{1}{60} b_4^2 O_4^2 + \frac{1}{60} b_4^4 O_4^4 \\ + \beta (g_x H_x S_x + g_y H_y S_y + g_z H_z S_z).$$

The Spin Hamiltonien parameters are given in Table 1. ($\pm 1,0$ gauss for b_n^m and $\pm 0,01$ for g-factors.)




Fig. 4. Stereographic projection showing the orientation of the magnetic axes and of the principal axes of the electric field gradient.

Mr_Q-microcline, Khibin massiv, EPR, Q-band. Mr_x-microcline, Khibin massiv, EPR, X-band. Mr_H-microcline (U. HÖCHLI, 1964), EPR, X-band. Mr_{Al}-microcline, electric field gradient at Al-sites (St. HAFNER and P. HARTMANN, 1965). Ab_{Al}-albite, electric field gradient at Al-sites (St. HAFNER and P. HARTMANN, 1965).

Olig-oligoclase, EPR, Q-band.

¹⁾ The parameters obtained by U. HÖCHLI (1964) for Fe³⁺ in microcline are $b_2^0 = 1130$ gauss, $b_2^2 = 543$ gauss, $b_4^4 = 1500$ gauss. There is no description of this microcline.

Table 1. *Spin Hamiltonien for Fe³⁺ in oligoclase and microcline (gauss)*

	b ₂ ⁰	b ₂ ²	b ₄ ⁰	b ₄ ²	b ₄ ⁴	g _x	g _y	g _z
oligoclase	1927	383	9	10	25	2,001	2,001	2,005
microcline	1138	1198	6	17	75	2,00	2,00	2,00

The values of the energy levels for the three magnetic axes were calculated and plotted in Figs. 1 and 2. The observed Q-band and X-band transitions are shown. In addition to the "allowed" iron lines, there are also "forbidden" lines.

The calculated angular variation of the spectrum Fe³⁺ in oligoclase in the planes zy, yz, xz of the magnetic axes at X-band frequency is shown in Fig. 3.

Fig. 4 shows the direction of the magnetic axes, and table 2 gives the direction cosines with respect to a set of orthogonal axes: a=[100], b=±(010), c=±[100] || (010).

Table 2. *Direction cosines of the magnetic axes and of the principal axes of eQq/h*

	a	b	c	Method	Reference
Oligoclase	g _x 0,587	-0,493	0,643	EPR Q-band	This paper
	g _y -0,697	0,073	0,713		
	g _z 0,388	0,871	0,301		
Microcline	g _x 0,692	-0,466	0,552	EPR Q-band	This paper
	g _y -0,621	0,042	0,783		
	g _z 0,406	0,871	0,275		
Microcline	g _x 0,653	-0,569	0,503	EPR X-band	This paper
	g _y +0,574	0,079	0,816		
	g _z 0,496	0,824	0,274		
Microcline	g _x —	—	—	EPR X-band	HÖCHLI, 1964
	g _y —	—	—		
	g _z -0,57	0,82	0,07		
Microcline	V _{xx} 0,007	-0,356	0,934	NMR	ST. HAFNER and P. HARTMANN
	V _{yy} 0,888	-0,428	-0,170		
	V _{zz} 0,461	0,831	0,313		
Albite	V _{xx} 0,788	-0,488	0,375	NMR	ST. HAFNER and P. HARTMANN
	V _{yy} -0,551	-0,288	0,783		
	V _{zz} 0,274	0,824	0,496		

A single site only of Fe^{3+} is found both in oligoclase and microcline. Two spectra of Fe^{3+} in microcline show the same relation as the microcline twins.

The linewidths are about 20 gauss for $-1/2 \rightarrow +1/2$ transition, 60 gauss for $\pm 1/2 \rightarrow \pm 3/2$, 80 gauss for $\pm 3/2 \rightarrow \pm 5/2$ both in oligoclase and microcline.

B. Low-sanidine ("Fe-orthoclase" from Madagaskar)

At frequency 35.000 Mc/s only the $1/2 \rightarrow -1/2$ transition could be observed, but there are two magnetically inequivalent sites. The linewidths are very considerable: 100, 150 and 350 gauss for the principal directions (12475, 12485 and 12200 gauss).

CONCLUSION

The orientation of the g-factors obtained from this investigation was compared with the directions of the principal axes of electric field gradients available from the nuclear magnetic resonance investigation (Fig. 4).

The similarity of the positions of the z-axes in microcline, oligoclase and albite would seem to indicate that Fe^{3+} is located at the Al-sites in the unit cell of the ordered feldspars. This leads, in other words, to the assignment of Fe^{3+} to the tetrahedral $T_1(0)$ sites. Changes from order to disorder are accompanied by random variations of the atom coordinates and by the consequent spreading of the directions of the magnetic axes. This gives rise to a broadening of the EPR-lines from about 20 gauss in microcline and oligoclase to 350 gauss in low-sanidine.

After the identification and calculation of the spectrum of Fe^{3+} in microcline, oligoclase and sanidine, many samples of feldspars were tested for Fe^{3+} using EPR at frequency 9500 Mc/s. Concentrations of 0,n – 0,00 n % of tetrahedral Fe^{3+} were observed in sanidines, orthoclases, microclines (white and red), amazonites, albites, oligoclases, labradorites, anorthites from different localities.

Thus, Fe^{3+} ions are shown to be common as substitutional impurities in feldspars and can be used for determining the composition, degree of order-disorder, and twinning in this group.

Further investigations of EPR of Fe^{3+} in feldspars are now in progress.

Acknowledgements

The authors are indebted to Prof. M. Soutif and Prof. P. Dueros for their interest, to R. M. Mineeva for help in calculating and to Dr. V. Gabis and B. Borutski for the specimens of Fe-orthoclase and microcline.

REFERENCES

- BOESMAN, E. et SCHOEMAKER, D. (1961): Resonance paramagnétique de l'ion Fe^{3+} dans la kaolinite. C. R. Acad. Sci., Paris, 252.
- BURNS, G. and NICHOLSON, W. J. (1962): Crystal field comparisons. In "Magnetic and electric resonance and relaxation". Proc. XIth Colloque Ampère, p. 461 to 465, Amsterdam.
- CASTNER, T. G., NEWELL, G. S., HOLTON, W. C. and SLICHTER, C. P. (1960): Note on the paramagnetic resonance of iron in glass. J. Chem. Phys., 32, 668—673.
- COOMBS, D. S. (1954): Ferriferous orthoclase from Madagaskar. Min. Mag., 30, 226, 409—427.
- HAFNER, St. und HARTMANN, P. (1964): Elektrischer Feldgradient und Sauerstoff-Polarisierbarkeit in Alkali-Feldspäten ($\text{NaAlSi}_3\text{O}_8$ und KAlSi_3O_8). Helv. Phys. Acta, 37, 348—360.
- HÖCHLI, U. (1964): Electron spin resonance of Fe^{3+} in feldspars. In "Electronic magnetic resonance and solid dielectrics". Proc. XIIth Colloque Ampère, p. 191—197. North-Holland Publishing Company, Amsterdam.
- LOW, W. and OFFENBACHER, E. L. (1965): Electron spin resonance of magnetic ions in complex oxides. In: Solid State Physic, v. 17, 136—216, Academic Press, N. Y. and L.
- MANOOGLIAN, A., HOLUJ, F. and CARSWELL, J. W. (1965): The electron spin resonance of Fe^{3+} in single crystals of spodumene. Canad. J. Phys., 43, 2262—2275.
- MARFUNIN, A. S. (1965): Radiospectroscopy of minerals. Geol. J., 5.
- MARFUNIN, A. S. et MICHOUPLIER, T. (1966): Résonance paramagnétique électrique de l'ion Fe^{3+} dans un monocrystal d'oligoclase. C. R. Acad. Sci. Tome 262.
- ZARIPOV, M. M. and SHEKUN, L. JA. (1964): Electronic paramagnetic resonance in crystals. In: Paramagnetic resonance, p. 5—41, Kazanj.
- ZUKERMAN, B. J., MEILMAN, M. L., SOROKINA, L. P. (1962): Radiospectroscopic orientation of crystals. Kristallografia, 7, N. 2, 224—228. (Engl. transl. in: Sov. Physics — Crystallographie.)

Manuscript received June 30th, 1966.