

Miscellanea : Taxonomic research on the etiology of hookworm : disease occurring on the shores of the Black Sea, Turkey

Autor(en): **Oytun, H. ükrü / Güralp, Nevzat**

Objekttyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **16 (1959)**

Heft 1

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-310806>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

Taxonomic Research on the Etiology of Hookworm Disease occurring on the Shores of the Black Sea, Turkey.

By H. ŞÜKRÜ OYTUN and NEVZAT GÜRALP.

University of Ankara, Faculty of Medicine, Department of Parasitology
(Director: Prof. Dr. H. S. Oytun).

During the summer of 1951, one of us visited the Turkish provinces on the shores of the Black sea and studied the hookworm problems of the area. Special attention was given to the occurrence and identification of these parasites and to prophylactic measures being taken. A report of the findings was submitted to the Ministry of Health and Social Help and later results of the work were published in both Turkish and German.

A taxonomic study has been made both of the parasites we obtained from infected people and of those sent to us later by Dr. MUHARREM, Director of Health and Social Help of Trabzon. This paper reports the results of this study.

Materials and Methods.

The specimens were obtained from stools of patients who had been treated with carbon tetrachlorid at either Trabzon State Hospital or the Necator Control Dispensary at Rize. The infected individuals were originally from Akçaabat, Vakfıkebir, Of, Sürmene, Yumra, Pazar, Tirebolu, Görele, Trabzon, Çaykaya and Giresun, the majority being from Akçaabat, Vakfıkebir and Of. They were from 10 to 50 years old.

Stools were mixed with water and the parasites were removed by means of a fine brush or a needle. They were washed first in tap water, then in 10% saline solution, a brush being used to clean them properly. They were fixed in 10% formalin and preserved in 70% alcohol-glycerin solution. They were cleared in Amann's lactophenol.

Results.

Hookworms were collected from the stools of 61 patients; 55 females and 6 males. All told, 1,869 hookworms were collected and identified.

Of these 1,791 (95.83%) were identified as *Necator americanus* (STILES 1902) and 78 (4.17%) as *Ancylostoma duodenale* (DUBINI 1843). No specimens of any of the other three species known to occur in man, namely *Ancylostoma caninum* (ERCOLANI 1859), *A. brasiliense* (DE FARIA 1910), and *A. malayanum* (LANE 1916) were found.

Only *N. americanus* occurred in 48 (78.69%) of the 61 hookworm infected patients. Both *N. americanus* and *A. duodenale* were found in the other 13 (21.31%) infected patients.

The largest number of hookworms collected from a patient was 55, of these, 39 were *N. americanus* and 16 *A. duodenale*.

The largest number of *N. americanus* collected from a single individual was 52 and of *A. duodenale* 16.

Discussion.

Workers in other countries have reported that women are more susceptible to hookworms than men are. Our findings tend to substantiate this.

A. caninum is known to occur in dogs in Turkey. Since this parasite has been found in human beings in other countries, it may be that a larger series of examinations would result in finding it as a parasite of man in Turkey also.

Summary.

A taxonomic survey has been made on the hookworms collected from the infected patients on the Black sea shores of Turkey and diagnosed as *Necator americanus* and *Ancylostoma duodenale*. *N. americanus* is much more prevalent than *A. duodenale*.

References.

1. AKIL MUHTAR (1926). Türkienin Karadeniz vilâyetlerinde Necator. İstanbul Tıp Fak. Mec. 8, 321-325.
 2. BELDING, D. (1942). Textbook of clinical Parasitology. New York.
 3. BRUMPT, E. (1949). Précis de Parasitologie. Paris.
 4. CHANDLER, A. C. (1955). Introduction to Parasitology. New York: John Wiley and Sons Inc..
 5. FAUST, E. C. (1949). Human Helminthology. Philadelphia.
 6. GARIN, CH., ROUSSENT, J. et GONTHIER, B. (1932). L'Ankylostomose. Paris: Masson et Cie.
 7. CRAIG CH. FR. and FAUST, E. C. (1948). Clinical Parasitology. Philadelphia.
 8. İSMAİL HAKKI (1926). Ankilostom-Nekator. İstanbul Tıp Fak. Mec. 8, 449-473.
 9. MİMOĞLU, M. ve AKYOL, M. (1955). Hatay Vilâyetin de Ancylostomiasis üzerinde çalışmalar. A. Ü. Vet. Fak. Dergisi. C. 2, No. 1/2.
 10. NEDİM NİMET (1936). Menşei Zonguldak olan 3 Necator vakası münasebeti ile. Anadolu Klin. Yıl 4. No. 2, S. 115-117.
 11. NEVEU-LEMAIRE (1936). Traité d'Helminthologie Médicale et Vétérinaire. Paris.
 12. ONAT, A. R. (1932). Necatoriase. İstanbul.
 13. OYTUN, H. Ş. (1949). Tıbbî Parazitoloji. Ankara.
 14. OYTUN, H. Ş. (1953). Genel Parazitoloji ve Helmintoloji. Ankara.
 15. OYTUN, H. Ş. (1953). Türkiyede Ancylostomiasis (Kancalı kurt) problemine dair araştırmalar. Türk İj. Tecr. Biyol. Derg. Cilt XIV. 1, 19-36.
 16. OYTUN, H. Ş. (1954). The problem of Hookworm in Turkey. Acta med. tunc. 5, 27-33.
 17. ÖKTEM, Z. (1948). Ancylostomiasis ve Epidemiyolojisi. İstanbul Tıp Fak. Monografi. S. 4.
 18. SPREHN, C. (1932). Lehrbuch der Helminthologie. Berlin.
 19. STOLL, N. R. (1947). This wormy world. J. Parasit. 33, 1-19.
 20. UNAT, E. K. (1953). Tıbbî Helmintoloji. Ders kitabı. İstanbul.
 21. VASIF, H. (1928). Ankilostomiyaz, Nekatoriyaz. T.C. Sıhhiye ve Muaveneti İctimaiye. No. 28. Ankara.
 22. WATSON, J. M. (1953). Human Trichostrongylosis and its relationship to Ancylostomiasis in southern Irak, with comments on world incidence. Parasitology. 43, 102-109.
-