

**Zeitschrift:** Acta Tropica  
**Herausgeber:** Schweizerisches Tropeninstitut (Basel)  
**Band:** 19 (1962)  
**Heft:** (7): Pests of crops in warm climates and their control

**Kapitel:** Tobacco, pyrethrum, spices and drugs

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 14.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**TOBACCO, PYRETHRUM,  
SPICES AND DRUGS**

**Tobacco**

**Pyrethrum**

**Pepper**

**Chillies**

**Quinine**

Tobacco ( <i>Nicotiana sp.</i> )	Pest	Product	Formulation and Dosage	Where and when to apply
677	<i>Meloidogyne sp.</i>	143	L: 120 l/ha to a depth of 20-25 cm at planting station or in the row	Field: 2-3 weeks before transplanting the tobacco seedlings
		146	L: 60 l/ha to a depth of 20-25 cm at planting station or in the row	Field: 2-3 weeks before transplanting the tobacco seedlings
		143	L: 550 l/ha to a depth of 25 cm at 40 cm intervals	Seedbed: Several weeks before planting
		146	L: 350 l/ha to a depth of 25 cm at 40 cm intervals	do.
		149	L: 500 cc/sq.m (cover the bed with polythene; control nematodes, weeds and some soil borne fungi)	do.
678	<i>Melolonthinae</i> <i>Rutelinae</i>	7	G 10%: 24 kg/ha	Broadcast on soil surface when preparing for planting and work in 6-10 cm
		10, 16	WP 40%: 6 kg/ha	
			G 10%: 60-80 kg/ha	
		13	WP 50% or ES 50%: 12 kg/ha ES 20%: 2 l/200 l water	
679	<i>Drasterius sp.</i> <i>Heteroderes sp.</i> <i>Agriotes sp.</i>	10	WP 50%: 3 kg/3000 l water/ha	Around the stems when plants are transplanted so that the soil is wet (setting-water treatment)
		16	WP 25%: 3 kg/3000 l water/ha	
		7	WP 25%: 10-12 kg/500 l water/ha	Broadcast on soil surface 1 month before planting and work in 15-20 cm
		10	WP 50%: 10 kg/500 l water/ha	
		13	WP 50%: 4-5 kg/500 l water/ha	
		17	G 5%: 10 kg/ha	
		27	G 10%: 20 kg/ha	

- 680 *Gryllotalpa africana* see *Gryllotalpa* (No. 7)
- 681 *Scapteriscus vicinus* see *Gryllotalpa* (No. 7)
- 682 *Gonocephalum acutangulum* **7**  
**10**  
WP 40%: 1.5 kg/6000 l water/ha  
WP 50%: 2.4 kg/6000 l water/ha  
see also *Drasterius* (No. 679) and *Opatrum* (No. 683)
- 683 *Opatrum sabulosum* **7**  
D 2.5%: 25 kg mixed with fertilizer  
see also *Drasterius* (No. 679) and *Gonocephalum* (No. 682)
- 684 *Scotia (Euroa, Agrotis) segetum* **7, 12**  
**13**  
D 5%: 40 kg/ha  
D 2%: 40 kg/ha  
WP 40%: 900 g/300 l water/ha  
WP 50%: 3 kg/300 l water/ha  
WP 50%: 450 g/300 l water/ha  
Bait = WP 50%: 100 g + bran 40 kg mixture, 40 kg/ha  
Bait = WP 40%: 200 g + 20 kg bran mixture  
(add sufficient water to make a soft porridge-like slurry)  
12.21 or 30.1 may also be used in this way
- 685 *Scotia (Agrotis) ypsilon* **9**  
Bait = D 5%: 10 kg + bran 125 kg mixture, dampen with water so that the materials are crumbly and not over-moist
- 686 *Feltia subterranea* see *Scotia* (No. 684)
- 687 *Brachytrypus membranaceus* see *Gryllulus* (No. 63) and *Brachytrypus* (No. 150)
- 688 *Paralixis truncatulus* **13**  
WP 50%: 2 kg/500 l water  
Spray stem when pests appear
- 60 cc to the plant, planted after the rains
- Put the material in the planting hole and mix with the soil before planting
- Apply dust to plant and surrounding soil
- Spray base of stem and surrounding soil
- Broadcast on soil surface
- Scatter the bait along the rows in the late afternoon so that it does not dry out before nightfall
- Broadcast on soil surface of 1 hectare

Pest	Product	Formulation and Dosage	Where and when to apply
689 <i>Trichobaris trinotata</i>	39 see also <i>Paralixus</i> (No. 688)	<b>ES 50%:</b> 750 cc/500 l water	On stems and leaves when first signs of damage are seen
690 <i>Phthorimaea heliopa</i>	12	<b>D 5%:</b> 40 kg/ha <b>WP 25% or ES 25%:</b> 3 kg/600 l water/ha <b>D 1%:</b> 40 kg/ha <b>ES 50%:</b> 300-450 cc/600 l water/ha	As preventive measure: Spray or dust advanced crops fortnightly As soon as first seedlings in seedbeds show swollen stems
691 <i>Nezara viridula</i>	13 35.3 see also <i>Cyrtopeltis</i> (No. 617) and <i>Nezara</i> (No. 841)	<b>WP 50%:</b> 1 kg/500 l water/ha <b>Dust 1%:</b> 25-30 kg/ha	On leaves and stems when pests appear
692 <i>Planococcus citri</i>	34 35 39	<b>ES 20%:</b> 1.25-1.5 l/750 l water/ha <b>ES 50%:</b> 2.5 l/750 l water/ha <b>ES 20%:</b> 1.5 l/750 l water/ha	When pests appear. Repeat after 10 days
693 <i>Ditylenchus dipsaci</i>	39	<b>ES 50%:</b> 750 cc/1000 l water/ha	On foliage and stems when first signs of damage are seen
694 <i>Epilachna 28-maculata</i>	12 13	<b>WP 50%:</b> 2 kg/500 l water/ha <b>WP 20%:</b> 2 kg/500 l water/ha	When beetles appear
695 <i>Lema bilineata</i>	12 see also <i>Epilachna</i> (No. 694)	<b>D 5%:</b> 20-25 kg/ha	When beetles appear
696 <i>Epithrix hirtipennis</i>	12 34	<b>D 5%:</b> 20-30 kg/ha <b>WP or ES 25%:</b> 1.5 kg/250 l water/ha <b>ES 25%:</b> 250 cc/250 l water/ha	The lowest leaves on the plant must be treated
697 <i>Mimaulus thesii</i>	7 13, 15	<b>WP 40%:</b> 1.25 kg/500 l water/ha <b>WP 50%:</b> 1 kg/500 l water/ha	When weevils appear

- 698 *Psara periusalis* **27** ES 60%: 500 cc/500 l water/ha  
When first signs of damage appear
- 699 *Phthorimaea operculella* **12** WP 50%: 3 kg/750 l water/ha  
As preventive measure: Spray every 14 days after first signs of damage appear  
When first signs of damage (mines) appear
- 700 *Diacrisia obliqua* see Thosea (No. 318)  
When caterpillars appear
- 701 *Protoparce sexta* **11** ES 25%: 4 l/500 l water/ha  
**21, 34** ES 25%: 2 l/500 l water/ha  
see also *Mimaulus* (No. 697)  
When caterpillars appear
- 702 *Heliothis (obsoleta) zea* **0** Bait = WP: 1 kg + maize meal  
9 kg (mixture) 10 kg/ha  
**11** Bait = WP 50%: 250 g + maize meal  
10 kg (mixture) 10 kg/ha  
**12** Bait = WP 50%: 250 g + maize meal  
10 kg (mixture) 10 kg/ha  
**21** D 5%: 20-25 kg/ha  
**30.1** WP or ES 50%: 2 kg/750 l water/ha  
ES 25%: 2 l/1000 l water/ha  
ES 60%: 0.5-1 l/1000 l water/ha  
To buds of plants when small caterpillars appear  
On buds and foliage; repeat at fortnightly intervals
- 703 *Prodenia litura* **12** WP 50%: 3-4 kg/750 l water/ha  
see also *Mimaulus* (No. 697) and *Prodenia* (No. 837)  
When caterpillars appear or first signs of damage are seen
- 704 *Laphygma exigua* see *Laphygma* (No. 660)
- 705 *Zonocerus elegans* **10** WP 50%: 2 kg/1000 l water  
**12** D 5%  
**13** WP 50%: 3 kg/1000 l water  
D 2%  
**35** WP 50%: 1 kg/1000 l water  
ES 50%: 3 l/1000 l water  
Cover a strip 10-15 m wide round the plot
- 706 *Catantops opulentus* see *Zonocerus* (No. 705)

Pest	Product	Formulation and Dosage	Where and when to apply
707 <i>Thrips tabaci</i>	27 39	ES 25%: 600-750 cc/500 l water ES 50%: 250 cc/500 l water	When first signs of damage appear
708 <i>Frankliniella fusca</i>		see also Thrips (No. 810)	
709 <i>Heliothrips haemorrhoidalis</i>		see Thrips (No. 810)	
710 <i>Anaphothrips obscurus</i>		see Scirtothrips (No. 527)	
711 <i>Cyrtopeltis tenuis</i>		see Thrips (No. 707 and 810)	
712 <i>Dicyphus minimus</i>	39	see Cyrtopeltis (No. 617) D 1%: 15-20 kg/ha ES 25%: 750-1000 cc/750 l water/ha	On foliage as needed
713 <i>Orosius argentatus</i>		see Empoasca (Nos. 636 and 814)	
714 <i>Empoasca tabaci</i>		see Empoasca (Nos. 636 and 814)	
715 <i>Bemisia tabaci</i>	28, 33.10	see Bemisia (No. 817) ES 40%: 625 cc/1200 l water/ha	When the plants look sickly and are full of very minute, white flying insects
716 <i>Trialeurodes tabaci</i>		see Bemisia (No. 817) and Bemisia (No. 715,	
717 <i>Myzodes persicae</i>	21 27 35 36 39 28 42	ES 25%: 1.25 l/750 l water/ha ES 25% or WP 25%: 1 l/750 l water/ha ES 25% or WP 25%: 1.25-2 l/750 l water/ha ES 20%: 1 l/750 l water/ha ES 50% or WP 50%: 300-500 cc/750 l water/ha ES 40%: 500 cc/750 l water/ha ES 50%: 400-800 cc/750 l water/ha	When aphids appear on plants. Do not apply insecticides for 5 days before transplanting

- 718 *Macrosiphon solani* see Myzodes (No. 717)
- 719 *Cyaneolytta pectoralis* see Mylabris (No. 856)
- 720 *Heliothis virescens* see Heliothis (No. 702)
- 721 *Heliothis assulta* see Heliothis (No. 702)
- 722 *Lasioderma serricornis* **2** synergised pyrethrum spray: Spray the room once a fortnight  
tel quel
- 13** WP 50%: 1 kg/100 l water } 100 cc/  
**17** WP 20%: 2.5 kg/100 l water } sq.m } Treatment of the walls of tobacco stores  
see also Balanogastris (No. 166)
- 723 *Setomorpha margalaestriata* see Lasioderma (No. 722)
- 724 *Ephesia elutella* see Lasioderma (No. 722)

Pyrethrum (*Chrysanthemum cinerariaefolium*)

Pest	Product	Formulation and Dosage	Where and when to apply
725 <i>Nysius</i> sp.	see Creontiades (No. 797)		On stems when pests appear
726 <i>Frankliniella dampfi</i>	see Thrips (No. 727)		
727 <i>Thrips tabaci</i>	<b>28, 48</b>	<b>ES 40%: 500 cc/500 l water/ha</b>	On foliage when pests appear. Repeat 2 weeks later
728 <i>Myzodes persicae</i>	see Myzodes (No. 717)		
729 <i>Tetranychus bucleni</i>	<b>28, 48</b>	<b>ES 40%: 500-750 cc/500 l water/ha</b> see Tetranychus (No. 820) or Thrips (No. 727)	Repeat after 14 days

			Pre-plant treatment
730	<i>Meloidogyne</i> sp.	see <i>Meloidogyne</i> (No. 101)	
731	<i>Radopholus similis</i>	see <i>Radopholus</i> (No. 102)	do.
732	<i>Disphinctus maesarum</i>	see <i>Heliothis</i> (No. 137)	
733	<i>Ferrisia virgata</i>	see <i>Ferrisia</i> (No. 72)	When mealybugs appear
734	<i>Saissetia coffeae</i>	see <i>Saissetia</i> (No. 455) and <i>Saissetia</i> (No. 488)	
735	<i>Lepidosaphes piperis</i>	<b>39</b> ES 50%: 750-1000 cc/1000 l water	On foliage and stems when scales appear. Repeat after 2 weeks
736	<i>Thosea sinensis</i>	see <i>Thosea</i> (No. 318)	
737	<i>Cricula trifenestrata</i>	see <i>Parasa</i> (No. 439)	
738	<i>Gynaikothrips Karny</i>	see <i>Thrips</i> (No. 810)	
739	<i>Elasmognathus greeni</i>	see <i>Habrochila</i> (No. 38)	
740	<i>Diplogomphus hewitti</i>	see <i>Habrochila</i> (No. 38)	
741	<i>Longitarsus nigrispennis</i>	<b>12</b> WP 50%: 4 kg/1000 l water <b>39</b> D 1%: 15 kg/ha	When beetles appear on plant. Repeat after 2 weeks
742	<i>Lophobaris piperis</i>	<b>13</b> WP 50%: 1 kg/1000 l water see also <i>Trichobaris</i> (No. 689)	On foliage and stems when weevils appear or when first signs of damage are seen
743	<i>Dasynus piperis</i>	see <i>Dasynus</i> (No. 352) and <i>Leptoglossus</i> (No. 529)	
744	<i>Elasmognathus hewitti</i>	see <i>Habrochila</i> (No. 38)	

Chillies ( <i>Capiscum sp.</i> )	Pest	Product	Formulation and Dosage	Where and when to apply
	745 <i>Meloidogyne</i> sp.	see	<i>Meloidogyne</i> (No. 304)	
	746 <i>Radopholus similis</i>	see	<i>Meloidogyne</i> (No. 304)	
	747 <i>Hodoterme</i> <i>mossambicus</i>	see	<i>Hodoterme</i> (No. 170)	
	748 <i>Scirtothrips dorsalis</i>	see	<i>Scirtothrips</i> (No. 664) and <i>Thrips</i> (No. 810)	
	749 <i>Empoasca lybica</i>	see	<i>Empoasca</i> (No. 814)	
	750 <i>Tetranychus</i> sp.	see	<i>Tetranychus</i> (No. 327)	
	751 <i>Anthonomus Eugeniae</i>	12	<b>D 5%:</b> 25-30 kg/ha <b>WP 50%:</b> 4 kg/1000 l water <b>WP 20% or ES 20%:</b> 10 kg/ 1000 l water	On foliage when weevils appear. Repeat weekly
	752 <i>Phthorimaea</i> <i>gudmanella</i>	22	<b>WP 20% or ES 20%:</b> 10-15 kg/ 1000 l water	On flowers and young pods when first signs of damage are seen
	753 <i>Zonosemata electa</i>	35	<b>WP 40%:</b> 4 kg/1000 l water	On foliage when flies appear or when first signs of damage are seen. Repeat at fortnightly intervals
	754 <i>Helopeltis westwoodi</i>	12	<b>D 5%:</b> 20-30 kg/ha <b>WP 50%:</b> 2-4 kg/1000 l water	On foliage when pests appear or when first signs of damage are seen

755 <i>Meloidogyne</i> sp.	see Meloidogyne (Nos. 1 and 101)	
756 <i>Coccus viridis</i>	see Saissetia (No. 455) and Saissetia (No. 488)	On trunks and leaves when scales appear
757 <i>Howardia biclavis</i>	see Howardia (No. 76)	
758 <i>Alcides cinchonae</i>	<b>WP 50%:</b> 5 kg/1000 l water <b>ES 20%:</b> 2-4 l/1000 l water	On leaves and twigs when weevils appear. Repeat after 2-3 weeks if necessary
759 <i>Helopeltis cinchonae</i>	<b>WP 50%:</b> 5 kg/1000 l water	On foliage when bugs appear
760 <i>Dactylispa manteroi</i>	<b>WP 50%:</b> 3 kg/1000 l water	On foliage when beetles appear
761 <i>Metanastria hyrtaca</i>	see Parasa (No. 79) and Parasa (No. 439)	
762 <i>Odonestis plagifera</i>	see Parasa (No. 79) and Protoparce (No. 701)	
763 <i>Deitephila nerii</i>	see Parasa (No. 79) and Protoparce (No. 701)	
764 <i>Boarmia crepuscularia</i>	see Parasa (No. 79) and Protoparce (No. 701)	
765 <i>Euproctis varia</i>	see Parasa (No. 79) and Protoparce (No. 701)	
766 <i>Heliothrips haemorrhoidalis</i>	see Scirtothrips (No. 527)	On first signs of damage
767 <i>Anaphothrips orchidii</i>	see Selenothrips (No. 85) and Scirtothrips (No. 527)	On first signs of damage
768 <i>Pachypeltis vittiscutis</i>	see Helopeltis (No. 754)	
769 <i>Tetranychus urticae</i>	see Tetranychus (No. 820)	
770 <i>Hemitarsonemus lata</i>	see Hemitarsonemus (No. 147)	On foliage when first signs of damage are seen

