Zeitschrift: Acta Tropica

Herausgeber: Schweizerisches Tropeninstitut (Basel)

Band: 25 (1968)

Heft: 2

Sonstiges

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 18.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

ACTA TROPICA Supplement nº 7

Pests of Crops in Warm Climates and their Control

by

R. WYNIGER

A handbook (size 17×24.5 cm) with about 600 photographs and hand drawings. 2 Vol. XXIV, 555 ± 162 pages. Artificial leather binding. Price 210.— Swiss francs.

Every year insects cause enormous damage to food plants all over the world. This is mainly the result of agriculture tending more and more to large areas of monocultures. The growing of only a few crops in a certain area, which disturbs its biocoenosis, and the continual improvement of the quality of crops to meet market demands are some of the factors which are responsible for the increase of insect attacks and consequently the reduction of yields.

The frequency of insect appearance depends on the presence of host plants and on the prevailing climate; warm and moist climates usually favour insect development. The great number of species and individuals which make up over 70% of the animal kingdom are a constant threat to all the crops, especially in the tropics, subtropics and hot countries of the temperate zone.

The purpose of control measures is the prevention or reduction of injuries done by insects. By permitting higher yields, pest control contributes to the economy of agriculture and thus helps to solve the world's problem of nutrition.

Methodical and successful pest control must be based on a thorough knowledge of the pests and of the damage they cause. The present book is intended for partical use, presenting in as condensed a form as possible the vast quantity of information which has appeared in publications all over the world. Its purpose is to assist agriculturists in hot countries of all the continents in their fight against plant pests.

The ideas embodied in this book were developed not only in the laboratory and at the work-desk, but, in particular, during the extended visits of the author to tropical Africa and Southern States of America. It, therefore, sums up the experience acquired by the author on the spot, from observing the mode of action of the pests in the actual plantations, by applying his extensive biological knowledge, in advising the planters, and by conducting his own practical experiments in pest-control with new methods and products. His camera was always with him and was able to record characteristic phases of damage and stages of pests: the author's own photographs, in fact, account for most of the about 600 excellent illustrations which are scattered throughout the book and which so very effectively support the text.

The first chapter gives a synopsis of the morphology, anatomy, development, and classification of insects and other pests. The second chapter enables the reader to identify and preserve them. The third and most extensive chapter describes briefly the morphology, ecology, and distribution of each pest that may be found on a crop and the damage it causes. Technical data concerning the formulation of chemical insecticides, and the construction and manipulation of application equipment are contained in the fourth chapter, which also deals with the action of insecticides on the plant, the organization and timing of phytosanitary campaigns, biological control, time of treatment, assessment of the effectiveness of chemicals and field trials, evidence of insecticidal residue on plants and stored products, and insect resistance to chemical substances. A short glossary of the terms used in the book, followed by conversion tables for weights, measures and temperature concludes this chapter.

APPENDIX

as a separate volume

2nd revised and enlarged edition

The fifth part is published as a separate appendix. It deals with control measures, precautions to be taken and advice to doctors as to therapeutical measures in case of poisoning. Since the publication of the 1st edition of the "Appendix to Pests of Crops in Warm Climates and their Control" in 1962, many new synthetic-organic compounds for general use specialized Pest Control have been developed and marketed.

The "Table of Active Ingredients" in the present edition has been compiled correspondingly and brought up-to-date. Moreover, chemosterilants, activators or synergists, repellents and rodenticides are included.

The Active Ingredients are numbered as in the 1st edition, with some exceptions. The names or synonyms of new compounds are in alphabetical order and given a decimal figure. While the sections "Precautions" and "Mixing Tables" have not been altered, the section "First Aid Measures" has been in parts.

Control methods and methods of application resulting from the rapid development of new substances with new properties have been taken into consideration as far as possible in the main section entitled "Recommendations for Control Measures".

Contents

I. Introduction

- 1. Structure, general characteristics and development of insects
- 2. Synopsis of the classification of the most important plant pests.

II. Identification of Plant Pests

- 1. How pests cause damage
- 2. How pests are identified in the field
- 3. Evidence of plant parasitic nematodes
- 4. Killing, preserving and packing insects
- 5. How to make simple microscope slide mounts.

III. Pests of Crops in Warm Climates

Beverages: Coffee, Cocoa, Tea, Kola-nut — Food Crops: Maize, Sorghum, Rice, Sugar cane, Beans, Cassava, Sweet Potato — Fruits: Pineapple, Date-palm, Banana, Papaw, Mango, Citrus — Oil Plants: Olive, Coco-nut an Oil-palm, Sesame, Castor, Ground-nuts — Tobacco — Pyrethrum — Spices and Drugs: Pepper, Chillies, Quinine — Rubber and Fibres: Rubber, Cotton, Sisal, Roselle (Hibiscus), Ramie, Jute, Kapok — Locusts — Termites — Ants.

IV. Methods and Equipment for Pest Control

Direct method

- 1. Mechanical control
- 2. Chemical control
- 3. Product formulation
- 4. Method of application
- 5. Action of pesticides on the plant
- 6. Organization of phytosanitary campaigns
- 7. Biological control
- 8. Timing of phytosanitary campaigns
- 9. Examination of insecticides and field trials
- 10. Assessment of insecticidal residues on plants and stored products. Method of assessing insecticide residues
- 11. Development and causes of resistance to insecticides
- 12. Glossary
- 13. Conversion tables for weights, measures and temperatures.


Appendix: V. Control Measures

- 1. Table of active ingredients
- 2. Precautions
- 3. First-aid measures
- 4. Mixing tables
- 5. Recommendations for control measures against the described pests.

RICE

stalk

215




Plants with dead hearts. Leaf sheaths stained yellow; stem breaks off easily, its inside being hollowed out and filled with excrement.

Chilo suppressalis Walk, (simplex Butl.) Purple lined borer,

LEPIDOPTERA; Pyralididae

see page 135 (Maize)



Young leaves withering. Foliage eaten away; hearts destroyed, causing death of plants. Leaf sheaths with bore-holes. Yield often greatly reduced, due to destroyed ears and withered inflorescences.

Scirpophaga innotata Walk. White rice borer.


LEPIDOPTERA; Pyralididae

216

White moth with a wing span of 25-30 mm, which flies at night and is attracted by light (optical bait = lamp). The eggs are fixed to the underside of the leaves and the egg-clusters covered with brown fluff. The creamy-white caterpillars feed on leaves or tunnel through the leaf sheaths into the central bud. The ears of older plants are gnawed away. Several (4-5) generations each season. The insects go through diapause before they pupate.

Distribution: Indonesia, Philippine Islands

COTTON



Frequent blotch mines on unopened bolls. Bore-hole 1-3 mm wide. Lint inside boll mostly destroyed. Bolls fall off. Seeds also damaged. Supervening boll-rot.

bolls

Pectinophora gossypiella Saund. Pink bollworm.

LEPIDOPTERA; Gelechiidae

835

Moth with yellowish-brown to ochrous and dark spotted forewings, expanding to 15-20 mm. The hindwings are silvery. The female deposits 200-400 eggs singly under the calyx, on terminal buds and on young immature bolls. The young caterpillars are creamy to yellow, older ones have two pink transverse dorsal bands on each segment. Head and prothoracic shield are dark brown. The caterpillars bore into the bolls feeding on lint and seeds. The larval development is rapid (about 30 days). Pupation takes place inside the bolls, seeds or underground. Total time of development: 5-6 weeks. The pupal stages of the insect can be disseminated with the seeds. Several generations.

Distribution: wherever cotton is grown



Half ripe, unopened bolls show boreholes surrounded by superficial mining. Damage similar to that caused by P. yossypiella. Seeds inside bolls attacked.

Argyroploce leucotreta Meyr. False codling moth (Orange moth).

LEPIDOPTERA; Tortricidae

836

518

Moth with variegated brown and grey forewings, a white spot in the centre, while the hindwings are light brown to grey. They expand to 18-20 mm. Oviposition takes place on half-ripe, unopened bolls. Peak in June. Several generations.

Distribution: Africa (widespread)

1. Table of Active Ingredients. Classification of Active Ingredients

Botanical	insec	tici	de	S	•	•	•	•	•	•	•	٠	٠	•	٠	٠	٠		10
Synthetic	-organ	nic	ins	sec	tic	ide	es.	141	ı	(2)			u	•		12		200	11
	chlor																		11
	phos	pho	ro	us	es	ter	S	•		200		N# (٠		•		•	12
	carba	ıma	ite	S	٠	•		•		•		•	•	٠	٠	•			18
	carba	izo	les		٠	•	٠	•		٠	8	•	٠			٠	*	٠	19
	phen																	•	19
	misc	. co	mĮ	001	ıno	ds	٠	*	٠		•	٠	•	٠	٠	ŀ	**	•	20
Inorganic	insec	tici	de	s	٠	•		•	•	•			٠	٠	•	•	*		20
Mineral o	ils .		٠	•		٠	٠			(80)	٠		ĸ	100		1.0		•	20
Fumigant	s .	٠	9 . 0				•		•		•				•	0.00		(4)	20
Synthetic	-orgai	nic	ac	ari	cio	les				•		30 * 10	•	•		3.50			21
J	chlor																		21
	phos																		21
	carba	ıma	ite	S										0.00		1.0		1.0	22
	sulfu	r co	om	ро	un	ds				150						18.00		•	22
	misc.	co	mĮ	ot	me	ls	•1	9.	•		٠	•	*	2.00	٠			1.0	23
Inorganic	acari	cid	es	ě	•		*	•	ř	•	٠	•		•	•			٠	23
Attractan	ts.	*	•	8	•8	14	ě	*	٠	14	٠	- T	ě		*			•	24
Repellent:	s .		*1			•	*8			1005		100		٠	*	111		100	24
Chemoste	rilant	s	•			18	•		*	i.	•	696	•	S. 1.10	,	1.5			26
Activator:	s or S	Syn	erț	jis	t s			8	•	•		•	•	•	Ť	•	·	•	27
Molluscoo	ides	•	į.					8	20	*	¥	(3 4)		•	•	٠	٠	•	28
Nematoci	des	: •	es:			*	•		*	•	*	**	*	**				36%	28
Rodentici	des			12	20			_				12							29

CONTROL MEASURES

	on name or chemical or other name used	Mode of action *	Acute oral toxicity DL 50 (mg/kg rats)	** Use against
16 .3	Isodrin	C, St	7–17	biting insects
16.4	Kepone see Chlordecone			
17	Lindane (gamma-BHC)	C, St, V	125	pests in stored crops
18	Methoxychlor	C, St	6000	insects on fruits and vegetables (similar to DDT-insecticide)
18.1	Mirex	C, St	235-702	biting insects
19	Perthane	C, St	6600	biting insects on vegetables (caterpillars) and fruit flies
19.1	Prolan see Dilan			
19 .2	Rothane see DDD			
19 .3	Strobane	C, St	200 – 250	biting insects
19.4	TDE see DDD			
20	Telodrin (Isobenzan)	C, St	7	biting and sucking insects insects in soil
21	Thiodan (Endosulfan)	C, St	30–79	biting and sucking insects on various crops
22	Toxaphene	C, St	40 – 283	biting insects on fruits and vegetables
	Synthetic-organic inse	cticides: Ph	osphorous esters	
22.1	Abate	C, St	1000-3000	sucking and biting insects
22 .2	Alamos (Slam)	C, St	> 1500	sucking and biting insects
22 .3	Aphidan	C, St, S	86 (Mice)	sucking insects
22 .4	Azinphos(ethyl) see Ethyl-Azinphos			
22 .5	Azinphos(methyl) (Guthion) (Gusathion)	C, St, P	7–18	sucking and biting insects and spider mites
22 .6	Azodrin see Dimethyl phosphate of 3-hy- droxy-N-methyl-cis- crotonamide			
23	Baitex (Entex) (Fenthion) (Lebaycid)	C, St	200-250	fruit flies
23 .1	Bidrin see Dicrotophos			

-	-			
			n	п
		H		

Pest	Product	Formulation and Dosage	Where and when to apply
824 Anthonomus grandis	12+19.3	$\mathbf{ES25\%:4.51+ES50\%:4.51/5001}$	
	12+22	ES 25%: 4.51 + ES 50%: 4.51/5001	
	31	water/ha FS 90%: 9 8 1/500 1 water/ha	
	34	ES 25%: 11/500 1 water/ha	
	35	ES 50%: 21/5001 water/ha	
	37	ES 50%: 500 cc/500 l water/ha	
	38	ES 50%: 11/500 l water/ha	
	53	WP 80%: 1.25 kg/500 l water/ha	
	27.1, 42	ES 50%: 800-1000 cc/500 l water/ha	
825 Bucculatrix	15	ES 50%: 1 1/500 l water/ha	On foliage when first damage is seen
thurberiella	22	ES 50%: 12 1/500 l water/ha	
	2.57	ES 25%: 1.5-2 1/500 l water/ha	
	700	ES 50%: 1.25 1/500 l water/ha	
	39	ES 50%: 11/500 l water/ha	
	53	WP 80%: 3 kg/500 l water/ha	
826 Heliothis (obsoleta)	9+12	ES 25%: $51 + ES 25\%$: $51/5001$	Apply when an average of 5 small
zea		water/ha	CO
	10+12	$ ext{ES 25}\%: 41 + ext{ES 25}\%: 51/5001$	Repeat every 5-7 days until under
	;	water/ha	control
	- F	FS 90/6: 9 1/500 1 water/Ha	
	1	D 10%: 30 kg/ha	
	12+19.3	$ ext{ES 25}\%: 4.5 1 + ext{ES 50}\%: 4.5 1/500 1$	
	19 + 99	water/ha FS 25%: 451+ FS 50%: 4.51/5001	
		water/ha	
	15	ES 50%: 11/500 l water/ha	
	533	WP 80%: 1.5 kg/500 l water/ha	
	30.1	ES 60%: 11/500 l water/ha	
827 Earias insulana	31 + 34	$\mathbf{ES25\%;1.21} + \mathbf{ES25\%;31/5001}$	do.
		water/ha	

sec also Heliothis (No. 826)

ACTA TROPICA

In Vorbereitung | Articles en préparation | Articles in preparation

- Brack, C. Elektronenmikroskopische Untersuchungen zum Lebenszyklus von Trypanosoma cruzi unter besonderer Berücksichtigung der Entwicklungsformen im Überträger Rhodnius prolixus.
- Freyvogel, T. A., Honegger, C. G. & Maretić, Z. Zur Biologie und Giftigkeit der ostafrikanischen Vogelspinne Pterinochilus spec.
- Hecker, H. & Aeschlimann, A. Contribution à la connaissance des symbiotes chez Ornithodorus moubata (Ixodoidea). Etude au microscope électronique.
- Jadin, J. M. & Creemers, J. Etude de la pinocytose dans les formes en rosaces de Leptomonas ctenocephali.
- Jadin, J, M. & Creemers, J. Ultrastructure et biologie des Toxoplasmes.

 III. Observations de Toxoplasmes intraérythrocytaires chez un mammifère.
- Otieno, N. C. Further contributions to a knowledge of "Poroid Agarics" from the tropics: the genera Campanella P. Henn. and Dictyoploca (Mont.) Heim.
- Steiger, U. Intrazelluläre Mikroorganismen bei der Waldameise im Ei-, Larven- und Puppenstadium.
- Stöcklin, W.H. Kukukuku. Medical Patrol into one of the last restricted areas in the New Guinea Highlands.