

<b>Zeitschrift:</b>	Acta Tropica
<b>Herausgeber:</b>	Schweizerisches Tropeninstitut (Basel)
<b>Band:</b>	31 (1974)
<b>Heft:</b>	2
<b>Artikel:</b>	Miscellaneum : Viruslike particles of "Glossina fuscipes fuscipes" Newst. 1910
<b>Autor:</b>	Jenni, L. / Steiger, R.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-311958">https://doi.org/10.5169/seals-311958</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Miscellaneum

Swiss Tropical Institute, Basel

## Viruslike Particles of *Glossina fuscipes fuscipes* Newst. 1910

L. JENNI and R. STEIGER

### *Abstract*

The ultrastructure of viruslike particles forming pseudo-crystalline arrays in the nuclei of midgut epithelial cells of *Glossina fuscipes fuscipes* (Tsetse fly, Diptera) is described.

These spherical particles measure about 350–390 Å in diameter. Their number increases with the age of the flies.

### **Introduction**

Intranuclear viruslike particles (VLP) have been found in various cell types of *Drosophila*: they occur in fat bodies, oenocytes and central nervous tissue (PHILPOTT et al. 1969), in neurons and glia (HERMAN et al. 1971), as well as in salivary glands, accessory glands and muscles (MIQUEL et al. 1972) of imagines; FILSHIE et al. (1967) described them in cuprophilic cells of larvae, and WEHMAN & BRAGER (1971) in tissue cultured imaginal discs of the same insect genus.

Similar particles were seen in nuclei of muscle and hypodermal cells of a nematode (FOOR 1972).


During ultrastructural investigations on *Trypanosoma brucei* in organs of *Glossina* we regularly found VLP in different cell types of the vector. In *Glossina fuscipes fuscipes* these particles were restricted to nuclei of midgut epithelial cells. They are distinct from the cytoplasmic VLP found in *G. morsitans* (JENNI 1973).

### **Material and Methods**

*Glossina fuscipes fuscipes* (both sexes) emerged in the E.A.T.R.O. insectary (Tororo, Uganda) from pupae, which were either collected in the field (Lugala, Lake Victoria) or derived from a laboratory colony of the same origin. These flies got one infective bloodmeal (day 1 after emergence) on a rat showing a suitable parasitaemia of *Trypanosoma brucei*, and were subsequently fed daily on a clean ox and kept in a climatized (25 °C/80 % RH) dark fly room.

Flies (generally from day 18 on) were dissected; the midguts, salivary glands and reproductive organs were then processed for electron microscopy according to routine methods (STEIGER 1973).

Lead citrate and uranyl acetate stained serial sections (about 600 Å thick) were examined and recorded in a Zeiss EM 9 and Philips EM 300.



## Results

VLP can be seen in nuclei of midgut epithelial cells (Figs. 1–4). These measure 350–390 Å in diameter. They appear either "hollow" or contain electron-dense centrally disposed material (Figs. 2, 3 inset). The VLP form quasi-crystalline arrays up to 2.8 μm in diameter (Figs. 3, 4), which seem to be confined to the nucleolar region. The size of the arrays increases with age of the flies (between day 20 and 30); they are rather disperse in younger flies and compact in older ones. Inclusions have been found in practically all nuclei of the midgut epithelium.

Different stages of VLP development can be followed: first, single particles are detected in a specific zone adjacent to the nucleolus (Fig. 1). The number of the VLP then increases and single (Fig. 3) or multiple (Fig. 4) quasi-crystalline patterns arise. The zone of VLP formation is encircled by hetero-chromatin and electron-dense granules (Figs. 1, 2, 4). Later, the arrays occur free in a lighter nucleoplasm (Fig. 3).

## Discussion

The viruslike particles found in the present study resemble morphologically those described in *Drosophila* (MIQUEL et al. 1972; PHILPOTT et al. 1969), *Aedes aegypti* (FILSHIE et al. 1967) and in a nematode (FOOR 1972). The latter author noticed a similar association of the VLP with the nucleolar region of the nucleus.

Although cytopathic effects could not be seen with certainty in our material, it would be tempting to relate the presence of VLP to the high mortality of young *G. fuscipes* (within 10 days after emergence) and to the low hatching rate we observed. A lot of young flies were found to have disrupted midguts, and in one case, where we checked the midgut ultrastructure of a young weak imago, large intranuclear VLP aggregates already could be seen. The occurrence of "viruses" in old flies would then rather be regarded as a latent infection.

Concerning a possible transovarial transmission of the VLP, crystalline inclusions in the reproductive organs have never been found so far.

We shall try to clarify the nature of the intranuclear VLP using new fly material of the same origin.

---

Fig. 1. 20 days after emergence. First single VLP (→) in the specific zone (z) close to the nucleolus (n). Heterochromatin (h), electron-dense granules (g). 36,000 ×.

Fig. 2. 25 days after emergence. Formation of crystalline arrays (c) in the juxtap-nucleolar zone. The spherical VLP appear either "hollow" (→) or electron-dense (→). 36,000 ×.

Fig. 3. 30 days after emergence. Completed VLP crystal (→) free in the nucleoplasm. Rough endoplasmic reticulum (rer). 5250 ×. Inset: Detail from a VLP array. 90,000 ×.

Fig. 4. 30 days after emergence. Multiple crystalline inclusion (→) in the nucleoplasm. 12,000 ×.

*Acknowledgements*

This work was supported by the "Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung" (Grant Nr. 4.096.073).

We wish to thank the E.A.T.R.O. (East African Trypanosomias Research Organization, Tororo/Uganda) for providing laboratory facilities and fly material.

The authors gratefully acknowledge the technical assistance of Miss B. Baumgartner and Mrs. E. Ramseyer.

*References*

- FILSHIE, B. K., GRACE, T. D. C., POULSON, D. F. & REHACEK, J. (1967). Viruslike particles in insect cells of three types. – *J. Invert. Path.* 9, 271–273.
- FOOR, W. E. (1972). Viruslike particles in a nematode. – *J. Parasit.* 58, 1065–1070.
- HERMAN, M. M., JOHNSON, M. & MIQUEL, J. (1971). Viruslike particles and related filaments in neurons and glia of *Drosophila melanogaster* brain. – *J. Invert. Path.* 17, 442–445.
- JENNI, L. (1973). Viruslike particles in a strain of *Glossina morsitans centralis*, Machado 1970. – *Trans. R. Soc. trop. Med. Hyg.* 67, 295.
- MIQUEL, J., BENSCH, K. G. & PHILPOTT, D. E. (1972). Viruslike particles in the tissues of normal and  $\gamma$ -irradiated *Drosophila melanogaster*. – *J. Invert. Path.* 19, 156–159.
- PHILPOTT, D. E., WEIBEL, J., ATLAN, H. & MIQUEL, J. (1969). Viruslike particles in the fat body, oenocytes, and central nervous tissue of *Drosophila melanogaster* imagoes. – *J. Invert. Path.* 14, 31–38.
- STEIGER, R. F. (1973). On the ultrastructure of *Trypanosoma (Trypanozoon) brucei* in the course of its life cycle and some related aspects. – *Acta trop.* 30, 64–168.
- WEHMAN, H. J. & BRAGER, M. (1971). Viruslike particles in *Drosophila*: constant appearance in imaginal discs *in vitro*. – *J. Invert. Path.* 18, 127–130.