

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 32 (1975)
Heft: 2

Nachwort: Blood digestion in haematophagous insects : concluding remarks
Autor: Freyvogel, Thierry

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Concluding remarks

As has been said above the aim of publishing these four papers under a common heading is to reveal correlations between some of the aspects dealt with and so, hopefully, to come to a better understanding of the capability of numerous arthropods to digest blood and to transmit pathogens.

So far, qualitative electron microscopy has not revealed fundamental differences in the organization of midgut cells in blood sucking vs. non blood sucking insects. Recent morphometric investigations, on the other hand, have demonstrated quantitative ultrastructural differences between epithelial cells of the anterior and the posterior part of the midgut in the female *Aedes aegypti*. The data obtained could be correlated with the physiological results available and would allow the attribution of different functions to the two parts of the midgut mentioned (HECKER et al., *Cell. Tiss. Res.* 152, 31–49, 1974).

Morphometric analysis is actually being extended to compare the midgut cells of blood sucking with those of non blood sucking insects. It is to be elucidated, which cellular structures are specially involved in blood digestion, and possibly, in the transmission of pathogens. It remains to be shown whether there are representative quantitative differences between the two groups of insects concerned. So far, males and females of *Aedes aegypti* have been compared (RUDIN & HECKER, in prep.). The results showed some structural specialisations in females, which may be seen in the context of blood intake and digestion. For example, differences are found in the relative surface area (surface density) of rough endoplasmic reticulum (rer) membranes in the “stomach” cells, the values for females being significantly higher than those for males.

The rer is thought to be closely connected with enzyme production. In fact, a fairly good correlation has been shown to exist between the structural appearance of the rer and the activity of certain enzymes, mainly proteinases. It is felt, however, that concentration on proteinases may reveal only one part of the picture; in addition, attention should be paid to other enzymes which are present in gut cells and, at times, in the gut lumen, such as esterases, including those with an acetylcholine-like activity (GEERING & FREYVOGEL, *Comp. Biochem. Physiol.* 49B, 775–784, 1974).

What has been said in some of these papers on vector endocrinology shows very impressively how much an insect's physiology is affected by each blood meal. To us, a thorough knowledge of the processes concerned is essential for a real understanding of the degree to which a pathogen's life cycle may be interlinked with its vector's physiology, in other words to an understanding of the host/parasite relationship.

Acknowledgement

It is a pleasure to thank Dr. A. Diehl, Neuchâtel, Dr. K. Geering and Dr. H. Hecker, Basle, for their valuable help in preparing the section on physiology and biochemistry of parasitic arthropods and for useful comments with respect to the present publication.