Zeitschrift: Acta Tropica

Herausgeber: Schweizerisches Tropeninstitut (Basel)

Band: 34 (1977)

Heft: 1

Artikel: Experiments on trypanosome transmission by "Auchmeromyia" larvae :

preliminary note

Autor: Geigy, R. / Kauffmann, M.

DOI: https://doi.org/10.5169/seals-312252

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Experiments on trypanosome transmission by *Auchmeromyia* larvae¹

Preliminary note²

R. GEIGY, M. KAUFFMANN

During field surveys in the Serengeti National Park (SNP) from 1970 to 1975 (Geigy et al., 1971; Geigy and Kauffmann, 1973; Beglinger et al., 1976) a high incidence of *Trypanosoma* (T.) brucei was found in hyaena (Crocuta crocuta). As the tsetse flies are known to feed only rarely on these carnivores (Moloo et al., 1971; Rogers and Boreham, 1973), other infection routes were looked for, beside the possible oral infection by devoring infected prey (Moloo et al., 1973). In certain areas of the SNP (Mwanza Road) great numbers of blood sucking Auchmeromyia larvae were found in culverts, in close contact with hyaenas, and it was shown that these larvae occasionally feed on hyaenas (Geigy and Boreham, 1976; Boreham and Geigy, 1976). Therefore attempts were made to elucidate the role of these larvae as possible mechanical transmitters by experiments carried out at the Swiss Tropical Institute Field Laboratory in Ifakara (Tanzania), where these larvae are frequently found in local mud houses.

For these experiments 2 known strains of *T. brucei* were used:

- a) STIB 245, isolated from hyaena SNP 1971
- b) STIB 366 A, a derivative of S 42–030, Molteno Institute, Cambridge, originating from warthog, Uganda 1966.

Hungry larvae of the second and third stage were fed on infected nembutalized rats and then allowed to complete their blood meal within 1 h on a clean rat. The white rats (SIV) derived from a breeding station in Switzerland. All in all 22 clean rats were exposed to a double feed, of which 13 developed a patent parasitaemia after 3 to 17 days. The other 9 rats remained negative.

In 4 cases larvae were triturated 1 to 24 h after an incomplete, very small single meal on an infected rat. Trypanosomes from all of these larvae caused a parasitaemia 3 to 4 days after i.p. injection into clean rats.

Correspondence: Prof. Rudolf Geigy, Swiss Tropical Institute, Socinstrasse 57, CH-4051 Basel, Switzerland

¹ This study was supported by the Swiss National Science Foundation, Grant No. 3'2360'74.

² A detailed report will appear in «A Festschrift for Dr. C. A. Hoare's 85th birthday» in the Journal of Protozoology, 1977.

In 4 other cases, double fed larvae were triturated and the suspension injected into clean rats. But in this case, no parasitaemia occurred, which is probably due to the fact, that all viable trypanosomes taken up with the infective first meal were eliminated by the second feed.

This indicates that *Auchmeromyia* larvae present in the culverts of the Mwanza road in the Serengeti may act as mechanical transmitters.

- 1 Beglinger R., Kauffmann M., Müller R.: Culverts and trypanosome transmission in the Serengeti National Park (Tanzania). II. Immobilization of animals and isolation of trypanosomes. Acta trop. (Basel) 33, 68–73 (1976).
- 2 Boreham P. F. L., Geigy R.: Culverts and trypanosome transmission in the Serengeti National Park (Tanzania). III. Studies on the genus *Auchmeromyia* Brauer and Bergenstamm (Diptera: Calliphoridae). Acta trop. (Basel) 33, 74–85 (1976).
- 3 Geigy R., Mwambu P. M., Kauffmann M.: Sleeping sickness survey in Musoma District Tanzania. IV. Examination of wild mammals as a potential reservoir for *T. rhodesiense*. Acta trop. (Basel) 28, 211–220 (1971).
- 4 Geigy R., Kauffmann M.: Sleeping sickness survey in the Serengeti area (Tanzania) 1971. I. Examination of large mammals for trypanosomes. Acta trop. (Basel) 30, 12–23 (1973).
- 5 Geigy R., Boreham P. F. L.: Culverts and trypanosome transmission in the Serengeti National Park (Tanzania). I. Survey of the culverts. Acta trop. (Basel) 33, 57–67 (1976).
- 6 Moloo S. K., Steiger R. F., Brun R., Boreham P. F. L.: Sleeping sickness survey in Musoma District, Tanzania. II. The role of *Glossina* in the transmission of sleeping sickness. Acta trop. (Basel) 28, 189–205 (1971).
- 7 Moloo S. K., Losos G. J., Kutuza S. B.: Transmission of *Trypanosoma brucei* to cats and dogs by feeding infected goats. Trans. roy. Soc. trop. Med. Hyg. 67, 287 (1973).
- 8 Rogers D., Boreham P. F. L.: Sleeping sickness survey in the Serengeti area (Tanzania) 1971. II. The vector role of *Glossina swynnertoni* Austen. Acta trop. (Basel) 30, 24–35 (1973).