Zeitschrift: Acta Tropica

Herausgeber: Schweizerisches Tropeninstitut (Basel)

Band: 36 (1979)

Heft: 2

Artikel: In vivo and in vitro effect of cyclophosphamide on "Borrelia duttoni"

Autor: Wright, D.J.M.

DOI: https://doi.org/10.5169/seals-312513

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Department of Microbiology, Charing Cross Hospital Medical School, London

In vivo and in vitro effect of cyclophosphamide on Borrelia duttoni

D. J. M. WRIGHT

In memoriam Oscar Felsenfeld

Summary

Sera taken from mice given cyclophosphamide 75 min previously fails to substantially inhibit borrelial multiplication in vitro. Mice given cyclophosphamide were infected with *Borrelia duttoni*. Inability to demonstrate specific antibody in these mice did not prevent recovery from infection, resolution of the bacteraemia or appearance of an antigenic variant. Cyclophosphamide may delete suppressor lymphocytes allowing enhanced production of antibodies which may not only protect, but induce antigenic variation. Alternatively, recovery from infection may be unrelated to the immune response.

Key words: Borrelia duttoni; antigenic variation; antibody; cyclophosphamide.

Introduction

Specific antibody appears in relapsing fever when the bacteraemia is greatest and may be responsible for terminating the infection (Wright and Ginger, 1973). Antibody may prevent the borreliosis developing (Calabi, 1959) or abort an attack if passively transfused to a patient with the active disease (Balteanu et al., 1948). Alternatively as in trypanosomiasis (Vickerman, 1974), antibody might induce antigenic variation thereby perpetuating the infection. To determine which was the dominant role of antibody, experimental infections were studied in mice whose antibody production had been suppressed by cyclophosphamide.

Correspondence: D. J. M. Wright, Department of Microbiology, Charing Cross Hospital Medical School, Fulham Palace Road, London W6 8RF, England

Materials and methods

Organisms. The isolates of B. duttoni used were the Wellcome mouse adapted strain, originally obtained from Dr. J. D. Fulton at the National Institute for Medical Research, London, and a human pathogenic strain from Professor R. Geigy at the Swiss Tropical Institute, Basle (their reference number STIB 503). This strain had been repeatedly passaged in mice and ticks. Isolates were prepared from this strain either after one mouse passage (type 1) or after passaging through the tick Ornithodoros moubata and harvesting after subsequent mouse passage (type 2). Inocula of Borrelia were obtained by diluting infected mouse blood with citrated glucose saline (0.15M saline, 1% glucose, 0.5% citrate with Sorensens buffer to pH = 7.2). Spirochaetes were counted in a Helber counting chamber (Southee, 1972).

Serology. Slide agglutination was tested on a suspension containing 10⁸ fresh Borrelia in citrated saline. Borreliolysins were also assayed. Assessment was microscopical. Titres were established by determining the highest dilution of antisera at which lysis or agglutination occurred. Borrelial immunofluorescence was tested with goat anti-mouse IgM conjugates (Meloy Laboratories Inc., Springfield, USA) as used (Wright and Ginger, 1973). Antisera were prepared against Borrelia from the first, parental (P) and relapse, filial-one (F₁) parasitaemias on both the STIB 503 type 1 and type 2 strains by inoculating pure isolates into mice and collecting sera after recovery from infection had taken place. These specific sera were used in the agglutination and borreliolysin tests.

Mice. All mice used in this study were male or female, 5 to 6 weeks old, 12–15 g weight, inbred CBA/CA strain fed on water ad libitum and on an antibiotic-free diet. When venesection was required, bleeding from the retro-orbital veins or exsanguination by cardiac puncture from the anaesthetised mouse was performed (Wright et al., 1974).

Experiments and results

1. Tolerance of Borrelia to cyclophosphamide. Borrelicidal level of cyclophosphamide (Wyeth Brothers, Maidenhead, Bucks.) was determined by giving intraperitoneally 1,000 mg/kg and 150 mg/kg respectively to two groups each of 12 mice. The mice were then exsanguinated by cardiac puncture 75 min later and the sera from 4 mice were pooled. Thus the active cytotoxic alkylating metabolite of cyclophosphamide was obtained (Sladek, 1973). 0.2 ml aliquots of the fresh pooled serum was added to a monolayer of human embryonic lung cells (Flow Laboratories, Scotland). The monolayers were suspended in Eagles minimum essential medium (Flow Laboratories) in 150×15 millimetre Pyrex glass tubing (Corning Glass Works, N.Y. 148390) and sealed by a siliconed bung. A cytopathic effect was sought after 48-h incubation at 37° C. Cytopathic effects were found after 24 h with a sixfold dilution of sera from the high dose cyclophosphamide regime, but not after the low dose regime (Table 1). Equal volumes of fresh serum were also added to spirochaetal suspensions of Wellcome strain of Borrelia and incubated for 2 h. The serum had been taken from the cyclophosphamide-treated mice and from the appropriate controls (Table 1). The Borrelia were then counted and the serum-Borrelia suspension mixture was then inoculated into four mice. The *Borrelia* in the peripheral mouse blood were then counted at 48 h. The "toxic" effect of the sera from mice who had received cyclophosphamide or *Borrelia* paralleled the cytopathic effect in cells

Table 1. The effect of cyclophosphamide (Cy) on cells and Borrelia duttoni

	Cytopathic effect on human embryonic lung	Borrelia count/ml	
		2-h incubation in vitro	48 h after infection in vivo**
Borrelial suspension (BS) alone	_	3×10^{8}	4×10^{9}
Cy 1,000 mg/kg sera*	+	2×10^7	2×10^7
Cy 1,000 mg/kg sera diluted 1/3	+	2×10^7	6×10^8
Cy 1,000 mg/kg sera* diluted 1/6	+	4×10^8	9×10^8
Cy 1,000 mg/kg sera* diluted 1/10	_	5×10^8	5×10^9
Cy 1,000 mg/kg sera* diluted 1/100	-	6×10^8	6×10^9
Cy 150 mg/kg sera*	=	3×10^8	8×10^9
Normal mouse sera*	_	4×10^8	6×10^9
Krebs saline plus BS	=	3×10^8	7×10^9
Human embryonic lung cells	-		

^{*} All sera taken 75 min after giving cyclophosphamide.

(Table 1). Incubation of a solution containing 40 mg of cyclophosphamide in Krebs saline with human embryonic lung cells or *Borrelia* produced no cytopathic or borrelicidal effect. The Wellcome mouse strain was uniformly fatal to mice and therefore not used in the following experiments.

2. The effect on murine borrelial infection of antibody suppression by cyclophosphamide. Twelve mice were given cyclophosphamide 150 mg/kg intraperitoneally then infected intraperitoneally with a 0.5 ml of a Borrelia duttoni (STIB 503 type 1) suspension of 10^7 organisms per ml. A further 4 mice were given normal saline, the diluent for cyclophosphamide, and then infected with Borrelia duttoni as above. Eight further mice were just infected without a preliminary injection. The first parasitaemia (P) was detected 4 days later. The mean number of organisms found in the parasitaemia was 5.4×10^9 borrelia per ml in the animals given cyclophosphamide and 2×10^9 /ml in the non-immunosuppressed animals. Eight out of 12 immunosuppressed animals succumbed to the infection, 1 out of 4 mice given Krebs saline and then infected and 4 out of 8 of

^{**} Mean of blood samples from 4 mice.

Table 2. Serological differences between the first (P) and second (F₁) borrelial parasitaemia

Source of Borrelia	Test sera				
	4 days post infection	Anti-P	11 days post infection (anti $P + F_1$)	Anti-F ₁	
P control	+	+	+	_	
P cyclophosphamide	+	+	+	_	
F ₁ control	-	-	+	+	
F_1 cyclophosphamide	-	_	+	+	

⁴ and 11 day post-infection sera tested for lysins and agglutinins. Anti-P and anti-F₁ tested for agglutinins only.

the infected controls. The second parasitaemia (F_1) was detected 4 days later in the control animals and 5 days later in the immunosuppressed mice. In this second parasitaemia (F_1) the maximal number of spirochaetes seen was 0.5– 1×10^6 organisms per ml. None of these animals died. No specific antibodies were detected either against the P or F_1 parasites in sera taken between parasitaemias or 11 days after the initial infection in the immunologically compromised mice. In contrast, antibody in the control mice appeared against the P spirochaetes on the 4th day and against the F_1 parasites on the 11th day following infection. Prior to infection all mice had a negative serology. Antibody titres were: lytic antibody $\frac{1}{100} = \frac{1}{400}$, agglutinating antibody $\frac{1}{40} = \frac{1}{320}$ and immunofluorescent antibody $\frac{1}{100}$ to $>\frac{1}{500}$. P spirochaetes in both the cyclophosphamidetreated and control animals were found to be serologically identical as was the F_1 variant found in the second parasitaemia (Table 2).

The experiment was repeated using an identical protocol with $B.\ duttoni$ (STIB type 2) strain for producing the experimental infection, i.e. the spirochaete that had been mouse to tick and then mouse passaged. The infecting dose, given intraperitoneally, was 0.2 ml. The first parasitaemia developed on the fifth day. 6 out of the 12 immunologically suppressed mice died and 7 out of the 12 controls. The P variant was found in both groups of mice. No F_1 parasitaemia was detected although by day 11 antibodies against the F_1 strain had appeared in the normal mice. The P parasitaemia again resolved and by the 11th day no antibodies were detected in the mice given cyclophosphamide.

Discussion

The mouse can recover from borrelial infection without measurable specific antibody being demonstrated. More surprisingly, antigenic variation may occur as seen in the initial mouse experiment. Although no borrelial antibody was detected in the cyclophosphamide treated mice, it is possible that antibo-

dies inducing antigenic variation could still develop. Cyclophosphamide may enhance this type of antibody production, perhaps by inhibiting suppressor thymus derived lymphocytes (Thomas et al., 1977). If production of antibody modulating antigen was thereby enhanced, this antibody would not only protect against infection, but would induce antigenic change in the immune compromised mice. The failure in mice to demonstrate delayed hypersensitivity to spirochaetes or to find haptenic cardiolipin antibodies in *T. pallidum* infected mice, which would presumably require T-lymphocyte help for production (Wright et al., 1974), does not necessarily exclude T-lymphocyte suppressant activity. The deletion of these cells may therefore allow increased production of the antibody which modulates the variation.

The possibility that there might be two antigenic variants of *Borrelia* inoculated simultaneously in the above experiments is not likely as the findings do not account for the "spontaneous" cure or explain why the F₁ variant could be passaged without P spirochaete antibody being produced. Alternatively it might be suggested that lytic antibody was produced in such small quantities that it was "complexed" with the circulating antigen, lysed the bacteria leaving no antibody to be detected (Schuhardt and Wilkerson, 1951). The inability to detect antibody in the immunosuppressed animals even after the second parasitaemia makes this suggestion unlikely.

The activated cyclophosphamide metabolites are unlikely to induce the antigenic change, as the same variants appeared whether or not the mice had received cyclophosphamide. Activated cyclophosphamide is of transient duration in the mouse (Berenbaum et al., 1973) and in the concentration of cyclophosphamide used in the animal experiments, showed no inhibition of in vivo growth of the mouse adapted strain of spirochaetes. Antigenic variation may be dependent on antibody, but still environmentally influenced. Paramecium undergo antigenic changes with differences in environmental temperature and salinity (Beale, 1974). The occurrence of cross agglutinating antibody in alternate relapses reported by Cunningham et al. (1934) does not rule out the possibility that antigenic variation may be an inherent, perhaps a genetically controlled characteristic.

Acknowledgments. I thank Mr. Derek Frost of the Department of Microbiology, Charing Cross Hospital Medical School, for his excellent technical assistance.

¹ Balteanu I., Russ M., Voiculescu M.: The spirochaetolytic property of relapsing fever convalescent serum. Arch. roum. Path. exp. 15, 310–312 (1948).

² Beale G. H.: Genetics of antigenic variation in Paramecium. In: Ciba Foundation symposium 25 (NS): Parasites in the unimmunized host – mechanism of survival, p. 29–30. Elsevier/Excerpta Medica/North Holland, Amsterdam 1974.

³ Berenbaum M. C., Cope W. A., Double J. A.: The effects of microsomal enzyme inhibition on the immunosuppressive and toxic effects of cyclophosphamide. Clin. exp. Immunol. *14*, 257–270 (1973).

- 4 Calabi O.: Presence of plasma inhibitors during the crisis phenomenon in experimental relapsing fever (*Borrelia novyi*). J. exp. Med. 110, 811–825 (1959).
- 5 Cunningham M. J., Theodore J. H., Fraser A. G. L.: Further observations on Indian relapsing fever, Part 1: Types of spirochaetes found in experimental infection. Ind. J. med. Res. 22, 105–155 (1934).
- 6 Schuhardt V. T., Wilkerson M.: Relapse phenonemon in rats infected with single spirochaetes (*Borrelia recurrentis var turicatae*). J. Bact. 62, 215–219 (1951).
- 7 Sladek N. E.: Bio-assay and relative cytotoxic potency of cyclophosphamide metabolites generated in in vitro and in vivo. Cancer Res. 33, 1150–1158 (1973).
- 8 Southee T.: Enumeration of *Treponema pallidum* using an electronic counter. Med. lab. Technol. 29, 329–331 (1972).
- 9 Thomas W. R., Watkins M., Asherson G. L.: Suppressor cells in contact sensitivity. Effect on T-cell helping antibody production to picryl chloride. Cell. Immunol. *34*, 319–325 (1977).
- 10 Vickerman K.: Antigenic variation in trypanosomiasis. In: Ciba Foundation symposium 25 (NS): Parasites in the unimmunized host mechanism of survival, p. 53–80. Elsevier/Excerpta Medica/North Holland, Amsterdam 1974.
- 11 Wright D. J. M., Gaugas J. M., Rees R. J. W.: *T. pallidum* in mice with suppressed-cell mediated immunity. Guy's Hosp. Rep. 123, 385–390 (1974).
- 12 Wright D. J. M., Ginger C. E.: Syphilitic immunofluorescence in experimental relapsing fever. Brit. J. vener. Dis. 49, 239–241 (1973).