

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 36 (1979)
Heft: 4

Artikel: Identification by the blood incubation infectivity test of "Trypanosoma brucei" subspecies isolated from game animals in the Luangwa Valley, Zambia
Autor: Awan, M.A.Q.
DOI: <https://doi.org/10.5169/seals-312537>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Central Veterinary Research Station, Mazabuka, Zambia

Identification by the blood incubation infectivity test of *Trypanosoma brucei* subspecies isolated from game animals in the Luangwa Valley, Zambia

M. A. Q. AWAN

Summary

A total of 7 stocks of *Trypanosoma brucei* subspecies, isolated from naturally infected game animals in the Luangwa Valley, Eastern Province, Zambia were examined using a modified version of the Blood Incubation Infectivity Test (BIIT). One stock giving consistent BIIT responses typical of *T. b. rhodesiense*, was obtained from warthog (*Phacochoerus aethiopicus*). Four other stocks, 2 from hyaena (*Crocuta crocuta*), 1 from a waterbuck (*Kobus ellipsiprymnus*) and 1 from a lion (*Panthera leo*) responded like *T. b. brucei*. One stock from a waterbuck and 1 from a giraffe (*Giraffa camelopardalis*) failed to infect mice after incubation in human serum for 30 min at 37° C when first tested, but after 5 or 6 further serial passages in mice and even with serum incubation time increased to 5 h, they retained infectivity.

Key words: *Trypanosoma brucei* subspecies; blood incubation infectivity test; game animals.

Introduction

The identification of *T. b. rhodesiense* in a bushbuck (*Tragelaphus scriptus*) by Heisch et al. (1958) and later in cattle by Onyango et al. (1966) provided the first evidence of the significance of game and domestic animals as reservoirs of trypanosomes pathogenic for man.

However, the lack of a suitable laboratory test with which to differentiate human-infective trypanosomes among the morphologically identical members of the *T. brucei* complex, when these were isolated from a non-human host, has precluded both quantitative and qualitative assessment of the game reservoirs

Correspondence: M. A. Q. Awan, Chemonics International, American Embassy, Bamako, Republic of Mali

Table 1. BIIT responses of 7 stocks of *Trypanosoma brucei* subspecies, isolated from naturally infected game animals in the Luangwa Valley, Zambia, after incubation in human serum in vitro at 37°C for 30 min or 5 h

Stock number	Date of isolation	Host	BIIT														
			30 min incubation			5 h incubation			Sample B (Control)			Sample C (Test)			Sample D (Control)		
			Passages before testing	Mouse number	Mean PP	Mouse number	Mean PP	Mouse number	Mean PP	Mouse number	Mean PP	Mouse number	Mean PP	Mouse number	Mean PP		
J 10	4. 11. 1973	Hyaena (<i>Crocuta crocuta</i>)	3	-	-	-	+	+	+	1.0	-	-	-	+	+	+	1.66
J 11	8. 11. 1973	Hyaena (<i>Crocuta crocuta</i>)	9	-	-	-	+	+	+	2.33	-	-	-	+	+	+	1.0
H 6	21. 8. 1974	Waterbuck (<i>Kobus ellipsiprymnus</i>)	6	-	-	-	+	+	+	4.66	-	-	-	+	+	+	5.6
H 3	13. 8. 1974	Lion (<i>Panthera leo</i>)	1	-	-	-	+	+	+	1.33	-	-	-	+	+	+	1.0
H 18	25. 11. 1974	Warthog (<i>Phacochoerus aethiopicus</i>)	1	+	+	+	+	+	+	6.0	+	+	+	+	+	+	1.0
H 1	21. 7. 1974	Waterbuck (<i>Kobus ellipsiprymnus</i>)	5	-	-	-	+	+	+	1.0	not tested						
H 1	21. 7. 1974	Waterbuck (<i>Kobus ellipsiprymnus</i>)	11	-	not tested						+	+	+	+	+	+	1.0
H 15	6. 10. 1974	Giraffe (<i>Giraffa camelopardalis</i>)	1	-	-	-	+	+	+	1.33	not tested						
H 15	6. 10. 1974	Giraffe (<i>Giraffa camelopardalis</i>)	6	not tested							+	+	+	+	+	+	2.0

- = indicates mouse did not become parasitaemic within 40 days

+ = indicates mouse developed parasitaemia within 40 days

PP = prepatent period in days

of *T. b. rhodesiense* in East and Central Africa. The discovery and development of the blood incubation infectivity test (BIIT) by Rickman and Robson (1970a, b) provided such a test which has led to the study of several *T. brucei* sub-group stocks isolated from game, domestic stock, human patients and tsetse flies (Awan 1971; Dillmann and Awan, 1972; Robson et al. 1972; Awan and Dillmann, 1973; Robson and Rickman, 1973; Targett and Wilson, 1973; Geigy et al., 1975).

In this paper we report further investigations of the game animal reservoirs of *T. b. rhodesiense* and *T. b. brucei* and describe the BIIT responses given by 7 stocks isolated from naturally infected game animals in the Luangwa Valley, Zambia.

Materials and methods

Trypanosome stocks. All the stocks originated from an area on the eastern bank of the Luangwa River, between Milyoti Gate and Nyamaluma Pontoon (i. e. between latitudes 13–13°30' S and between longitudes 31° 30'–32° E) at an altitude of 600–800 metres; they were isolated in mice and preserved in liquid nitrogen using 7.5% glycerine as cryoprotectant.

Blood Incubation Infectivity Test (BIIT). A stabilate of the stock to be tested was brought to ambient temperature and inoculated intraperitoneally to 3 mice. Cardiac blood from a single donor mouse provided all the samples for any one test.

Identification of the test stock as *T. brucei* subspecies was confirmed before testing by morphological examination of a Giemsa stained, thin tail-blood film. Each stock was tested by incubating 2 samples (A and B) for ½ h and 2 samples (C and D) for 5 h at 37° C (Awan and Dillmann, 1973). Only the “test” samples were incubated in the human serum, while the “control” samples were incubated in the absence of human serum. Samples were examined by wet-film microscopy before and after incubation at 37° C, and three mice were then inoculated with aliquots of each sample. Post-inoculation microscopy of recipient animals, by wet-film examination of tail blood samples, was continued up to 40 days if aparasitaemic.

Results

Wet-film examination of samples. No difference could be detected in trypanosome motility in samples examined before and after incubation at 37° C, in the absence of human serum. By contrast, in most test samples of BIIT positive stocks, after incubation in serum, trypanosome number and motility were both markedly reduced. This adverse effect of human serum in vitro was reflected in a significant prolongation of the prepatent periods in the recipient animals (Table 1).

BIIT responses (Table 1). Of the 7 stocks tested, 4 (i. e. J10, J11 from hyaena, H6 and H3 from waterbuck and lion, respectively) were uninfected to mice after serum incubation of 30 min or 5 h. One stock, H18 (warthog) retained infectivity in each of the 2 tests, i. e. with incubation of 30 min or 5 h, although one of the 3 test mice inoculated with the sample of stock H18 incubated for 5 h remained negative up to day 38.

The remaining 2 stocks, H1 (waterbuck) and H15 (giraffe), were uninfec-

tive after 30 min incubation, but both retained infectivity when tested after 6 and 5 further passages (respectively) in mice, although the incubation time was increased to 5 h.

Discussion

Because of its consistent infectivity it is likely that the stock from warthog was *T. b. rhodesiense*. This is the first time that this parasite has been identified in these animals. In a preliminary study also the warthog stock retained its infectivity (Dillmann and Townsend, 1976). Similarly, the 4 stocks consistently losing infectivity were almost certainly *T. b. brucei*. The loss of infectivity also seems to confirm the earlier findings of Awan (1971), that incubating *T. b. brucei* trypanosomes in human serum in vitro at 37° C for 30 min is sufficient to inhibit the normal infectivity of these organisms for small laboratory rodents.

The inconsistency in the results of stocks H1 and H15 seems not to be due to the duration of incubation of the samples in human serum. It may be due to the possible presence of mixed populations of *T. b. rhodesiense* and *T. b. brucei* in newly isolated uncloned stocks from naturally infected game animals (Geigy et al., 1975). During the course of passage in mice the proportion of the two populations probably fluctuates and the resultant BIIT responses are dependent on the dominating population type in a donor mouse.

In our experiments there appears to be some in vitro activity of human serum on *T. b. rhodesiense* stocks, as evidenced by prolongation of the subsequent prepatent periods.

Acknowledgments. I am grateful to Miss R. Dryoff for technical assistance, Mr. L. R. Rickman for his constructive criticism and the Director of Veterinary and Tsetse Control Services, Zambia, for permission to publish this paper.

- 1 Awan M. A. Q.: The use of human plasma in the blood incubation infectivity test to differentiate *Trypanosoma brucei* and *T. rhodesiense*. Trop. anim. Hlth Prod. 3, 83–186 (1971).
- 2 Awan M. A. Q., Dillmann J. S. S.: *Trypanosoma brucei* infection in a lion (*Panthera leo*) in Zambia. Trop. anim. Hlth Prod. 5, 75–78 (1973).
- 3 Dillmann J. S. S., Awan M. A. Q.: The isolation of *Trypanosoma brucei* from *Hippopotamus amphibius* in the Luangwa Valley, Zambia. Trop. anim. Hlth Prod. 4, 135–137 (1972).
- 4 Dillmann J. S. S., Townsend A. J.: A trypanosomiasis survey on various species of wild animals in the Luangwa Valley, Zambia. Report of Veterinary Wildlife Research Section, Department of Veterinary and Tsetse Control Services 1, 173 (1976).
- 5 Geigy R., Jenni L., Kauffmann M., Onyango R. J., Weiss N.: Identification of *T. brucei* subgroup strains isolated from game. Acta trop. (Basel) 32, 190–205 (1975).
- 6 Heisch R. B., McMahon J. P., Manson-Bahr P. E. C.: The isolation of *Trypanosoma rhodesiense* from a bushbuck. Brit. med. J. II, 1203–1204 (1958).
- 7 Onyango R. J., van Hove K., de Raadt P.: The epidemiology of *T. rhodesiense* sleeping sickness in Alego location, Central Nyanza, Kenya. I. Evidence that cattle may act as reservoir hosts of trypanosomes infective to man. Trans. roy. Soc. trop. Med. Hyg. 60, 175–182 (1966).
- 8 Rickman L. R., Robson J.: The blood incubation infectivity test: a simple test which may serve to distinguish *Trypanosoma brucei* and *T. rhodesiense*. Bull. WHO 42, 650–651 (1970a).

- 9 Rickman L. R., Robson J.: The testing of proven *Trypanosoma brucei* and *T. rhodesiense* strains by the blood incubation infectivity test. Bull. WHO 42, 911–916 (1970b).
- 10 Robson J., Rickman L. R.: Blood incubation infectivity test results for *Trypanosoma brucei* subgroup isolates in the Labwe Valley, South Nyanza, Kenya. Trop. anim. Hlth Prod. 5, 187–191 (1973).
- 11 Robson J., Rickman L. R., Allsopp R., Scott D.: The composition of *Trypanosoma brucei* subgroup in non-human reservoirs in the Lambwe Valley, Kenya, with particular reference to the distribution of *T. rhodesiense*. Bull. WHO 46, 765–770 (1972).
- 12 Targett G. A. T., Wilson V. C. L. C.: The blood incubation infectivity test as a means of distinguishing between *Trypanosoma brucei* and *T. brucei rhodesiense*. Intern. J. Parasit. 3, 5–11 (1973).

