

Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	37 (1980)
Heft:	1
Artikel:	Colonization of "Trypanosoma brucei gambiense" within transplanted Ehrlich's tumors of "Microtus montanus"
Autor:	Seed, J.R. / Seed, T.M.
DOI:	https://doi.org/10.5169/seals-312635

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Laboratory of Microbiology and Parasitology, Department of Biology,
Texas A & M University, College Station, TX 77843, USA
Division of Biological and Medical Research, Argonne National Laboratory,
9700 South Cass Avenue, Argonne, Illinois 60439, USA

Colonization of *Trypanosoma brucei gambiense* within transplanted Ehrlich's tumors of *Microtus montanus*

J. R. SEED, TH. M. SEED

Summary

Trypanosoma brucei gambiense has been observed growing in extravascular sites throughout a solid Ehrlich ascites tumor (EAT) in *Microtus montanus*. These trypanosomes appear in clusters in the tumor. The possible importance of this observation to the host-parasite relationship is discussed.

Key words: colonies; *Trypanosoma brucei gambiense*; *Microtus montanus*; extravascular sites.

There are recent reports which demonstrate that *Trypanosoma congolense* and *T. brucei* are located in extravascular sites in chronically infected animals (Luckins and Gray, 1978; Ssenyonga and Adam, 1975). In this paper, we present evidence for the presence of extravascular colonization by *T. b. gambiense* within transplanted Ehrlich's tumor of chronically infected *Microtus montanus*. In many ways, these colonies *in vivo* resembled the uneven distribution (or clouds) of trypanosomes growing over and between feeder-layer cells *in vitro* (Hirumi et al., 1977).

Materials and methods

Microtus montanus were obtained from an outbred colony maintained at Texas A & M University in College Station, Texas and maintained as previously described (Seed and Negus, 1970). All animals used were sexually mature and maintained on a 12 hour light-12 hour dark photoperiod regime. The maintenance and harvesting of both the trypanosome and the Ehrlich-ascites tumor cell (EATC) line have been described (Ackerman and Seed, 1976). Both cell types were washed in glucose (1%) phosphate buffered saline (pH 7.4) and then counted with the aid of a hemacytometer.

Correspondence: Prof. John R. Seed, Department of Biology, Texas A & M University, College Station, Texas 77843, USA

The *Microtus* were infected intraperitoneally with 5×10^5 organisms of the Wellcome TS strain (TTrT-1) of *T.b.* The pathogenicity of this strain in *Microtus* has been described (Seed and Negus, 1970). Twelve days later, the infected animals were inoculated with 2×10^7 EATC subcutaneously; and sacrificed 8 days later.

The solid Ehrlich ascites tumors (EAT) were then removed and fixed in 1.25% (v/v) glutaraldehyde plus 4% sucrose in 0.05 M phosphate buffer. The tissues were later post-fixed in 1% buffered osmium tetroxide, dehydrated, and embedded in Epon according to standard procedures. Thick sections were prepared for light microscopy and examined after staining with 1% toluidine blue for 15 min at 45°C. Thin sections were cut and doubly stained with uranyl acetate and lead citrate. These preparations were examined with a Siemens 101 TEM.

Results and discussion

The light micrograph in Fig. 1 shows an extravascular concentration of trypanosomes within a solid Ehrlich tumor. The trypanosomes appeared to be distributed in clusters throughout the section. These clusters varied in both their size and location within the tumor, and were often a considerable distance from the vascular bed. In addition, there was no obvious correlation between the location of the clusters and the type of host or tumor cell present.

Similar observations were made using the transmission electron microscope. The electron micrograph also shows that the trypanosomes within the cluster appear morphologically intact with respect to their surface coat, cell membrane, and other organelles (Fig. 2). The trypanosomes within the tumors do not resemble published reports of either trypsinized or lysed cells, and dividing forms were observed within the tumors.

It is apparent that *T. b. gambiense* grow predominantly in extravascular sites throughout a solid EAT. We suggest that this may be a good model for further studies on the biology of trypanosomes in these sites.

Acknowledgments. This work was supported in part by the U.S. Department of Energy and by a grant from the Trypanosomiasis component of the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases. The kind assistance of the Argonne National Laboratory is also acknowledged. A major portion of this work was completed while the senior author was a visiting scientist there.

- Ackerman S. B., Seed J. R.: Effect of *Trypanosoma brucei gambiense* infections in *Microtus montanus* on susceptibility to Ehrlich's tumor. *Infect. Immun.* 13, 388-390 (1976).
- Hirumi H., Doyle J. T., Hirumi U.: African trypanosomes: cultivation of animal-infective *Trypanosoma brucei* in vitro. *Science* 196, 992-994 (1977).
- Luckins A. G., Gray A. R.: An extravascular site of development of *Trypanosoma congolense*. *Nature (Lond.)* 272, 613-614 (1978).
- Ssenyonga G. S. Z., Adam K. N. G.: The number and morphology of trypanosomes in the blood and lymph of rats infected with *Trypanosoma brucei* and *T. congolense*. *Parasitology* 70, 255-261 (1975).

Fig. 1. A light micrograph ($\sim 320 \times$) of a solid EAT from an infected *Microtus*. The long arrow points to one of many trypanosomes observed within the tissue section. The short arrow points to one of several small capillaries within the section.

Fig. 2. A transmission electron micrograph ($\sim 19000 \times$) of trypanosomes within a solid EAT.



