

**Zeitschrift:** Acta Tropica  
**Herausgeber:** Schweizerisches Tropeninstitut (Basel)  
**Band:** 37 (1980)  
**Heft:** 3

**Artikel:** Chemotherapy of experimental echinococcosis  
**Autor:** Eckert, J. / Burkhardt, B.  
**DOI:** <https://doi.org/10.5169/seals-312666>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 16.09.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Department of Parasitology, University of Zurich, Switzerland

## Chemotherapy of experimental echinococcosis<sup>1</sup>

J. ECKERT, B. BURKHARDT

### Introduction

Since 1974 it has been demonstrated that mebendazole, fenbendazole, flubendazole and some other benzimidazole derivatives exhibit certain anthelmintic effects against metacestodes of *Echinococcus granulosus* and *E. multilocularis* in intermediate host animals (Lit. see Barandun, 1978; Eckert et al., 1978; Burkhardt, in preparation).

Previous experiments (Eckert et al., 1978) with *Meriones unguiculatus* indicated that cysts of *E. granulosus* can be severely damaged or killed by long-term treatment of 80 days duration with 500 ppm mebendazole or fenbendazole applied in the food. Metacestodes of *E. multilocularis* were significantly inhibited in proliferation by long-term oral mebendazole treatments (500 ppm) over 60–200 days. In most cases, however, the parasites were capable to resume growth after termination of drug application as proved by transplantation experiments (Barandun, 1978; Eckert et al., 1978).

In recent studies the influence of chemotherapy on metastases formation of larval *E. multilocularis* in *Meriones* was examined.

### Materials and methods

Ninety *Meriones unguiculatus*, 6 months old (75 males, 15 females) were used. Of these, 60 animals were infected by implantation of 0.1 g tissue of larval *E. multilocularis* (strain B) per animal into the subcutis of the neck region. Three experimental groups were formed: *Group I*: 30 infected animals, treated from day 7 postinfection (p.i.) for 300 days with 500 ppm mebendazole<sup>2</sup> applied in medicated food pellets. On the basis of food consumption the daily drug dosis was estimated to be about 30–50 mg/kg body weight. *Group II*: 30 infected animals served as untreated control. *Group III*: 30 noninfected and untreated animals.

<sup>1</sup> Supported by the Swiss National Science Foundation (Grant No. 3.959.-0.78)

<sup>2</sup> 5(6)-benzoyl-2-benzimidazole carbamate (Janssen, Beerse, Belgium)

Survival rates of *Meriones* with subcutaneous *E. multilocularis* infection. Group I: treated from day 7-307 postinfection (300 days) with 500 ppm mebendazole, Group II: infected, untreated, Group III: uninfected, untreated.




Fig. 1

## Results and discussion

**Survival rates** (Fig. 1): Only 4 (13%) of the 30 infected but untreated animals of group II survived until the end of the experiment as compared with 17 (57%) of the infected and treated group I. The survival rate in the noninfected and untreated group III was 60% and thus not significantly different from the treated group I.

**Metastases formation:** Within 2–10 months p.i. most of the 30 untreated control animals of group II developed large parasite tissue masses in the subcutis of the cervical region, in the regional and thoracic lymph nodes and partly in the lungs (Fig. 2). The average group weight of metacestodes from all locations was 7.8 g with an individual maximum of 19.1 g (Fig. 3). Metastases in thoracic lymph nodes, in the lung tissue or the pleural cavity, respectively, were present in 21 animals or 70% of the untreated control group II.

In 28 *Meriones* of the treated group I only small remnants of parasite tissue with an average weight of 0.03 g were detectable at the implantation site (Figs. 3 and 4). All organs were free of macroscopically visible metastases.

**Transplantation experiments:** Ten days after termination of treatment parasite tissue isolated from survived untreated and treated animals, respectively, was transplanted into the peritoneal cavity of helminth-free *Meriones*.





Fig. 2. *Meriones* (dorsal view) of untreated group II 134 days postinfection. Large parasite masses at implantation site (1) and in lymphnodes (2).

Weight\* of *E. multilocularis* metacestode tissue from *Meriones* infected subcutaneously.



\*total metacestode weight from all locations. n = number of animals start / end of experiment.  
d = % difference of mean metacestode weights between treated animals and untreated controls.  
p = probability

Fig. 3




Fig. 4. *Meriones* (dorsal view) of group I 230 days postinfection and after 223 days of mebendazole treatment. Arrow 1: remnants of parasite tissue at implantation site.

After 70 days living parasite tissue was present in 8 animals which had been infected with transplants from untreated controls and in all 5 animals which had received transplants from *Meriones* treated for 300 days.

These results indicate that long-term treatment with high oral doses of mebendazole can prevent proliferation and metastases formation of larval *E. multilocularis* in *Meriones* under drug treatment, but it has no complete parasitocidal effect.

Barandun G.: Untersuchungen zur Chemotherapie der alveolären Echinokokkose und der *Mesocystoides corti*-Infektion bei Labortieren. Vet. Diss., Zürich 1978.

Burkhardt B.: Beiträge zur experimentellen Chemotherapie der larvalen Echinokokkose mit Untersuchungen zur biologischen Verfügbarkeit von Mebendazol bei Nagetieren. Vet. Diss., Zürich (in preparation).

Eckert J., Barandun G., Pohlenz J.: Chemotherapie der larvalen Echinokokkose bei Labortieren. Schweiz. med. Wschr. 108, 1104–1112 (1978).