

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 37 (1980)
Heft: (11): Santé de l'enfant d'âge scolaire en Côte d'Ivoire

Artikel: Summary
Autor: [s.n.]
DOI: <https://doi.org/10.5169/seals-312675>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.09.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Summary

The purpose of this study was to examine the health status of 430 school-age children living in four villages of the forest region of the Ivory Coast. Basic anthropometric and hematological data as well as vitamin status were determined, and the evolution of the nutritional status was examined in relation to parasitic infection and diet.

An antiparasitic treatment for intestinal helminths and schistosomiasis provoked, over a period of one year, a marked change in growth as well as in blood levels of several vitamins. In contrast, the hematological values remained surprisingly constant compared to those in untreated children.

Various environmental aspects such as family, social and ethnic background have been presented as well as some epidemiological data on primary tuberculosis, toxoplasmosis, cytomegaly and hepatitis (HBs antigen). Daily food intake was determined through weighing of the ingredients, the composition of which was calculated using FAO tables. It was shown that caloric intake was only 75%, protein consumption 80% and lipid intake 30% of recommended levels of intake. The diet contained insufficient amounts of thiamin, riboflavin and niacin while vitamin C and carotenoid content was adequate.

A quantitative parasitological survey revealed a high prevalence of ascariasis and ancylostomiasis in all villages, schistosomiasis in two of them, onchocerciasis and trichuriasis in one village each. The intestinal worm load decreased with age. Performing serological tests for schistosomiasis and filariasis the difficulty arose to accord sensitivity and specificity in polyparasitized children. Malaria was holoendemic in all villages proven by high splenic and parasitic indices. An increase with age of *P. falciparum* was observed with a concomitant decrease of a mixed *P. falciparum* and *P. malariae* infection.

Moderate malnutrition among 30% of the children, as indicated by anthropometric measurements (weight-for-height, height-for-age and skin fold), appeared to be the consequence of the various helminthic infestations (schistosomiasis included). Malaria seemed to have an inhibitory effect on growth.

A hematological study revealed microcytic anemia among 30% of the children which correlated well with malarial infection but not with necatorosis and schistosomiasis. An association between anemia and a diet low in protein and iron cannot be excluded, but there are clearcut correlations between anemia and low vitamin A, B₂ and C blood levels. No interaction between the various hemoglobinopathies and malarial parasitic rates could be demonstrated.

The biochemical determination of vitamin status (A, B, B₁, B₂, B₆, B₁₂, folate, niacin) showed more or less pronounced deficiencies of vitamin C, riboflavin, vitamin A and pyridoxin. These vitamin deficiencies were, with the exception of pyridoxin, associated with corresponding clinical signs. Vitamin A deficiency despite a high carotenemia is not easy to understand.

Treatment of intestinal helminthiases and schistosomiasis led to a considerable increase in vitamin C blood levels while other vitamins remained unchanged. Malaria infection levels were correlated with most vitamin concentrations and it was also noticed that onchocerciasis was associated with low vitamin A levels. It appears that riboflavin deficiency is of dietary origin whereas vitamin A and C deficiencies are connected with parasitic diseases.

An epidemiological study of hepatitis, toxoplasmosis and cytomegaly showed that all three of them were highly prevalent. The exceptionally high frequency of HBs antigen occurred in the village with onchocerciasis. The extent of prevalence of tuberculosis primary infection is consistent with previous studies in the Ivory Coast but it appeared that the tuberculinic reaction was weaker in children with schistosomiasis.