

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 38 (1981)
Heft: 4

Artikel: Metabolic end products of helminths : their degradation and excretion by the host : biochemical approach to the diagnosis of ascariasis and other helminthiases
Autor: Soprunov, F.F. / Lurje, A.A. / Laynis, Y.Y.
DOI: <https://doi.org/10.5169/seals-312845>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Martsinovsky Institute for Medical Parasitology and Tropical Medicine, Moscow, USSR

Metabolic end products of helminths: their degradation and excretion by the host

Biochemical approach to the diagnosis of ascariasis and other helminthiases

F. F. SOPRUNOV, A. A. LURJE, YU. YA. LAYNIS, N. YA. SOPRUNOVA,
H. H. ALIEVA

Summary

Accumulation, degradation and excretion by the host of helminth metabolic end products, such as branched-chain volatile fatty acids are poorly studied although they play an appreciable role in host-parasite interactions and, being detectable in excreta, provide a reliable diagnostic test for at least some helminthiases. The amount and the structure of the end products of intestinal Nematoda metabolism depend on pH, rH, pCO₂, pO₂, and composition of the media. In the human intestine *Ascaris* excretes 40–60 mmol volatile fatty acids/100 g wet weight daily. α -methylbutyric, α -methylvaleric and isovaleric acids are slowly degraded. In concentrations of 10⁻³–10⁻⁴ M they exert an inhibitory effect on O₂ consumption, oxydative phosphorylation and Mg²⁺-ATP'ase activity in mitochondria. Given by mouth to laboratory animals or injected intraperitoneally, ¹⁴C-labeled branched-chain volatile fatty acids rapidly appear in saliva and urine and are detectable by chromatography and IR spectrography.

Key words: *Ascaris*; metabolism; volatile fatty acids; diagnostic; saliva.

Introduction

Metabolic pathways of helminths differ in many aspects from the pathways of their hosts (Bueding, 1966; von Brand, 1972; van den Bossche, 1972; Soprunov, 1978). Some end products of helminth metabolism are unusual for the host (Moyle and Baldwin, 1952; Saz and Gerson, 1962; Bueding, 1966; Greichus and Greichus, 1966), for example the branched-chain volatile fatty

Correspondence: Prof. F. F. Soprunov, Martsinovsky Institute for Medical Parasitology and Tropical Medicine, M. Pirogovskaya 20, Moscow 119435, USSR

Table 1. End products of helminths

Species	End products (acids)	References
Nematodes:		
<i>Ascaris lumbricoides</i> (<i>Ascaris suum</i>)	Formic (C ₁), acetic (C ₂), propionic (C ₃), butyric (C ₄), α -methyl-butyric, valeric (C ₅), α -methyl-caproic, lactic, carbonic (ethyl alcohol)	Flury, 1912; Schulte, 1917; Epps et al., 1950; Bueding and Yale, 1951; Moyle and Baldwin, 1952; Bueding, 1953; Ellison et al., 1960; Harpur and Waters, 1960; Ueno, 1960; Pushkarev, 1965; Greichus and Greichus, 1966; Harpur and Leigh-Brown, 1971; Laynis, 1971, 1975 Toryu, 1933 Yanagisawa and von Brand, 1965
<i>Parascaris equorum</i>	Propionic, valeric, lactic	
<i>Angiostrongylus cantonensis</i>	Lactic	Shestak, 1973
<i>Mecistocirrus digitatus</i>	Acetic, propionic, α -methyl-butyric, lactic	Bueding, 1949 Bueding and Oliver-Gonzales, 1950 Glockin and Fairbairn, 1952; Fairbairn, 1954 von Brand and Browman, 1963 Kmetec and Bueding, 1965; Bueding, 1966 Crowley and Warren, 1963; Warren and Poole, 1970 Prichard and Rothwell, 1972 Prichard and Rothwell, 1972; von Brand, 1952; Castro and Fairbairn, 1969 Vaastra, 1969
<i>Litomosoides carinii</i>	Acetic, lactic, acetyl-methyl-carbinol	
<i>Dracunculus insignis</i>	Lactic	
<i>Heterakis gallinae</i>	Acetic, propionic, lactic, pyruvic, long-chain fatty acids	
<i>Dirofilaria immitis</i>	Lactic	
<i>Trichuris vulpis</i>	Formic, acetic, propionic, butyric, valeric, α -methyl-valeric, succinic, lactic, carbonic	
<i>Ancylostoma caninum</i>	Acetic, propionic, iso-butyric, iso-caproic, carbonic	
<i>Trichostrongylus colubriformis</i> (larvae)	Acetic, propionic, butyric	
<i>Trichinella spiralis</i>	Formic, acetic, propionic, butyric, valeric, caproic (C ₆)	
<i>Dictyocaulus viviparus</i>	Acetic	

Table 1 (continued)

Trematodes:		
<i>Fasciola hepatica</i>	Acetic, propionic, butyric, iso-butyric, α -methyl-valeric, valeric, lactic	Mansour, 1959; Moss, 1970; Senutayte, 1970; Lahoud et al., 1971
<i>Fasciola gigantica</i>	Lactic	Goil, 1957, 1961
Cestodes:		
<i>Hymenolepis diminuta</i>	Lactic, succinic	Read, 1956; Fairbairn et al., 1961
<i>Moniezia expansa</i>	Lactic, succinic, long-chain fatty acids	von Brand, 1933; Smyth, 1947
<i>Echinococcus granulosus</i>	Acetic, propionic, valeric, succinic, carbonic, long-chain fatty acids, ethyl alcohol	Agosin, 1957

acids (VFA) excreted by many species (Table 1). These end products are found in the body of vertebrates only if these harbour helminths (Karnaukhov and Soprunova, 1976; Alieva et al., 1980).

Accumulation, degradation and excretion of end products of parasites by the host are poorly studied although the identification of VFA in excreta of the host could serve as a highly sensitive and reliable diagnostic test for at least some forms of helminthiasis.

Materials and methods

Amounts and structure of end products excreted by *Ascaris lumbricoides*, *A. suum*, *Ascaridia galli*, *Fasciola hepatica* and some other parasitic worms have been investigated in our laboratory during the last 15 years (Pushkarev, 1966; Laynis, 1971; Senutayte, 1971; Lazdyna et al., 1976). The fate of these end products in the body of the host, their possible damaging action on the tissues and their excretion rate were extensively studied (Soprunova, 1968, 1971; Soprunova and Andreeva, 1968; Lazdyna and Buykis, 1970 a/1970 b; Laynis, 1971, 1975; Soprunova and Lurje, 1972; Soprunova et al., 1973; Karnaukhov and Soprunova, 1976; Alieva and Lurje, 1979; Alieva and Soprunov, 1979; Alieva et al., 1980). This work was carried out in laboratory animals (rats, rabbits, pigs and chickens) and in infected patients in our clinic and in foci of helminthic diseases.

Compensative potentiometry was used to follow the excretion of acid end products by the worms maintained in Hedon-Fleig medium at 37° C; the isolation of VFA from biological material was effected by distillation with steam and specific extraction with ether, benzene and other organic solvents. Individual VFA were identified by paper chromatography of hydroxamate derivatives, gas chromatography and infra-red spectroscopy; isotopic labeling was used for following VFA transformation and excretion and polarography in closed cells with rotating platinum electrode for the registration of the inhibitory effect of VFA on O₂ consumption and oxidative phosphorylation in isolated mitochondria (medium for polarography: 0.02 M KH₂PO₄, 0.05 M tris-buffer-pH-7.5, 0.001 M MgCl₂, 0.15 M KCl, 0.15 M glucose + 0.5 mg hexokinase, 0.001 M ATP, 0.001 M EDTA; concentration of substrates – 0.025 M; inhibitor – FNa); the micro-Astrup method was employed for determining the shift of physico-chemical properties in saliva. For details see previous publications (Pushkarev, 1966; Soprunova, 1968; Laynis, 1971; Soprunova and Lurje, 1972; Alieva and Lurje, 1979; Alieva et al., 1980).

Results

End products of Ascaris

Carbohydrate degradation pathways of intestinal worms are to some extent labile. Amounts and chemical structure of excreted VFA closely depend on the properties of the medium (e. g. pH, pCO₂, pO₂, concentration of glucose and mineral salts). For the effect of the redox potential, see Soprunov (1978, p. 28). Table 2 shows the effect of oxygen on VFA production.

Under conditions close to those prevailing in the lumen of the intestine the daily excretion of VFA by *Ascaris* is of the order of magnitude of 20–60 mmol/100 g wet weight. The structure of these metabolic end products is presumably similar to the VFA extracted from the hemolymph and tissues of fresh collected worms (Table 3).

Table 2. Amount of VFA excreted by *Ascaris suum* in dependence of medium conditions

Medium conditions for <i>Ascaris</i>	Total amount mmol/(10 g · 24 h)	VFA (mol %)					
		C ₁	C ₂	C ₃	C ₄	C ₅	C ₆
Aerobic	1.89 ± 0.32	4.7	27.5	18.5	4.3	27.0	18.0
Anaerobic (CO ₂)	2.01 ± 0.37	4.6	20.2	13.8	3.4	35.6	22.4
Anaerobic (N ₂)	2.17 ± 0.47	8.0	21.8	16.1	4.1	30.0	20.0

Table 3. VFA in haemolymph and tissues of *A. suum*

VFA	VFA (mol %)	
	in tissues of <i>Ascaris</i>	in haemolymph of <i>Ascaris</i>
C ₂	9.8 ± 0.7	13.7 ± 1.7
C ₃	4.0 ± 0.1	5.0 ± 0.3
iso-C ₄	1.2 ± 0.2	1.1 ± 0.3
n-C ₄	2.2 ± 0.1	2.4 ± 0.1
α-Me-C ₄ / iso-C ₅ ..	41.7 ± 1.4	40.0 ± 2.3
n-C ₅	9.2 ± 1.1	5.2 ± 0.9
α-Me-C ₅	31.2 ± 2.1	32.4 ± 1.4
iso-C ₆ / n-C ₆	0.7 ± 0.2	0.2 ± 0.1

A reliable diagnostic test for ascariasis can be based on the identification of α-methylbutyric, α-methylvaleric and isovaleric acids in the excreta of the host.

Damaging action of VFA on the host

Table 4 shows the inhibitory effect of ascaris VFA on O₂ consumption and on oxidative phosphorylation in isolated mitochondria. Table 5 shows the influence of VFA on ATP-ase activity.

We have found that as much as 60–80% of the total amount of VFA, excreted by the worms, are retained and metabolized in the body of the host (piglets). Structure and damaging effect of presumable intermediary products are unknown.

Excretion of VFA by the host

Given by mouth to laboratory animals (rats) or injected intraperitoneally, VFA rapidly appeared in saliva and urine (Table 6). On pigs experimentally invaded with *A. suum* it was shown that VFA appeared in urine before the worms reached maturity (before ova were found in faeces). In two pigs harbouring at autopsy only 1 and 2 male worms the urine test was positive. In urine and

Table 4. Influence of VFA ($1 \cdot 10^{-3}$ M) on O_2 consumption and oxidative phosphorylation in isolated liver mitochondria of rats (%)

VFA	ΔO	ΔP	$\Delta (P:O)$
C ₂	-2.8	-23.3	-20.9
C ₃	-2.8	-6.0	-2.7
n-C ₄	-0.2	-18.7	-19.7
iso-C ₄	-8.1	-26.5	-24.3
n-C ₅	+2.2	-15.1	-18.0
iso-C ₅	+3.2	-22.6	-26.5
α -Me-C ₄	+5.0	+1.0	-3.5
n-C ₆	-3.6	-12.0	-11.6
iso-C ₆	-0.7	-28.0	-27.7
VFA from <i>A. suum</i> tissues	+0.5	-14.7	-11.4
VFA from <i>A. suum</i> haemolymph	-14.0	-25.9	-13.6

Table 5. Influence of VFA ($5 \cdot 10^{-3}$ M) on Mg²⁺-dependent ATP-ase activity in rat liver mitochondria

VFA	ATP-ase activity $\mu\text{g P}/(\text{mg of protein} \cdot 30 \text{ min})$	Stimulation (%)
Without VFA	2.55 \pm 0.07	0
C ₂	4.64 \pm 0.11	+82.4
C ₃	4.92 \pm 0.14	+93.4
n-C ₄	4.73 \pm 0.14	+85.6
iso-C ₄	3.65 \pm 0.13	+43.0
n-C ₅	4.31 \pm 0.16	+69.0
iso-C ₅	3.50 \pm 0.12	+37.2
α -Me-C ₄	3.32 \pm 0.16	+30.3
Me ₃ -C ₂	3.20 \pm 0.08	+25.0
n-C ₆	4.85 \pm 0.07	+90.0
iso-C ₆	4.34 \pm 0.10	+69.8
α -Me-C ₅	3.26 \pm 0.10	+27.7
VFA from <i>A. suum</i> tissues	3.60 \pm 0.14	+41.0
VFA from <i>A. suum</i> haemolymph	3.83 \pm 0.13	+50.0

saliva of patients with ascariasis VFA are readily detected by paper and gas chromatography after extraction with suitable organic solvents (Soprunkova, 1968).

In our experiments the quantity of VFA detected in urine, saliva and faeces did not exceed 10–20% of the injected amount or the amount presumably excreted by the worms in the intestine. Excretion through the lungs and the skin was not studied.

Table 6. Excretion of VFA in rat urine

VFA given p. o.	Amount of VFA detected in urine, mmol/(m · 24 h)	VFA (%)					
		C ₂	C ₃	C ₄	α-Me-C ₄	C ₅	α-Me-C ₅
α-Me-C ₄ (0.25 mg/g of animal weight)	0.24 ± 0.05	57.3 ± 2.2	22.8 ± 7.2	7.0 ± 2.3	12.9 ± 4.1	—	—
α-Me-C ₅ (0.25 mg/g)	0.20 ± 0.04	51.7 ± 2.8	23.4 ± 5.0	9.2 ± 2.1	—	traces	15.7 ± 4.1
Volatile fraction from coeloma fluid of <i>A. suum</i> (0.50 mg/g)	0.25 ± 0.09	47.0 ± 14.3	19.5 ± 2.6	7.0 ± 4.4	11.5 ± 3.7	traces	15.0 ± 3.9
Without VFA	0.15 ± 0.08	57.5 ± 6.2	32.5 ± 5.7	10.0 ± 2.3	—	traces	—

Table 7. Comparative data of coproovoscopy and biochemical tests

Colour test in urine*		Coproovoscopy	Test in urine
106	patients with acute and chronic diseases (hepatitis, gastritis and others)	106–	106–
14	patients with ascariasis	14+	14+
76	patients with other helminthiasis (opistorchosis, trichocephalosis, enterobiosis and others)	76+	76–
Test in saliva**		Coproovoscopy	Δ pH
309	healthy children	309–	1.04 \pm 0.02
69	children with ascariasis***	69+	1.37 \pm 0.02
32	children with mixed invasion (ascariasis + other helminths)	32+	1.44 \pm 0.02
32	children with other helminthiasis (children harbouring <i>Trichocephalus trichiuris</i> , <i>Strongyloides stercoralis</i> , <i>Enterobius vermicularis</i> , <i>Hymenolepis nana</i> , <i>Taeniarhynchus saginatus</i> and <i>Diphyllobothrium latum</i>)	32+	1.25 – 1.34

* Patients in the clinic of the institute

** School children in foci of helminthiasis

*** In cases of heavy ascariasis the mean value was 1.55 \pm 0.09.

Simple diagnostic tests for VFA detection

Excess of NaOH is added to collected samples of urine or saliva. After drying on a water bath, residues are sent to the laboratory for chromatographic determination of VFA. In the absence of facilities for chromatography, the following tests may be useful:

a) Colour test in urine

1. Put 10 ml urine into a tube with a glass stopper containing 5 g anhydrous MgSO₄. Shake up to dissolution.
2. Add 1 ml H₂SO₄ conc. and 12 ml C₆H₆. Shake vigorously by hand for 4–5 min.
3. After separation into two layers transfer 10 ml of the upper benzene extract into a tube with 10 ml 3 mM NaOH. (N. B. Be careful not to contaminate the pipette.) Shake vigorously by hand for 4–5 min.
4. Discharge the upper layer. Transfer the alkaline solution (10 ml) into a clean tube.
5. Titrate the re-extract with the indicator solution up to colour change: blue \rightarrow orange. The reaction is considered positive if ≤ 6 drops are needed for colour change.

To prepare the indicator-solution dissolve 0.1 g bromthymol blue in 20 ml of hot ethanol and dilute up to 100 ml with 0.1 N H₂SO₄. The colour changes when 9–10 drops (0.3 ml) of this indicator are added to 10 ml of 3 mM NaOH sol.

b) Test in saliva

Collected samples of saliva in capillary tubes (80 microl). Close tubes rapidly at both ends with mastic (BMA-102) and send to laboratory in refrigerated box. Diluted samples 1/10 with physiol. sol. and introduce in vessels of p-CO₂-microelectrodes (membrane-Silicon fume D-606; sensitivity 10⁻⁴–10⁻⁵ mM H⁺). Register drop of pH (Δ pH) after 2 minutes.

Evaluation:

	Δ pH	<1.08	reaction negative (no VFA in saliva)
1.08	< Δ pH	<1.16	reaction negative (doubtful)
	Δ pH	>1.16	reaction positive (VFA in saliva)

Biochemical methods are used for mass examination in USSR, Roumania and Poland. Some results and comparative data are given in Table 7. Biochemical methods have been used for routine mass examination. As a rule they give a higher percentage of positive findings than coproovoscopy (Karnaukhov and Soprunova, 1976, Alieva et al., 1980).

Discussion

Amounts and structure of specific metabolic end products of *Ascaris* and their dependance on the culture medium have been studied by many authors. In this respect our data seem to roughly resemble those published previously.

However, there is a great discrepancy between the results in Table 2 and the results given in a previous publication (Soprunov, 1978, p. 28). Even if one takes into consideration that CO₂ accounts for 10 to 15% of the total amount of organic acids excreted by *Ascaris* at a redox potential of –400 mV and pH 8.0, it is obvious that the excretion rate of VFA in buffered alkaline flowing media with a low oxidative potential is 10 to 20 times higher (45.5 micromol/h/g wet weight) than in media where an accumulation of VFA takes place at a neutral pH (Table 2). The properties of the medium and the conditions in our previous experiments are rather close to those usually existing in the lumen of the human intestine and consequently the excretion of VFA by *Ascaris* in vivo must be much higher than usually expected.

To our knowledge, no data have been previously published about an accumulation of specific metabolic end products of helminths in the tissues of the host and no research has been carried out on the host's degradation pathways and the excretion of these products. We have found that at least 20 to 40% of the

VFA produced by *Ascaris* are excreted unchanged by the host through urine and saliva. This phenomenon can be used for diagnostic purposes.

We have developed simple, sensitive and reliable diagnostic methods which are much easier to use than microscopic or immunological assays. They may be useful, where microscopy fails. Polish authors (Pirog and Pietron, 1974) found the biochemical method to be 3 times more effective than copro-ovoscopy. The colour test proved to be sensitive and reliable (Roman et al., 1970; Pirog and Pietron, 1974).

Since the metabolic end products of helminths vary from species to species, it seems reasonable to assume that an analysis of the composition of VFA in saliva and urine should allow the development of differential diagnostic tests for various forms of helminthic infections.

- Agosin M.: Studies on the metabolism of *Echinococcus granulosus*. II. Some observations on the carbohydrate metabolism of hydatid cyst scolices. *Exp. Parasit.* 6, 586–593 (1957).
- Alieva H. H., Lurje A. A.: Excretion of volatile fatty acids in the saliva of rats. *Med. Parazit. (Mosk.)* 48, Nr. 4, 36–39 (1979).
- Alieva H. H., Soprunov F. F.: Changes in the acid-base values of mixed human saliva in intestinal helminthic diseases. *Med. Parazit. (Mosk.)* 48, Nr. 6, 35–39 (1979).
- Alieva H. H., Lurje A. A., Soprunov F. F.: Volatile fatty acids and their excretion from white rats with saliva and other ways. *Parazitologiya* 14, 71–75 (1980).
- Van den Bossche H.: Comparative biochemistry of parasites. Academic Press, New York 1972.
- Von Brand T.: Untersuchungen über den Stoffbestand einiger Cestoden und den Stoffwechsel von *Moniezia expansa*. *Z. vergl. Physiol.* 38 B, 43–52 (1933).
- Von Brand T.: Chemical physiology of endoparasitic animals. Academic Press, New York 1952.
- Von Brand T.: Parasitenphysiologie. G. Fischer Verlag, Stuttgart 1972.
- Von Brand T., Browman J. B. R.: Studies on the aerobic and anaerobic metabolism of larval and adult *Taenia taeniaeformis*. *Exp. Parasit.* 11, 276–297 (1963).
- Bueding E.: Metabolism of parasitic helminths. *Physiol. Rev.* 29, 195–218 (1949).
- Bueding E.: Formation of tiglic and *n*-valeric acids by bacteria free *Ascaris lumbricoides*. *J. biol. Chem.* 202, 505–512 (1953).
- Bueding E.: Biochemistry of parasites. Academic Press, New York/London 1966.
- Bueding E., Oliver-Gonzales J.: Aerobic and anaerobic production of lactic acid by the filarial worm *Dracunculus insignis*. *Brit. J. Pharmacol.* 5, 62–64 (1950).
- Bueding E., Yale H.: Production of α -methylbutyric acid by bacteria free *Ascaris lumbricoides*. *J. biol. Chem.* 193, 411–432 (1951).
- Castro G. A., Fairbairn D.: Carbohydrates and lipids in *Trichinella spiralis* larvae and their utilization in vitro. *J. Parasit.* 55, 51–58 (1969).
- Crowley R., Warren L. G.: Production of volatile acids by *Ancylostoma caninum*. *J. Parasit.* 49, 52–53 (1963).
- Ellison T., Thompson W. A. B., Strong F. M.: Volatile fatty acids from axenic *Ascaris lumbricoides*. *Arch. Biochem.* 91, 247–254 (1960).
- Epps W., Weiner M., Bueding E.: Production of steam volatile acids by bacteria free *Ascaris lumbricoides*. *J. infect. Dis.* 87, 149–151 (1950).
- Fairbairn D.: The metabolism of *Heterakis gallinae*. II. Carbon dioxide fixation. *Exp. Parasit.* 3, 52–63 (1954).
- Fairbairn D., Wertheim G., Harpur R. P., Schiller E. L.: Biochemistry of normal and irradiated strains of *Hymenolepis diminuta*. *Exp. Parasit.* 11, 248–263 (1961).

- Flury F.: Zur Chemie und Toxikologie der Ascariden. Arch. exp. Path. Pharmacol. 73, 214–232 (1912).
- Glockin V. C., Fairbairn D.: The metabolism of *Heterakis gallinae*. I. Aerobic and anaerobic respiration: carbohydrate sparing action of carbon dioxide. J. cell. comp. Physiol. 39, 341–356 (1952).
- Goil M. M.: Carbohydrate metabolism in trematode parasites. Z. Parasitenk. 18, 36–39 (1957).
- Goil M. M.: Physiological studies on trematodes *Fasciola gigantica* carbohydrate metabolism. Parasitology 51, 335–337 (1961).
- Greichus A., Greichus Y. A.: Chemical composition of volatile fatty acids of male *Ascaris lumbricoides* before and after starvation. Exp. Parasit. 19, 85–90 (1966).
- Harpur R. P., Leigh-Browne G.: *Ascaris lumbricoides* succinate and tiglate in haemolymph. Exp. Parasit. 29, 208–214 (1971).
- Harpur R. P., Waters W. R.: Production of carbon dioxide and volatile acids by muscle from *Ascaris lumbricoides*. Canad. J. Biochem. 38, 1009–1020 (1960).
- Karnauchov V. K., Soprunova N. Ya.: The diagnostic value of biochemical urianalysis (determination of volatile fatty acids) in ascariidosis. Med. Parazit. (Mosk.) 45, 450–452 (1976).
- Kmetec E., Bueding E.: Production of succinate by the canine whipworm *Trichuris vulpis*. Biochem. Physiol. 15, 271–274 (1965).
- Lahoud H., Prichard R. K., McManus W. R., Schofield P. J.: The dissimilation of leucine, isoleucine and valine to volatile fatty acids by adult *Fasciola hepatica*. Comp. Biochem. Physiol. 38, 379–391 (1971).
- Laynis Yu. Ya.: Volatile fatty acids excreted by *Ascaris suum* and their influence on the oxidative phosphorylation in mitochondrion of vertebrate animals. Tr. vsesoyuz. Inst. Helmintologii 17, 91–98 (1971).
- Laynis Yu. Ya.: Changes in vertebrate hosts mitochondria induced by the volatile fatty acids excreted by ascarides. Thesis, Riga 1975 (in Russian).
- Lazdynya M. A., Buykis A. Kh.: Influence of the carbohydrate metabolism endproducts of ascarides on white blood picture in experimental animals. In: Problems of parasitology in the Baltic Republics, p. 95–98, Riga 1970 a (in Russian).
- Lazdynya M. A., Buykis A. Kh.: Clinical and morphological changes in laboratory animals with the allergy induced by some components of or the whole volatile fraction of carbohydrate metabolism endproducts of ascarides. In: Problems of parasitology in the Baltic Republics, p. 156–158, Riga 1970 b (in Russian).
- Lazdynya M. A., Zarinya R. K., Soprunova N. Ya., Feodorova E. Ya.: The composition of volatile fatty acids of *Ascaridia galli* and their effect on *Eimeria tenella*. Med. Parazit. (Mosk.) 45, 189–192 (1976).
- Mansour T. E.: Studies on carbohydrate metabolism of liver fluke *Fasciola*. Biochim. biophys. Acta (Amst.) 34, 456–464 (1959).
- Moss G. D.: Endproducts of carbohydrates in *Fasciola hepatica*. Parasitology 60, 1–19 (1970).
- Moyle V., Baldwin E.: Volatile fatty acids of *Ascaris lumbricoides* from the pig. Biochem. J. 51, 504–509 (1952).
- Pirog Z., Pietron W.: Biochemiczna diagnostyka askarydozy i jej swoistosc. Wiadomości Parazytologiczne 20, 849–853 (1974).
- Prichard R., Rothwell T. L. W.: Volatile fatty acid production by parasitic fourth-stage *Trichostrongylus colubriformis*. J. Parasit. 58, 1161–1167 (1972).
- Pushkarev I. A.: Volatile fatty acids produced by ascarides *Ascaris suum* by cultivation under aerobic and anaerobic conditions. Med. Parazit. (Mosk.) 34, 551–556 (1965).
- Pushkarev I. A. Volatile fatty acids of *Ascaris suum*. In: Parasites of animals and the control of them, p. 75, Riga 1966 (in Russian).
- Read C. P.: Carbohydrate metabolism of *Hymenolepis diminuta*. Exp. Parasit. 5, 325–344 (1956).
- Roman N., Radulescu S., Yancy L., Simionescu O., Scarlat S., Sima A.: Diagnostical biochemic in ascariidosis. Microbiologia (Buc.) 15, 255–258 (1970).

- Saz H. J., Gerson K.: Identification of α -methylvalerate as a product of *Ascaris lumbricoides* fermentation. *Exp. Parasit.* 12, 204–210 (1962).
- Schulte H.: Versuche über Stoffwechselforgänge bei *Ascaris lumbricoides*. *Arch. ges. Physiol.* 166, 1–44 (1917).
- Senutayte Ya. Yu.: Peculiarity of glycolysis in *Fasciola hepatica*. In: Problems of parasitology in the Baltic Republics, p. 114–116, Riga 1970 (in Russian).
- Senutayte Ya. Yu.: Peculiarity of carbohydrate metabolism in *Fasciola hepatica*. *Tr. vsesoyuz. Inst. Helmintologii* 17, 87–89 (1971).
- Shestak E. A.: Peculiarity of carbohydrate metabolism in *Mecistocirrus digitatus* and the inhibitory action of Phenothiazin. Thesis, Moscow 1973 (in Russian).
- Smyth J. D.: The physiology of tapeworms. *Biol. Rev. Cambridge phil. Soc.* 22, 214–236 (1947).
- Soprunov F. F.: *Biochemie der Helminthen*. G. Fischer Verlag, Jena 1978.
- Soprunov F. F., Benedictov I. I., Salmenkova E. A., Grigorovitch Yu. A.: Features of biochemical metabolism of ascaride of swine (*Ascaris suum*). In: Problems of medical parasitology and prevention of infectious diseases, ed. by Sh. D. Moshkovski, p. 472–494, Moscow 1964 (in Russian).
- Soprunova N. Ya.: Biochemical diagnosis of ascaridosis. *Med. Parazit. (Mosk.)* 37, 76–78 (1968).
- Soprunova N. Ya.: Time of detection of volatile fatty acids in the urine of piglets artificially infected with *Ascaris suum*. *Med. Parazit. (Mosk.)* 40, 333–336 (1971).
- Soprunova N. Ya., Andreeva G. A.: Examination of a boarding school in Obninsk of ascaridosis by a biochemical method. *Med. Parazit. (Mosk.)* 37, 359–360 (1968).
- Soprunova N. Ya., Lurje A. A.: Gas-chromatographic determination of volatile fatty acids excreted by helminths in the urine of patients with helminthic diseases. *Med. Parazit. (Mosk.)* 41, 137–141 (1972).
- Soprunova N. Ya., Soprunov F. F., Lurje A. A.: Nachweis von Helminthen-Metaboliten im Harn des Wirtes als ein neuer diagnostischer Test für Helminthiasen. *Angew. Parasit.* 14, 11–17 (1973).
- Toryu J.: Contribution to the physiology of the *Ascaris*. Glycogen content of the *Ascaris megaloccephala*. *Sci. Rep. Tohoku Univ. First Ser.* 4, 65–74 (1933).
- Ueno J.: A method for microdetermination of lower aliphatic fatty acids by paper chromatography and its application to study the metabolism of *Ascaris lumbricoides* v. *suis*. *J. Biochem.* 48, 161–168 (1960).
- Vaastra J. W.: Intermediary metabolism of the cattle lungworm *Dictyocaulus viviparus*. *Hoppe-Seyler's Z. physiol. Chem.* 350, 701–709 (1969).
- Warren L. G., Poole W. R.: Biochemistry of the dog hookworm. II. Nature and origine of the excreted fatty acids. *Exp. Parasit.* 24, 408–416 (1970).
- Yanagisawa T., von Brand T.: Carbohydrate metabolism in *Angiostrongylus cantonensis*. *J. Parasit.* 51, 418–423 (1965).