

Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	40 (1983)
Heft:	1
Artikel:	Host-parasite interactions which influence the virulence of "Trypanosoma (Trypanozoon) brucei brucei" organisms
Autor:	Black, S.J. / Jack, R.M. / Morrison, W.I.
DOI:	https://doi.org/10.5169/seals-313107

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

International Laboratory for Research on Animal Diseases (ILRAD), Nairobi, Kenya

Host-parasite interactions which influence the virulence of *Trypanosoma (Trypanozoon) brucei brucei* organisms

S. J. BLACK, R. M. JACK, W. I. MORRISON

Summary

Subclones were prepared in mice from *T. b. brucei* ILTat 1.4 parasites. Subclones which did not differentiate to stumpy forms in mice were highly virulent and did not stimulate detectable antibody responses. A subclone which did give rise to stumpy forms in mice, was less virulent and did stimulate an antibody response specific for the trypanosome surface glycoprotein. Clones and subclones of *T. b. brucei* parasites which did not give rise to stumpy forms in mice, did give rise to stumpy forms in Bovidae. Plasma from cattle infected with those parasites did not stimulate differentiation of *T. b. brucei* parasites in mice. Murine pleomorphic and monomorphic *T. b. brucei* parasites retained their respective phenotypes in co-infected mice. Both types of parasites were equally pleomorphic in Bovidae. We conclude that some clones of *T. b. brucei* remain monomorphic in mice as a result of a high avidity interaction between slender forms and host molecules which inhibit differentiation of *T. b. brucei* parasites.

Key words: *T. b. brucei*; slender; stumpy; mice; cattle; differentiation; regulation.

Introduction

The rate of differentiation of blood stream *T. b. brucei* parasites from rapidly dividing slender forms through intermediate and stumpy forms to senescent parasites (Black et al., 1982a) influences the parasite population growth rate and the kinetics of the host antibody responses (Sendashonga and Black, 1982). Antibody responses are induced by senescent parasites derived from the stumpy form population but not by dividing parasites. Recent studies suggest that

Correspondence: Doctor Sam J. Black, International Laboratory for Research on Animal Diseases, P.O. Box 30709, Nairobi, Kenya

the host plays a role in regulating the rate of parasite differentiation: The rate of differentiation of the same *T. b. brucei* parasites differs between inbred strains of mice because a host mechanism, which influences the rate of parasite differentiation, is activated to different degrees in these mice during the first parasitaemic wave (Black et al., 1982 b). In this presentation we further investigate the role of the host in regulating parasite differentiation and speculate on the mechanisms involved.

Materials and Methods

Mice. – Male and female BALB/c mice, bred and maintained at ILRAD and aged between 3 and 6 months were used in the study.

Cattle. – Female Hereford cattle, aged 6 months, were bred and maintained at ILRAD under trypanosome-free conditions and were serologically negative for anti-trypanosome antibody prior to use in experiments.

Trypanosomes. – *T. b. brucei* ILTat 1.4 and 1.2 were derived from Lump 227 stock (UHEM-BO/64/EATRO/795). *T. b. brucei* GUTat 3.1 was derived from Treu 667 stock and cloned in BALB/c mice. The properties of these clones of *T. b. brucei* in mice have been described (Sendashonga and Black, 1982). Methods for the purification of trypanosomes from blood have been described (Lanham and Godfrey, 1970). Slender and stumpy differentiation forms were identified in air-dried, methanol-fixed, Giemsa-stained, thin films of infected blood by the method of Wijers (1959). At least 100 parasites were examined per slide. Parasitaemia in mouse blood was assessed by counting parasites in a haemocytometer. Parasitaemia in bovine blood was assessed by the dark ground method of Murray et al. (1977).

Solid phase radioimmunoassay (SRIA). – Serum antibodies to variable surface glycoprotein (VSG) were measured in a SRIA which has been described (Black et al., 1982 a). Briefly, parasites which had been centrifuged onto wells in a flexible polystyrene microtiter plate (10^5 organisms/well, 500 g for 10 min) and fixed with 0.25% glutaraldehyde for 5 min were used as target antigen. The assay was conducted as for SRIA against other antigens using affinity labelled ^{125}I anti-mouse IgM or IgG as a second step reagent (Tsu and Herzenberg, 1980). The \log_{10} reciprocal titer of antibodies was calculated by comparing the 50% binding points of pre-infection and post-infection sera. Results are presented as the arithmetic mean ± 1 standard deviation (SD).

Indirect immunofluorescence. – Purified ILTat 1.4 trypanosomes were spread as a thin film on glass slides, air-dried for 10 min, fixed in cold (4°C) acetone for 10 min and incubated with a staining buffer containing 0.01 M NaPO₄ (pH 8.1), 0.15 M NaCl and 1% bovine serum albumin for 30 min. The parasites were reacted for 30 min with monoclonal antibodies V3/16.15.1 and V3/66.1.1 which recognize internal and external epitopes of ILTat 1.4 VSG respectively (Pearson et al., 1980; Black et al., 1982 a). The slides were washed with 20 ml of staining buffer and exposed to staining buffer containing fluorescein conjugated anti-mouse Ig (Nordic Immunological Laboratories B.V., Tilburg, The Netherlands) for 30 min. The slides were washed and stained trypanosomes were examined on a UV microscope (Leitz, Wetzlar, W. Germany) using a 63/1.3 oel Phaco 3 Fluoresz objective and GW 6.3 \times eyepieces.

Analyses of ILTat 1.4 trypanosomes in thin smears prepared from the buffy coat of washed mouse blood (3 \times in RPMI 1640 medium) were conducted in the same manner. Trypanosomes present in the preparations which did not express the ILTat 1.4 VSG were identified by examining the slides under UV and phase contrast illumination. Results are presented as the arithmetic mean ± 1 SD and at least 200 parasites were examined in each preparation.

Irradiation. – BALB/c mice were given 800 rads of γ irradiation delivered by a $^{137}\text{Cesium}$ source. This dose depletes the B- and T-cell dependent areas of the spleen in 6 days and kills the recipients in 10 to 14 days.

Results

*Subclones of *T. b. brucei* ILTat 1.4 Parasites are heterogeneous with respect to differentiation rates and their capacities to induce detectable antibody responses in BALB/c mice*

BALB/c mice, inoculated with 50 or more *T. b. brucei* ILTat 1.4 parasites die during the first parasitaemic wave with a parasitaemia of $>10^9$ parasites/ml. At the time of death of the host, the parasite population in the blood is composed of approximately 95% slender form, 4.8% intermediate form and 0.2% stumpy form parasites. Seventeen subclones of *T. b. brucei* ILTat 1.4 parasites were prepared in lethally irradiated BALB/c mice. All of the subclones expressed the ILTat 1.4 VSG to a level greater than 99.9% of the parasites examined, using the monoclonal antibodies V3/16.15.1 and V3/66.1.1.

Sixteen of the subclones, represented by No. 1 in Table 1, did not give rise to stumpy form parasites in BALB/c mice and did not stimulate a VSG-specific antibody response. In contrast, subclone 17 (Table 1) did give rise to stumpy form parasites in BALB/c mice and stimulated a VSG-specific antibody response. The BALB/c mice were inoculated intravenously with 50 parasites. Recipients of subclone 1 died after 8 days whereas recipients of subclone 17 died after 15 days.

**T. b. brucei* ILTat 1.4 subclone 1, ILTat 1.2 and GUTat 3.1 parasites give rise to stumpy forms in Bovidae*

Two calves were inoculated intravenously with either 10^5 *T. b. brucei* ILTat 1.4 subclone 1 or 10^5 *T. b. brucei* GUTat 3.1 parasites (a mouse pleomorphic parasite which when recloned only gives rise to pleomorphic parasites in mice). The blood parasitaemia and morphological types were assessed daily and results are presented in Table 2. Both sets of parasites differentiated at a similar rate in cattle although the pre-patent period of the ILTat 1.4 subclone 1 parasites was shorter than that of the GUTat 3.1 parasites. Differences in the pre-patent periods might have resulted from the proportionally greater number of slender form parasites present in the *T. b. brucei* ILTat 1.4 subclone 1 inoculum. Similar results were obtained using another mouse monomorphic clone of *T. b. brucei*, ILTat 1.2 (Sendashonga and Black, 1982). *T. b. brucei* GUTat 3.1, ILTat 1.4 subclone 1 and ILTat 1.2 parasites also gave rise to stumpy forms in sheep, domestic cattle, waterbuck, eland and african buffalo (Grootenhuis, Whitelaw and Black, unpublished).

*The same *T. b. brucei* ILTat 1.4 and 1.2 parasites which are pleomorphic in Bovidae, are monomorphic in mice*

T. b. brucei ILTat 1.4 subclone 1 parasites which were harvested at 5 days from the Hereford calf described in Table 2, were cloned into lethally irradiated BALB/c mice. All of the 10 subclones which were obtained expressed the ILTat

Table 1. Immune responses to and differentiation of subclones of *T. b. brucei* ILTat 1.4

<i>T. b. brucei</i> ILTat 1.4 Subclone No.	Day of analysis	Log ₁₀ parasites/ml blood ± 1SD	% ± 1SD of parasite morphological types		Log ₁₀ reciprocal titer ± 1SD antibodies on <i>T. b. brucei</i> ILTat 1.4	
			Intermediate	Stumpy	IgM	IgG
17	4	5.0 ± 0.3	0	0	0	0
	6	7.5 ± 0.2	0	0	0	0
	7	8.8 ± 0.3	0.5 ± 0.2	0	0	0
	4	4.0 ± 0.2	0	0	0	0
	6	6.2 ± 0.3	3.0 ± 2.0	0.5 ± 0.3	0.5 ± 0.2	0
	8	8.2 ± 0.3	19.0 ± 3.0	3.0 ± 2.0	1.0 ± 0.5	0.5 ± 0.2
	9	8.9 ± 0.2	32.0 ± 4.0	5.0 ± 3.0	1.5 ± 0.2	0.7 ± 0.2
	10	8.1 ± 0.4	nt	nt	2.8 ± 0.4	1.5 ± 0.2

50 parasites were inoculated i.v.

No antibody binding was detected on *T. b. brucei* ILTat 1.2.

nt = not tested

Table 2. *T. b. brucei* ILTat 1.4 subclone 1 and GUTat 3.1 parasites differentiate at similar rates in Hereford calves

<i>T. b. brucei</i>	Day after infection	Log ₁₀ parasites/ml blood	% of parasite morphological types:	
			Intermediate	Stumpy
ILTat 1.4 Subclone 1	4	5.0	7	0
	5	6.1	19	5
	6	6.4	38	50
	7	6.0	11	85
	8	3.0	nt	nt
GUTat 3.1	5	5.1	10	0
	6	6.2	24	6
	7	5.6	36	55
	8	5.9	13	82
	9	3.0	nt	nt

6-month-old female Hereford calves were inoculated intravenously with 10⁵ parasites.

nt = not tested

1.4 VSG as detected by indirect immunofluorescence with monoclonal antibodies V3/16.15.1 and V3/66.1.1 and did not differentiate from slender to stumpy form parasites in intact or irradiated BALB/c mice. *T. b. brucei* parasites inoculated into mice from the third parasitaemic wave of the calf also did not give rise to stumpy form trypanosomes. The studies were repeated using ILTat 1.2 parasites and similar results were obtained.

*Plasma from *T. b. brucei* infected cattle does not contain molecules which stimulate differentiation of *T. b. brucei* ILTat series parasites in mice*

Plasma prepared from normal cattle, from cattle in which ILTat 1.4 parasites had switched to stumpy forms or from cattle which had just cleared the first parasitaemic wave of *T. b. brucei* ILTat 1.4 parasites, did not induce *T. b. brucei* ILTat 1.2 parasites to switch to intermediate or stumpy forms in intact or irradiated BALB/c mice. The mice were inoculated intraperitoneally with 10² *T. b. brucei* ILTat 1.2 parasites and after the first day were inoculated intravenously with 0.25 ml of the bovine plasma every day until death which occurred between 6 and 7 days after infection. The reciprocal experiment was performed using mice infected with *T. b. brucei* ILTat 1.4 and plasma from cattle infected with *T. b. brucei* ILTat 1.2 or GUTat 3.1. No stumpy form parasites were detected nor was there any increase in the percentage of intermediate form parasites detected.

Table 3. Influence of co-infection on differentiation of *T. b. brucei* ILTat 1.4 and GUTat 3.1 in mice.

<i>T. b. brucei</i> inoculation	Day after infection	Log ₁₀ parasites/ml blood ± 1SD	% GUTat 3.1 ± 1SD	% stumpy forms
10 ² ILTat 1.4*	4	6.3 ± 0.2	0	0
	5	8.0 ± 0.2	0	0
	6	9.3 ± 0.2	0	0
5 × 10 ³ GUTat 3.1	4	6.0 ± 0.1	100	0
	5	7.6 ± 0.1	100	8 ± 2
	6	7.4 ± 0.1	100	58 ± 17
	7	<5.0		
10 ² ILTat 1.4* and 5 × 10 ³ GUTat 3.1	4	6.5 ± 0.1	41 ± 10	0
	5	8.2 ± 0.2	40 ± 16	0
	6	9.2 ± 0.2	0.9 ± 0.7	0.6 ± 0.4

* ILTat 1.4 parasites were identified by indirect immunofluorescence using monoclonal antibodies reactive with ILTat 1.4 VSG. No fluorescent stumpy form parasites were observed. GUTat 3.1 parasites were identified under phase contrast and UV illumination as those trypanosomes which did not express ILTat 1.4 VSG. All recipients of ILTat 1.4 died between 6 and 7 days after inoculation.

*Pleomorphic and monomorphic *T. b. brucei* parasites retain their respective phenotypes in co-infected BALB/c mice*

BALB/c mice were inoculated intraperitoneally with 10² *T. b. brucei* ILTat 1.4, 10³ *T. b. brucei* GUTat 3.1 or 10² *T. b. brucei* ILTat 1.4 and 10³ *T. b. brucei* GUTat 3.1. Each group contained 10 mice. After 4, 5 and 6 days, the parasitaemia in each mouse was counted, thin blood films were prepared for analysis of parasite morphology and blood was collected from the retro-orbital plexus of each mouse into heparinized pipettes. The blood was washed and thin smears of buffy coat were prepared. The smears were stained for indirect immunofluorescence with antibodies reactive with ILTat 1.4 VSG and were observed under UV and phase contrast illumination. Parasites with ILTat 1.4 VSG were taken to be ILTat 1.4 parasites whereas those without ILTat 1.4 VSG were taken to be GUTat 3.1 parasites. Results are presented in Table 3 which shows that ILTat 1.4 parasites in co-infected mice continued to multiply and remained as slender forms whereas the GUTat 3.1 parasites switched to stumpy forms and were overgrown. Differentiation of *T. b. brucei* GUTat 3.1 parasites to stumpy forms was delayed by 1 day in co-infected hosts. In the co-infected hosts, the proportional representation of GUTat 3.1 parasites by 6 days after infection was approximately 1% of the total parasites detected (i.e. about 2 × 10⁷/ml) and most if not all, of these parasites had switched to stumpy forms. The experiment was performed several times in both intact and irradiated (650 rads) BALB/c mice with similar results.

Discussion

The studies show that a cloned population of *T. b. brucei* parasites can contain members with the same VSG which are heterogeneous in virulence, rates of differentiation to intermediate and stumpy forms and capacities to induce detectable antibody responses in mice. The studies confirm previous observations which show that the degree of pleomorphism of *T. b. brucei* parasites is not linked to VSG type (Barry et al., 1979) and that antibody responses might be induced by senescent parasites derived from the stumpy form population but not by actively dividing parasites (Sendashonga and Black, 1982).

T. b. brucei parasites which were monomorphic in mice, were pleomorphic in cattle and when cloned back into mice from the first or third parasitaemic wave in a calf, were monomorphic. The results confirm previous observations (Black et al., 1982 b) which suggest that differentiation of *T. b. brucei* parasites is under host control. Plasma collected from cattle in which the murine monomorphic *T. b. brucei* parasites had differentiated to stumpy forms, did not induce differentiation of those parasites when injected into mice. The observations suggest that the bovine plasma did not contain molecules which stimulate differentiation *T. b. brucei* parasites. We tentatively suggest that differentiation of *T. b. brucei* parasites is negatively regulated by the host, i. e. there are variable concentrations of molecules which encourage growth of the parasites and prevent differentiation.

Murine monomorphic and pleomorphic *T. b. brucei* parasites were shown to retain their respective phenotypes in co-infected mice although differentiation of the pleomorphic parasites was slightly retarded. This suggests that murine molecules which encourage growth of *T. b. brucei* parasites are more efficiently perceived or utilized by monomorphic than pleomorphic parasites. Because murine monomorphic *T. b. brucei* were pleomorphic in cattle and did not grow more quickly than murine pleomorphic parasites when inoculated into cattle, it is likely that murine and bovine molecules which encourage *T. b. brucei* growth differ in some respect.

The rate of differentiation of *T. b. brucei* parasites directly influences parasitaemia (Black et al., 1982 b) and the kinetics of development of the host protective antibody response (Sendashonga and Black, 1982). Elucidation of the host molecules which inhibit *T. b. brucei* differentiation and the trypanosome receptor site for those molecules, might suggest new methods for control of infections with trypanosomes and new ways of examining the trait of trypanotolerance.

Acknowledgments

We would like to thank Mr. Francis McOdimba for excellent technical assistance and Ms. Leah Agina for manuscript preparation. – This is ILRAD Publication Serial No. 213.

Barry J. D., Le Ray D., Herbert W. J.: Infectivity and virulence of *Trypanosoma brucei* for mice: dissociation of virulence and VAT in relation to pleomorphism. *J. comp. Path.* 89, 465–470 (1979).

Black S. J., Hewett R. S., Sendashonga C. N.: *Trypanosoma brucei* variable surface coat is released by degenerating parasites but not by actively dividing parasites. *Parasite Immunol.* 4, 233–244 (1982 a).

Black S. J., Sendashonga C. N., Lalor P. A., Whitelaw D. D., Jack R. M., Morrison W. I., Murray M.: Regulation of the growth and differentiation of *Trypanosoma brucei* in resistant (C57 B1/6) and susceptible (C3H/He) mice. *Parasite Immunol.* (in press) (1982 b).

Lanham S. M., Godfrey D. G.: Isolation of salivarian trypanosomes from man and other mammals using DEAE cellulose. *Exp. Parasit.* 28, 521–534 (1970).

Murray M., Murray P. K., McIntyre W. I. M.: An improved parasitological technique for the diagnosis of African trypanosomiasis. *Trans. roy. Soc. trop. Med. Hyg.* 71, 325–326 (1977).

Pearson T. W., Pinder M., Roelants G. E., Kar S. K., Lundin L. B., Mayor-Withey K. S., Hewett R. S.: Methods for derivation and analysis of anti-parasite monoclonal antibodies. *J. immunol. Meth.* 34, 141–154 (1980).

Sendashonga C. N., Black S. J.: Humoral immune responses against *Trypanosoma brucei* variable surface antigens are induced by degenerating parasites. *Parasite Immunol.* 4, 245–257 (1982).

Tsu T. T., Herzenberg L. A.: Solid phase radioimmunoassays. In: *Selected methods in cellular immunology*, ed. by B. B. Mishell and S. M. Shiigi, p. 377–397. Freeman, San Francisco 1980.

Wijers D. J. B.: Polymorphism in *Trypanosoma gambiense* and *Trypanosoma rhodesiense* and the significance of intermediate forms. *Ann. trop. Med. Parasit.* 53, 59–68 (1959).