Zeitschrift: Acta Tropica

Herausgeber: Schweizerisches Tropeninstitut (Basel)

Band: 40 (1983)

Heft: 3

Artikel: The influence of schistosomiasis on the serum concentrations of retinol

and retinol binding protein of a rural population in Liberia

Autor: Stürchler, D. / Holzer, B. / Hanck, A. DOI: https://doi.org/10.5169/seals-313135

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 03.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

The influence of schistosomiasis on the serum concentrations of retinol and retinol binding protein of a rural population in Liberia

D. STÜRCHLER¹, B. HOLZER², A. HANCK³, A. DEGRÉMONT¹

Summary

121 persons from rural areas of central Liberia were examined for possible interactions between helminthic infections and the retinol (vitamin A) and retinol binding protein (= RBP) status. About \(\frac{1}{2} \) of this mainly adult population had retinol and RBP serum concentrations $\leq 400 \,\mu\text{g/l}$ and $\leq 40 \,\mu\text{g/ml}$, respectively. Only one person had a serum retinol concentration $< 100 \mu g/1$ which is critical for the appearance of xerophthalmia. The retinol and RBP concentrations showed a linear, positive correlation. 19 parasitologically helminth free individuals had mean serum retinol and RBP concentrations of 414 μ g/l and 43 μ g/ml, respectively, while the corresponding values of 20 individuals with schistosomiasis as the only helminthic infection were 339 μ g/l and 35 µg/ml. 65 other persons with mixed intestinal and/or tissue helminthiases, and 9 persons with pure Onchocerca volvulus infection exhibited intermediate mean concentrations. 32 persons without serological evidence of helminthic infections had higher serum concentrations of retinol and RBP than 32 individuals in whose sera at least one raised antibody titer was found in the enzyme-linked immunosorbent assay with 3 different helminthic test antigens. In addition to age, sex and pregnancy schistosomiasis might be a further risk factor of latent retinol and RBP deficiency among adults and teenagers living in a schistosomiasis endemic area.

Key words: retinol; vitamin A; retinol binding protein; schistosomiasis; *Schistosoma haematobium; Schistosoma mansoni; Onchocerca volvulus;* immunity; enzyme-linked immunosorbent assay; Liberia.

Correspondence: Dr. D. Stürchler, University of Washington, School of Public Health, Department of Epidemiology, SC-36, Seattle, WA 98195, USA

¹ Medical Department, Swiss Tropical Institute, Basel, Switzerland

² Liberian Institute for Biomedical Research, Robertsfield, Liberia

³ Pharmaceutical Division, F. Hoffmann-La Roche, Basel, Switzerland

Introduction

Retinol (vitamin A) is a fat soluble vitamin which is essential for the maintenance of the normal functions of epithelial and mucous membranes in humans. In well nourished individuals the serum concentrations of retinol and of the retinol binding protein (= RBP) are above 400 μ g/l and 40 μ g/ml, respectively (WHO, 1976; De Luca et al., 1979). In retinol deficiency the reserves in the liver get exhausted and the serum concentration is between 100 and 400 μ g/l. In clinically manifest retinol deficiency the serum concentration is below 100 μ g/l (WHO, 1976). In protein deficient individuals the RBP serum concentration is likely to be lower than 35 to 40 μ g/ml (De Luca et al., 1979).

Subclinical and clinical retinol deficiencies are prevalent in many tropical countries, often in association with protein deficiency, diarrhoea and with viral, bacterial and parasitic infections (Scrimshaw et al., 1968). In the rain forest area of West Africa retinol deficiency seems rare because red palm oil, which is rich in retinol and its precursors, is widely consumed. In this same area schistosomiasis, onchocerciasis and ancylostomiasis are endemic (Saladin et al., 1980; Stürchler et al., 1980). On the occasion of a field study on the epidemiology and morbidity of schistosomiasis in Liberia we wanted to find possible correlations between the serum concentrations of retinol and of BRP and the prevalence of schistosomiasis and other parasitoses in a rural population.

Study population, Materials and Methods

The field investigation took place during the dry season of 1980/81, in central Liberia where previous work had identified three different zones of schistosomiasis transmission (Saladin et al., 1980): I: no transmission, II: transmission of *Schistosoma haematobium* only, and III: transmission of both *S. haematobium* and *S. mansoni*.

From zones I, II and III the population was sampled for the schistosomiasis morbidity study (Holzer et al., 1983). These were random samples from zone II (Zeansue village) and III (Synea village). In zone I (Flehla village) the town chief indicated the individuals who were to be examined. Of Synea only pooled sera were available for retinol determination. In addition a group of outpatients consulting at the Phebe hospital in Suakoko for symptoms suggesting schistosomiasis were investigated. The data of 79 females and 42 males with an age range of 5 to 35 years were available for investigation.

The details of the clinical, laboratory and parasitological methods are described in Holzer et al. (1983) and Saladin et al. (1983). All participants completed a questionnaire and had a physical examination. Urine was filtered for eggs of *S. haematobium* and tested with paper strips for proteinuria and haematuria and with a dip-slide agar for bacteriuria. Stool was analysed with formolether concentration (Knight et al., 1976) for eggs of *S. mansoni* and other helminths. Skin-snips from the left iliac crest for microfilariae of *Onchocerca volvulus* were placed into normal saline solution and read after 3 and 6 hours. Hematocrit, total white blood cell count and blood eosinophilia were determined with capillary blood.

Sera were processed in Basel. The enzyme-linked immunosorbent assay (= ELISA) was used for the serodiagnosis of schistosomiasis, filariasis, toxocariasis and amebiasis (Speiser, 1982). Doubtfully positive ELISA extinction values were 0.15–0.3 for schistosomiasis, 0.3–0.5 for amebiasis, and 0.5–0.7 for filariasis and toxocariasis. The indirect immunofluorescent antibody test (= IFAT) was used for the serodiagnosis of malaria. Serum retinol concentrations were determined as

Table 1. Serum concentrations of retinol and retinol binding protein (= RBP) in the study population. Adequate retinol and RBP concentrations are $>400 \,\mu\text{g}/\text{l}$ and $>40 \,\mu\text{g}/\text{ml}$, respectively.

	Retinol in μ g/l		RBP in μg/ml	
	N	mean (±SD)	N	mean (±SD)
Total population	121	379.1 (139.2)	116	38.3 (13.1)
Sex				
Females	79	362.2a (138.2)	75	36.6 ^b (12.2)
Males	42	410.8a (137.3)	41	41.7 ^b (14.1)
Age: full years				
5–15	13	332.3° (94.7)	13	31.8 ^d (7.9)
16–25	67	371.1 (108.1)	63	37.8 (11.5)
26–35	41	407.4° (185.8)	40	41.3 ^d (15.8)
Location				
Flehla (zone I)	30	470.7*		
Zeansue (zone II)	39	406.6e (128.2)	37	$40.4^{\rm f}$ (11.0)
Synea (zone III)	39	387.1 (119.3)	39	40.2 (11.7)
Suakoko (outpatients)	43	346.8e (160.6)	40	34.6 ^f (15.5)

$$a-a$$
 $t = 1.8, 2p: n.s.$ $b-b$ $t = 2.1, 2p < 0.05$ $c-c$ $t = 1.9, 2p: n.s.$ $d-d$ $t = 1.9, 2p: n.s.$ $e-e$ $t = 1.9, 2p: n.s.$ $f-f$ $t = 1.9, 2p: n.s.$

follows: 100 µl serum samples were taken up in n-hexane with 2% isopropanol, then high pressure liquid chromatography was performed with a LiChrosorb (E. Merck, Darmstadt) Si 60 5 μ m 25 cm as fixed phase. Detection was done with UV 325 nm (Uvikon LCD 725, Kontron AG, Zürich). Quantification was carried out with planimetry of the peaks in relation to an external standard. RBP was determined by immunodiffusion assay.

For probability statements t was calculated.

Results

121 persons were investigated. None of them was severely ill at the time of examination or presented with signs of marasmus or kwashiorkor. A low hematocrit ($\leq 35\%$ in females, $\leq 38\%$ in males) was found in 27 females and in 7 males. 58 persons (48%) had eggs of S. haematobium in the urine, 51 persons had microhematuria (among them 41 were females), 16 had proteinuria (++ or +++) and 12 had bacteriuria (> 10^6 germs/ml). In a single stool sample of 45 persons (37%) eggs of S. mansoni were found. 37 persons had ancylostomiasis, 29 trichuriasis and 15 ascaridiasis. 54 persons (45%) were infected with Onchocerca volvulus.

Except for a 19-year-old girl the whole study population was reactive to Plasmodium falciparum antigen in the IFAT. 32 persons did not respond to the various helminthic test antigens in the ELISA.

Table 2. Serum retinol and retinol binding protein (= RBP) in *females* by pregnancy and hematocrit

	Retinol in μ g/l		RBP in μ g/ml	
	N	mean (±SD)	N	mean (±SD)
History of pregnancy				
Yes	6	318.8 (55.0)	6	33.2 (4.2)
No	73	365.8 (142.5)	69	36.8 (12.6)
Hematocrit				
>35%	52	378.0 (135.5)	49	37.9 (11.0)
≤35%	27	331.8 (140.7)	26	33.8 (14.0)

Table 3. Serum concentrations of retinol and retinol binding protein (= RBP) in relation to parasitological findings

	Retinol in μ g/l		RBP in μ g/ml	
	N	mean (±SD)	N	mean (±SD)
Helminth free persons	19	413.8a,b (147.1)	19	43.0 ^{c,d} (14.2)
All with helminths	102	369.7 ^a (130.5)	97	37.1° (12.0)
Single helminthiases				
Schistosomiasis	20	339.0 ^b (137.9)	20	34.5 ^d (10.9)
Onchocerciasis	9	394.0 (151.6)	9	37.0 (14.9)
ntestinal nematodes	8	370.8 (109.6)	7	34.9 (14.6)
Multiple helminthiases*	65	375.2 (128.9)	61	38.2 (11.8)

^{*} Schistosoma sp. plus O. volvulus plus intestinal nematodes: 21

$$a-a$$
 t = 1.2, p: n.s.
 $b-b$ t = 1.6, 2p: n.s.

$$c-c$$
 t = 1.7, 2p: n.s.

$$d-d$$
 t = 2.6, 2p < 0.02

Retinol and retinol binding protein (= RBP) by sex, age and village (Table 1). Serum retinol concentrations were $\leq 400 \, \mu g/l$ in 77 persons (64%), those of RBP were $\leq 40 \, \mu g/ml$ in 73 (63%). The lowest retinol concentration (72 $\mu g/l$) was observed in a woman consulting at Suakoko for intestinal schistosomiasis. Females had lower mean retinol and RBP concentrations than males. Pregnant females had lower values than non-pregnant females (Table 2). Adults had a better retinol and RBP status than children and teenagers. Schistosomiasis outpatients and the persons from Synea (zone III) had mean retinol serum concentrations $\leq 400 \, \mu g/l$.

Schistosoma sp. plus intestinal nematodes: 20

Schistosoma sp. plus O. volvulus: 14

O. volvulus plus intestinal nematodes: 10




Fig. 1. Scatterdiagram of the serum concentrations of retinol (= R) and of retinol binding protein (= RBP), by parasitological findings. A = helminth free subjects; B = Schistosoma free subjects; C = Schistosoma-infected subjects. C₁: with S. haematobium only; C₂: with S. haematobium and S. mansoni; C₃: with S. mansoni only. Vertical traces indicate medians.

Retinol, RBP and biochemical results. Males and females (Table 2) with low hematocrits had lower mean retinol and RBP concentrations than their mates with adequate hematocrits. The presence of neither proteinuria nor bacteriuria seemed to influence the retinol status of the population. However, men with microhematuria had lower mean retinol and RBB concentrations (359 μ g/l and 37 μ g/ml, respectively) than men without hematuria (with corresponding concentrations of 418 μ g/l and 42 μ g/ml, respectively).

Retinol, RBP and parasitological results (Table 3, Fig. 1). Helminth free individuals (11 females, 8 males) had the highest mean retinol and RBP concentrations and persons with pure schistosomiasis had the lowest. Persons with pure onchocerciasis or with polyparasitism had intermediate values. 5 persons (4 females, 1 male) with pure *S. haematobium* infection had mean retinol and RBP concentrations of 422 μ g/l and 43 μ g/ml, respectively. 5 other persons (again 4 females, 1 male) with pure *S. mansoni* infection had corresponding values of 353 μ g/l and 34 μ g/ml, respectively. 10 of 19 helminth free persons, and 16 of 20 with pure schistosomiasis had retinol concentrations <400 μ g/l.

Table 4. Serum concentrations of retinol and retinol binding protein (= RBP), and the presence of anti-helminthic antibodies as measured by the enzyme-linked immunosorbent assay (= ELISA)

	Retinol in μ g/l		RBP in μ g/ml	
	N	mean (±SD)	N	mean (±SD)
No anti-helminthic antibodies	32	425.7 ^{a.c} (140.7)	29	42.2 ^{b,d} (13.1)
All with anti-helminthic antibodies				
Only doubtful reactions*	6	361.8 (84.1)	6	36.3 (6.8)
At least one strong reaction*	32	321.9^a (101.4)	31	35.2^{b} (11.2)
Anti-schistosomal antibodies only				
Doubtful and strong reactions*	35	371.8 (146.1)	35	36.8 (14.1)
Only strong reaction*	12	339.3° (147.6)	12	33.8 ^d (13.8)
Anti-filarial and/or anti-toxocaral anti	ibodies (only		
Doubtful and strong reactions*	16	403.7 (131.4)	15	37.5 (13.9)
Only strong reaction*	8	351.6 (106.8)	5	ä36.8 (6.1)

^{*} For cut-off points see Materials and Methods

$$a-a$$
 $t = 3.4, 2p < 0.002$ $b-b$ $t = 2.3, 2p < 0.05$ $c-c$ $t = 1.8, 2p: n.s.$ $d-d$ $t = 1.8, 2p: n.s.$

Retinol, RBP and serological results (Table 4). Persons without anti-helminthic antibodies demonstrable in ELISA had higher mean retinol and RBP concentrations than persons whose sera gave a positive ELISA reaction with one or more helminthic test antigens. When among the 32 non-responders (19 females, 14 males) those (11 subjects) with parasitologically demonstrated *Schistosoma* infection were removed, the mean retinol and RBP concentrations in the remaining 21 persons without serological or parasitological evidence of schistosomiasis rose to $453 \pm 136 \,\mu\text{g}/\text{l}$ and $44 \pm 14 \,\mu\text{g}/\text{ml}$, respectively. 8 persons with neither parasitological nor serological evidence of any helminthiasis had mean retinol and RBP concentrations of $479 \,\mu\text{g}/\text{l}$ and $49 \,\mu\text{g}/\text{ml}$, respectively.

11 persons exhibited antibodies against *Entamoeba histolytica* in ELISA (extinction ≥ 0.3). They had mean retinol and RBP values of 276 μ g/l and 25 μ g/ml, respectively.

The relation of retinol and RBP. Within the range of observed values retinol and RBP showed a linear, positive correlation (r = 0.6, p < 0.001).

Discussion

This study forms part of an investigation of various aspects of schistosomiasis in central Liberia (Saladin et al., 1980; Saladin et al., 1983; Holzer et al., 1983). The sampling procedure and the physical examination were designed for symptoms and signs caused by schistosomiasis, not by retinol deficiency.

However, general nutritional aspects were regarded including the measurement of height and weight.

The population at risk of retinol deficiency are mainly pre-school children in tropical countries (Oomen et al., 1964; WHO, 1976; Sommer et al., 1981; WHO, 1982). Mechanisms leading to retinol deficiency are: a) inadequate composition or quantity of the diet, b) inadequate storing or cooking of food stuffs (Bauernfeind, 1978), c) impaired conversion of retinol equivalents to retinol within the intestinal tract, d) malabsorption of retinol, particularly when diarrhea or intestinal parasitoses such as ascaridiasis are present, and e) non-delivery to the periphery because the retinol binding protein (= RBP) is deficient (WHO, 1976; De Luca, 1978; De Luca et al., 1979).

In Africa retinol deficiency is more frequent and more often manifests itself clinically in the dry and savannah regions than in the rain forest area, because in the latter the consumption of red palm oil and mango fruits prevents deficiency (Oomen et al., 1964; Le François et al., 1980). The staple food in central Liberia is rice with red palm oil sauce and leafy vegetables. Fruits, fish, shrimps and meat are eaten occasionally. 100 g of raw mangos contain 190 μ g of retinol equivalents compared to 15,000 µg in 100 g of red palm oil (Périssé and Polacchi, 1979). It is, therefore, not surprising that only 1 individual had a critically low serum retinol concentration and that signs of clinically manifest deficiency were not found. However, about \(^2\)3 of the population investigated, mainly older children and adults, had biochemical signs of both retinol and RBP deficiency. Lauber and Haller (1980) made similar findings among children in the rain forest of the Ivory Coast. It is possible that the biochemical criteria of retinol deficiency have to be re-evaluated for some tropical areas. It is, however, more likely, that one of the above mentioned deficiency mechanisms is responsible for the discrepancy between a seemingly adequate diet and the biochemically inadequate retinol and RBP status observed.

Here intestinal infections come into play. The interactions between retinol, the host's immune response and viral, bacterial and parasitic infections are multi-directional and complex (Scrimshaw et al., 1968; Lee and Aboko-Cole, 1979; Glazebrook and Davis, 1979; Beisel et al., 1981; WHO, 1982). In retinol deficient persons the impaired epithelial barrier might facilitate the penetration of helminth larvae through the skin or the intestinal mucosa. Such a mechanism has been demonstrated in experimentally infected retinol deficient animals (Darip et al., 1979). Like in experimental conditions (Zile et al., 1979; Nauss et al., 1979) in retinol deficient persons the cellular immunity might be depressed. On the other hand intestinal parasites may impair the absorption of retinol. Our own preliminary investigations (Stürchler et al., 1981) point to the possibility that growing helminth larvae and reproductive adult worms might need some of the host's retinol.

The epidemiological, parasitological and serological results of this study are all in favour of an inverse correlation between schistosomiasis on the one

hand and the retinol and RBP status on the other hand. However, this field study does not sort out the contribution of the various parasite species found with certainty. Also, those persons without serological and/or parasitological evidence of schistosomiasis or other helminthiases and with a better biochemical retinol and RBP status than the infected ones might differ from the rest of the population by a better socioeconomical status and by better nutrition. Other studies gave contradictory results. Haller and Lauber (1980) found a correlation of low retinol serum concentrations with onchocerciasis and ascaridiasis, but not with schistosomiasis, in children of the Ivory Coast, while Mansour et al. (1979) observed lower retinol serum concentrations in adult Egyptians with variously severe forms of intestinal schistosomiasis with or without chronic salmonellosis than in uninfected controls.

A causal relationship could only be demonstrated with certainty by animal experiments or by studying the in vitro retinol uptake of adult schistosomes.

Acknowledgments

The authors gratefully acknowledge the valuable help of Dr. A. P. Hanson, Director of the Liberian Institute of Biomedical Research, in conducting this study and in offering adequate facilities at his Institute. The same thank they express to Prof. Dr. T. A. Freyvogel, Director of the Swiss Tropical Institute. They also thank Dr. B. Saladin and Mrs. K. Saladin for the results of the parasitological stool analyses, Mr. F. Speiser for the serodiagnostic results and Dr. M. Tanner for the critical review of the manuscript. Part of the study has been supported by a grant of the R. Geigy Foundation.

- Bauernfeind J. C.: The technology of vitamin A. Invitational presentation, XI. Int. Congr. of Nutrition, Rio de Janeiro, August 27–September 1, 1978.
- Beisel W. R., Edelman R., Nauss K., Suskind R. M.: Single-nutrient effects on immunologic functions. J. Amer. med. Ass. 245, 53-58 (1981).
- Darip M. D., Sirisinha S., Lamb A. J.: Effect of vitamin A deficiency on susceptibility of rats to *Angiostrongylus cantonensis*. Proc. Soc. exp. Biol. (N.Y.) *161*, 600–604 (1979).
- De Luca H. F.: Vitamin A. In: The fat-soluble vitamins, p. 1–67. Plenum Press, New York 1978.
- De Luca L. M., Glover J., Heller J., Olson J. A., Underwood B.: Recent advances in the metabolism and function of vitamin A and their relationship to applied nutrition. Int. Vitamin A Consultative Group, The Nutrition Foundation, New York 1979.
- Glazebrook P. W., Davis G. H. G.: A review of the interactions of infection and nutrition. Asian J. infect. Dis. 3, 93–118 (1979).
- Haller L., Lauber E.: Etude des interactions entre les taux sériques de vitamines et les parasitoses communément répandues en zone tropicale. Acta trop. (Basel) *37*, Suppl. 11, 110–119 (1980).
- Holzer B., Saladin K., Saladin B., Dennis E., Degrémont A.: The impact of schistosomiasis among rural populations in Liberia. Acta trop. (Basel) 40, 239–259 (1983).
- Knight W. B., Hiatt R. A., Cline B. L., Ritchie L. S.: A modification of formol-ether concentration technique for increased sensitivity in detecting *Schistosoma mansoni* eggs. Amer. J. trop. Med. Hyg. 25, 818–823 (1976).
- Lauber E., Haller L.: Déterminations des taux sanguins des vitamines A, B₁, B₂, B₆, B₁₂, C, niacine et folates. Etude de leur relation avec la clinique et l'alimentation. Acta trop. (Basel) *37*, Suppl. 11, 90–109 (1980).

- Lee C. M., Aboko-Cole G. F.: The interaction of nutrition and infection: a succinct review. J. nat. med. Ass. (N.Y.) 71, 765–768 (1979).
- Le François P., Chevassus-Agnes S., Benefice E., Dyck J. L., Maire B., Parent G., Seymat G., Ndiaye A. M.: Vitamin A status of populations in three West African countries. Int. J. Vit. Nutr. Res. 50, 352–363 (1980).
- Mansour M. M., Mikhail M. M., Farid Z., Bassily S.: Chronic salmonella septicemia and malabsorption of vitamin A. Amer. J. clin. Nutr. 32, 319–324 (1979).
- Nauss K. M., Mark D. A., Suskind R. M.: The effect of vitamin A deficiency on the in vitro cellular immune response of rats. J. Nutr. 109, 1815–1923 (1979).
- Oomen H. A. P. C., McLaren D. S., Escapini H.: A global survey on xerophthalmia. Trop. geogr. Med. 16, 271–315 (1964).
- Périssé J., Polacchi W.: Paper presented at the international vitamin A consultative group meeting, Cebu, Philippines, 26–29 November 1979.
- Saladin B., Saladin K., Dennis E., Degrémont A.: Preliminary epidemiological survey of schistosomiasis in central and southern Liberia. Acta trop. (Basel) *37*, 53–62 (1980).
- Saladin B., Saladin K., Holzer B., Dennis E., Hanson A., Degrémont A.: A pilot control trial of schistosomiasis by chemotherapy of target population combined with focal mollusciciding in central Liberia. Acta trop. (Basel) 40, 271–295 (1983).
- Scrimshaw N. S., Taylor C. E., Gordon J. E.: Interactions of nutrition and infection. Wld Hlth Org. Org. Monogr. Ser. No. 57 (1968).
- Sommer A., Tarwotjo I., Hussaini G., Susanto D., Soegiharto T.: Incidence, prevalence and scale of blinding malnutrition. Lancet *1981/*1, 1407–1408.
- Speiser F.: Serodiagnosis of tissue dwelling parasites: application of a multi-antigen enzyme-linked immunosorbent assay (ELISA) for screening. Ann. Soc. belge Méd. trop. 62, 103–120 (1982).
- Stürchler D., Stahlel E., Saladin K., Saladin B.: Intestinal parasitoses in eight Liberian settlements: prevalences and community anthelminthic chemotherapy. Tropenmed. Parasit. *31*, 87–93 (1980).
- Stürchler D., Wyss F., Hanck A.: Retinol, onchocerciasis and *Onchocerca volvulus*. Trans. roy. Soc. trop. Med. Hyg. *75*, 617 (1981).
- World Health Organization: Vitamin A deficiency and xerophthalmia. WHO Techn. Rep. Ser. No. 590 (1976).
- World Health Organization: Control of vitamin A deficiency and xerophthalmia. WHO Techn. Rep. Ser. No. 672 (1982).
- Zile M. H., Bunge E. C., De Luca H. F.: On the physiological basis of vitamin A-stimulated growth. J. Nutr. *109*, 1787–1796 (1979).

18 Acta Tropica 269

