Zeitschrift: Acta Tropica

Herausgeber: Schweizerisches Tropeninstitut (Basel)

Band: 41 (1984)

Heft: 2

Artikel: Comparative study of the susceptibility to infection with "Trypanosoma

simiae" of "Glossina morsitans" and "G. tachinoides"

Autor: Agu, W.E.

DOI: https://doi.org/10.5169/seals-313287

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Nigerian Institute for Trypanosomiasis Research, Vom, Plateau State, Nigeria

Comparative study of the susceptibility to infection with *Trypanosoma simiae* of *Glossina morsitans* and *G. tachinoides*

W. E. AGU

Summary

The susceptibility to infection with *Trypanosoma simiae* of *Glossina morsitans* and *G. tachinoides* was compared. A total of 592 *G. tachinoides* and 348 *G. morsitans* were used in trying to transmit *T. simiae* to pigs. *G. morsitans* were very good at transmitting *T. simiae* infection to pigs while *G. tachinoides* were very poor. The epidemiological importance of the results is discussed.

Key words: *Trypanosoma simiae; Glossina morsitans; G. tachinoides;* pigs; trypanosomes; parasitaemia; transmitting; epidemiology.

Introduction

The distribution of different *Glossina* species in Nigeria is well known (Davis, 1977). However, evidence of differences in susceptibility to trypanosome infections between the species is scanty in Nigeria. Harley and Wilson (1968) and Harley (1971) have shown marked differences in infection rates between species of *Glossina* infected with the same species of trypanosome. There is indication that genetic and/or environmental factors are involved in controlling fly susceptibility (Maudlin, 1982).

The objective of this study was to compare the susceptibility to infection with *Trypanosoma simiae* of *Glossina morsitans* and *G. tachinoides*.

Materials and Methods

Animals. 6 large-white pigs were used for the study. They were housed in concrete pens and fed on concentrates and vegetables. Salt licks were also provided.

Correspondence: Dr. W. E. Agu, Nigerian Institute for Trypanosomiasis Research, P.M.B. 3, Vom, Plateau State, Nigeria

Tsetse flies. 592 G. tachinoides were used. 141 were wild flies caught at Yankari, Bauchi State, Nigeria; 451 were hatched in the laboratory from pupae collected also at Yankari. 348 G. morsitans were used, consisting of 320 wild flies caught at Yankari and 28 flies hatched from pupae collected at the same place.

Experimental procedure

The experiment was divided into two parts. 141 wild *G. tachinoides* were fed on pigs Nos. 10 and 14 for 30 days. Blood films were examined daily to determine if the pigs had become infected. 320 wild *G. morsitans* were fed on pig No. 4 and similar daily examinations made.

One pig was inoculated intramuscularly with Trypanosoma simiae which had been stored in liquid nitrogen. When the pig showed a heavy infection of T. simiae (40 trypanosomes per microscope field in a wet preparation, \times 10 eye-piece and \times 40 objective), 451 newly emerged (one day old) G. tachinoides and 28 newly emerged (one day old) G. morsitans were fed on the pig twice daily (morning and evening) for 3 days. The G. tachinoides were then fed on pig No. 16 for 27 days while the G. morsitans were fed on pig No. YR for 17 days. Roberts (1971) has shown that cyclical development of T. simiae in tsetse flies takes between 16 and 25 days. Blood films were examined daily to determine if the pigs had become infected.

Results

No trypanosome was seen in the peripheral blood of pigs Nos. 10 and 14, fed on by wild *G. tachinoides*, even after over 50 days of examining their blood each day.

The results of the *G. morsitans* experiment were different. On the 6th day of feeding wild *G. morsitans* on pig No. 4, *T. simiae* identified morphologically and by rodent inoculation (Hoare, 1936) appeared in the peripheral blood of the pig. The pig died 13 days later after showing heavy parasitaemia.

In the second part of the experiment, no trypanosome was seen in pig No. 16 fed on by *G. tachinoides* while *T. simiae* parasites were demonstrated in the peripheral blood of pig No. YR, fed on by *G. morsitans*. The parasitaemia commenced 21 days after the first feeding by the flies. The pig died on the 4th day after parasitaemia became patent.

Discussion

This work agrees with that of others (Desowitz and Watson, 1953; Stephen and Gray, 1960; Isoun, 1968) in showing that *G. morsitans* is an efficient transmitter of *T. simiae* in pigs. The literature is scanty on experimental transmission of *T. simiae* in pigs by *G. tachinoides*. The present work shows that *G. tachinoides* is not a suitable vector of *T. simiae* in pigs. Baldry (1964), in a survey of tsetse flies at Nsukka, Nigeria noted that *G. tachinoides* was very prevalent and showed a marked preference for the local pig as a host. His dissection studies on the flies showed 6.6% infection rate with *T. vivax*, *T. congolense* and *T. brucei*. Killick-Kendrick and Godfrey (1963) examined pigs in close contact with *G. tachinoides* at Nsukka, Nigeria and found that 86% of the 35 pigs examined were infected with *T. congolense* and/or *T. brucei*. The authors did not identify

T. simiae in the pigs, however, commented that some of the T. congolense infections identified in the pig survey could be due to T. simiae of low virulence. Janssen and Wijers (1974) reported that the virulence of T. simiae in pigs depended on the species of tsetse transmitting the trypanosomes; G. brevipalpis transmitted virulent infections while G. pallidipes transmitted chronic infections. Work is continuing to find out if this is so with G. morsitans and G. tachinoides. Roberts (1971), however, showed that G. tachinoides could transmit T. simiae to pigs though the infection rate in G. morsitans was higher than in G. tachinoides. He found by dissection that infection rates of up to 6% and 8.6% may be obtained in G. tachinoides and G. morsitans, respectively.

No explanation can so far be given for the poor vectorial capacity of G. tachinoides for T. simiae in pigs demonstrated in the present work. Harley and Wilson (1968) showed that G. fuscipes was a poor vector for T. congolense when compared with G. pallidipes. They thought that the reason for the difference was the frequent feeding of G. fuscipes on hosts such as man, reptiles and birds, which were not susceptible to T. congolense infection. Genetic factors have been found to control the susceptibility of vectors to infections (Wakelin, 1978; Maudlin, 1982).

My results are significant in the epidemiology of *T. simiae* in pigs, particularly in areas of poor traditional pig husbandry where the pigs roam and scavenge in the compound. Obviously, in areas where *G. morsitans* is prevalent, there would be very limited success in establishing piggeries, except in fly-proof buildings. In areas where *G. tachinoides* are found, there would be a good chance of a successful out-door pig farm.

Acknowledgments

I wish to thank Mr. M. M. Baah, Mr. D. Bayei, Mallam Adamu Shittu and Mr. Ayo Adesiyan for their invaluable technical assistance. I also thank the Director of the Nigerian Institute for Trypanosomiasis Research, Alhaji Y. Magaji for permission to publish.

- Baldry D. A. T.: Observation on a close association between *Glossina tachinoides* and domestic pigs near Nsukka, Eastern Nigeria. 11. Ecology and trypanosome infection rates in *G. tachinoides*. Ann. trop. Med. Parasit. *58*, 32–44 (1964).
- Davis H.: Tsetse flies in Nigeria. 3rd edition. Oxford University Press, Ibadan 1977.
- Desowitz R. S., Watson H. J. C.: The maintenance of a strain of *Trypanosoma simiae* in rabbits: the effect on splenectomy on the course of infection. Ann. trop. Med. Parasit. 47, 324–334 (1953).
- Harley J. M. B.: Comparison of the susceptibility to infection with *Trypanosoma rhodesiense* of *Glossina pallidipes*, *G. morsitans*, *G. fuscipes* and *G. brevipalpis*. Ann. trop. Med. Parasit. 65, 185–189 (1971).
- Harley J. M. B., Wilson A. J.: Comparison between *Glossina morsitans*, *G. pallidipes* and *G. fuscipes* as vectors of trypanosomes of the *Trypanosoma congolense* group: the proportions infected experimentally and the numbers of infective organisms extruded during feeding. Ann. trop. Med. Parasit. *62*, 178–187 (1968).

- Hoare C. A.: Morphological and taxonomic studies on mammalian trypanosomes. 11. *Trypanosoma simiae* and acute porcine trypanosomiasis in tropical Africa. Trans. roy. Soc. trop. Med. Hyg. 29, 619–645 (1936).
- Isoun T. T.: The pathology of *Trypanosoma simiae* infection in pigs. Ann. trop. Med. Parasit. 62, 188–192 (1968).
- Janssen J. A. H. A., Wijers D. J. B.: *Trypanosoma simiae* at the Kenya coast. A correlation between virulence and the transmitting species of *Glossina*. Ann. trop. Med. Parasit. 68, 5–19 (1974).
- Killick-Kendrick R., Godfrey D. G.: Observations on a close association between *Glossina tachi-noides* and domestic pigs near Nsukka, Eastern Nigeria. 1. *Trypanosoma congolense* and *T. brucei* infections in the pigs. Ann. trop. Med. Parasit. 57, 225–231 (1963).
- Maudlin I.: Inheritance of susceptibility to *Trypanosoma congolense* infection in *Glossina morsitans*. Ann. trop. Med. Parasit. 76, 225–227 (1982).
- Roberts C. J.: The lack of infectivity to cattle of a strain of *Trypanosoma simiae* transmitted by *Glossina morsitans* and *G. tachinoides*. Ann. trop. Med. Parasit. 65, 319–326 (1971).
- Stephen L. E., Gray A. R.: Suramin complexes. VI. The prophylactic activity of antrycide-suramin complex and antrycide chloride against *Trypanosoma simiae* in pigs. Ann. trop. Med. Parasit. *54*, 493–507 (1960).
- Wakelin D.: Genetic control of suceptibility and resistance to parasitic infection. Advanc. Parasit. *16*, 219–308 (1978).