

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 42 (1985)
Heft: 2

Artikel: A new defined medium for cultivating "Leishmania" promastigotes
Autor: Melo, Norma Maria / Peixoto de Azevedo, H. / Roitman, I.
DOI: <https://doi.org/10.5169/seals-313462>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

¹ Departamento de Parasitologia, ICB/Universidade Federal de Minas Gerais,
30000 Belo Horizonte MG, Brasil

² Departamento de Biologia Celular, Universidade de Brasília, 70910 Brasília DF, Brasil

A new defined medium for cultivating *Leishmania* promastigotes*

NORMA MARIA MELO¹, H. PEIXOTO DE AZEVEDO², I. ROITMAN², W. MAYRINK¹

Summary

A chemically defined medium to grow the promastigotes of 19 stocks of *Leishmania* is described. The medium was developed by making qualitative and quantitative modifications of the medium AR-103 devised for *Trypanosoma cruzi*. The new medium, designated as MD-29, can be used in studies on the nutritional requirements of promastigotes.

Key words: *Leishmania*; cultivation of *Leishmania* promastigotes; chemically defined culture medium.

Introduction

The complexity of the media used to cultivate members of the genus *Leishmania* has complicated the feasibility of comparative studies on nutritional requirements, metabolism and other characteristics of these organisms, especially those isolated from mammals. The cultivation of *Leishmania* in a defined medium opens up new possibilities to determine biochemical and immunological characteristics.

A chemically defined medium (AR-130) for the cultivation of *Trypanosoma cruzi* was developed (Azevedo and Roitman, 1983); quantitative and qualitative (ATP and AMP were omitted and iron nitrate was added) modification of this medium led to the design of a chemically defined medium, herein designated as MD-29 in which several strains of *Leishmania* can be cultivated.

* This work is part of a Thesis for the Doctor Degree in Parasitology, by the senior author.

Correspondence: Prof. Isaac Roitman, Departamento de Biologia Celular, Universidade de Brasília, 70910 Brasília DF, Brazil

Table 1. Defined medium MD-29 for *Leishmania* cultivation

Components	mg/l	Components	mg/l
Fructose	2000	D Choline	12
A Na ₃ PO ₄ ·12H ₂ O	5000	E L-Alanine	285
β-glycerophosphate Na salt ..	20000	L-Arginine	550
Sodium acetate 3H ₂ O	540	L-Aspartic acid	100
NaCl	4000	L-Cysteine HCl	100
KCl	400	L-Glutamic acid	1045
EDTA Na salt	80	L-Glycine	350
Trisodium citrate 2H ₂ O ...	600	L-Histidine	350
FeNO ₃ ·9H ₂ O	0.0000005	L-Isoleucine	260
Sodium succinate 6H ₂ O ...	270	L-Leucine	760
B Adenosine	20	L-Lysine HCl	750
Adenine HCl	50	L-Methionine	175
Guanosine	20	L-Phenylalanine	373
Guanine HCl	1.5	L-Proline	780
Hypoxantine	1.5	L-Hydroxyproline	50
Xantine	1.5	L-Serine	285
C Folic acid	30	L-Threonine	270
D-α tocopherol succinate ...	4	L-Tryptophan	140
DL-α lipoic acid	0.4	L-Tyrosine etil ester	360
Nicotinic acid	0.125	L-Valine	485
Nicotinamide	10	F Haemin	1.0
D-Biotin	10	Double distilled H ₂ O to make up to	1000 ml
Riboflavin	1	pH adjusted to 7.5 with 1n HCl	
Inositol	2		
D-Ca pantothenate	10		
Pyridoxine HCl	0.125		
Pyridoxal	10		
Thiamine	10		
Ascorbic acid	0.25		

Notes: All ingredients were obtained from Sigma Chemical Company. A, B, C, and E were added from a dry mix stored at -4°C. D was added from a frozen stock solution. The medium was Seitz filtered before an autoclaved solution of haemin (F) was added aseptically. Fructose was added to the completely prepared medium.

Material and Methods

The composition of medium MD-29 is given in Table 1. Nineteen strains of *Leishmania* have been cultivated in medium MD-29. Cultures in the chemically defined medium were initiated with amastigotes from cutaneous lesions (BH6, BH121, BH32, BH132, BH216, BH224, LV78, PH8) and the spleen of experimentally infected hamster (BH46), and from promastigotes grown at 23°C in LIT (liver infusion-tryptose) medium (Camargo, 1964) (BH46, BH121, BH32, BH132, BH216, BH224, LV78, PH8, BH49, LV9, LV4, LV414, LRC-L7, LRC-L121, LRC-L134, LRC-L259, LUMP1482 and LUMP1701).

A fine pointed Pasteur pipette was used to aspirate material from infected tissue. In order to isolate parasites from cutaneous lesions, the animal was anesthetized with chloroform, hair on the

Table 2. *Leishmania* strains cultivated in defined medium MD-29

Designation and source	Species	Place of isolation	Number of passages	Mean growth* ($\times 10^7/\text{ml}$)
BH6 (Man)	<i>L. mexicana mexicana</i>	Brazil	187	4.0
BH46 (Man)	<i>L. donovani</i>	Brazil	208	4.5
BH49 (Man)	<i>L. mexicana</i>	Brazil	152	3.2
BH121 (Man)	<i>L. mexicana</i>	Brazil	218	4.3
LV9 (Man)	<i>L. donovani</i>	Ethiopia	106	3.6
LV4 (Nyctomyssumichrasti)	<i>L. mexicana mexicana</i>	Belize	106	6.0
LV78 (<i>Proechimys</i> sp.)	<i>L. mexicana amazonensis</i>	Brazil	106	4.0
LV414 (<i>Tarentolae</i> sp.)	<i>L. tarentolae</i>	Africa	106	5.3
PH8 (<i>Lutzomyiaflaviscutellata</i>)	<i>L. mexicana amazonensis</i>	Brazil	126	3.5
BH32 (<i>Akodon cursor</i>)	<i>Leishmania</i> sp.	Brazil	59	6.1
BH132 (<i>Proechimys dimidiatus</i>)	<i>Leishmania</i> sp.	Brazil	55	3.7
BH216 (<i>Akodon cursor</i>)	<i>Leishmania</i> sp.	Brazil	55	3.3
BH224 (<i>Proechimys dimidiatus</i>)	<i>Leishmania</i> sp.	Brazil	55	3.0
LRC-L7 (Man)	<i>L. tropica</i>	India	50	3.2
LRC-L121 (<i>Agama stellio</i>)	<i>L. agamiae</i>	Israel	52	3.5
LRC-L134 (Man)	<i>L. aethiopica</i>	Ethiopia	50	3.1
LRC-L259 (Man)	<i>L. mexicana amazonensis</i>	Brazil	53	3.4
LUMP1482 (<i>Coendu rothschildi</i>)	<i>L. hertigi hertigi</i>	Panama	51	3.1
LUMP1701 (<i>Coendu</i> sp.)	<i>L. hertigi deanei</i>	Brazil	51	3.2

* Final cell density after 9 days incubation at 25°C

BH = Belo Horizonte

LRC = Leishmaniasis Research Centre – Jerusalem

LV = Liverpool

LUMP = London University Medical Protozoology

PH8 = Obtained from Ralph Lainson – Instituto Evandro Chagas – Belém

border of the lesion was removed, and the skin surface was sterilized with merthiolate before inserting the pipette into the margin of the lesion. When parasites were to be isolated from the spleen, the animal was killed and the abdomen opened to expose the organ. When promastigotes were used to initiate cultures, 1 ml was withdrawn from 48-h LIT cultures containing $2-3 \times 10^7$ flagellates/ml, and added to the defined medium in 20 × 150 mm screwcap test tubes to give a final volume of 5 ml.

Subcultures were made at 9-day intervals by transferring 1 ml of the culture to 5 ml of the chemically defined medium. Cultures were incubated at 25°C and growth was assessed qualitatively and quantitatively by observations on the appearance and mobility of promastigotes and by assessing their numbers by means of a Coulter Counter, model ZB.

Results

A summary of the results obtained with 19 strains of *Leishmania* is given in Table 2. The defined medium supports the growth of strains isolated from both vertebrate and invertebrate host, with a yield of more than 3×10^7 promastigotes/ml in a 9-day period.

Discussion

Medium MD-29 can be used for mass produce promastigotes for analytical biochemical studies. In fact, the medium was used already for the production of promastigotes for studies on surface polysaccharides (Fiorini et al., 1981), nutritional requirements (Melo et al., 1982) and other important aspects of in vitro development as the maintenance of the pathogenicity of *Leishmania* (Melo, 1983).

As far as we know, this is the first time that strains of *L. tropica*, *L. aethiopica*, *L. hertigi*, *L. mexicana* and New World *L. donovani* have been cultivated in a defined medium. The defined media previously described (Trager, 1957; Steiger and Steiger, 1976, 1977; Berens and Marr, 1978) were used to culture only *L. tarentolae*, Old World *L. donovani* and *L. braziliensis*. Medium RE III of Steiger and Steiger (1977) did not support the development of *L. tropica*, *L. mexicana texana* or *L. hertigi* (Fish et al., 1978).

Leishmania species seem to be fastidious in their nutritional requirements since many strains of *Leishmania* are not able to grow in a defined medium – containing 10 amino acids, 7 vitamins, purine and hemin – (Roitman et al., 1972), that supports the growth of many lower trypanosomatids (unpublished results).

The availability of a defined medium contributes to an improved understanding of the biology of *Leishmania*. It helps to develop rational methods of chemotherapy; provides a source of cells free of extraneous antigenic material; and may lead to a better knowledge of the genetics of the parasites. The knowledge of the nutritional requirements by using the present medium will allow to design a minimal medium for *Leishmania*. Studies on antigens for immuno-diagnosis derived from promastigotes, cultured in our defined medium, are in progress.

Acknowledgments

This investigation received financial support from the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases and Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq, Brasil.

- Azevedo H. P., Roitman I.: Cultivation of *Trypanosoma cruzi* in defined medium. In: Genes and antigens of parasites, ed. by C. Morel, p. 29–36. WHO Special Publication (1983).
- Berens R. L., Marr J.: An easily prepared defined medium for cultivation of *Leishmania donovani* promastigote. J. Parasit. 64, 160 (1978).
- Camargo E. P.: Growth and differentiation of *Trypanosoma cruzi*. I. Origin of metacyclic trypanosomes in liquid medium. Rev. Inst. Med. trop. S. Paulo 6, 93–100 (1964).
- Fiorini J. E., Azevedo H. P., Angluster J., De Souza W., Roitman I., Alviano C. S.: Poissacarídeos de superfície em *Leishmania mexicana* e *Leishmania donovani*. 8º Reunião Anual Pesquisa Básica Doença de Chagas, Caxambu, Brasil, Abstr. 110 (1981).
- Fish W. R., Holz G. G., Beach D. H.: Cultures of trypanosomatids. J. Parasit. 64, 546–547 (1978).

- Melo M. N., Azevedo H. P., Roitman I., Mayrink W.: Requerimento de aminoácidos e vitaminas de duas cepas de *Leishmania*. 7º Congr. Brasil. Parasitologia, Porto Alegre, Brasil, 113–114 (1982).
- Melo M. N.: Cultivo de *Leishmania* em meio definido: estudos de seus requerimentos nutricionais. Ph. D. Thesis, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, 113 p. (1983).
- Roitman C., Roitman I., Azevedo H. P.: Growth of an insect trypanosomatid at 37°C in a defined medium. J. Protozool. 19, 346–349 (1982).
- Steiger R. F., Steiger E.: A defined medium for cultivating *Leishmania donovani* and *Leishmania braziliensis*. J. Parasit. 62, 1010–1011 (1976).
- Steiger R. F., Steiger E.: Cultivation of *Leishmania donovani* and *Leishmania braziliensis* in defined media: nutritional requirements. J. Protozool. 24, 437–441 (1977).
- Trager W.: Nutrition of a hemoflagellate (*Leishmania tarentolae*) having a inter-changeable requirement of choline or pyridoxal. J. Protozool. 4, 269–276 (1957).

