

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 42 (1985)
Heft: 3

Artikel: Survival period of field isolates of "Trypanosoma vivax" in refrigerated blood : short communication
Autor: Ekwuruke, J.O. / Ikede, B.O. / Opasina, B.A.
DOI: <https://doi.org/10.5169/seals-313478>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

¹ International Livestock Centre for Africa, Humid Zone Programme, P.M.B. 5320, Ibadan

² Department of Veterinary Pathology, University of Ibadan, Ibadan, Nigeria

Survival period of field isolates of *Trypanosoma vivax* in refrigerated blood

Short communication

J. O. EKWURUKE¹, B. O. IKEDE², B. A. OPASINA¹

The simplest and most reliable parasitological technique for the diagnosis of *T. vivax* and *T. congolense* infections in domestic animals is to centrifuge unclotted blood in capillary tubes and look for the parasites packed between the buffy coat and the red cell layer (Murray et al., 1977). The content of this section of each capillary tube is expressed onto a slide and examined using phase contrast or darkground microscopy. The level of parasitaemia can be determined by a scoring system (Paris et al., 1982). However, in field situations, there can be considerable time lag between blood collection and examination unless a portable generator is available to provide electricity. The usual practice, therefore, is for blood samples to be cooled with ice and transported to the laboratory for examination. Paris and others working with cattle recommended that samples should be examined within 12 h if the parasites are to survive in reasonable numbers (Paris et al., 1982; Murray et al., 1983). Our studies indicate that the time could be much longer without any appreciable change in the trypanosome score, at least when examining small ruminants.

During routine screening of animals at Fasola in southwest Nigeria for trypanosomiasis 9 sheep and 9 goats naturally infected with *T. vivax* were identified. 5 ml of blood was obtained from each animal using EDTA vacutainer tubes. Samples were immediately cooled in ice and transported over 70 km to the laboratory in Ibadan where they were refrigerated at 4°C. Two, 16 and 24 h after collection, trypanosome scores were determined as described by Paris et al. (1982) and shown on Table 1. Parasites were found to be viable beyond 24 h although there was some reduction in the trypanosome scores (Table 2). Another set of infected blood samples from 3 sheep and 4 goats was then examined as before but at 6 hourly intervals until the 68th hour after collection.

The results are shown in Table 3. It was found that the mean score remained constant for 20 h in sheep and 26 h in goats infected with *T. vivax*. Even at the 50th hour the mean score was still above 50% of the initial value with both species of animals. The results thus indicate that, with small ruminants infected with *T. vivax*, reliable trypanosome scores can be obtained up to 24 h after sample collection so long as the samples are properly refrigerated. Qualitatively, infection could be detected up to the third day after collection; trypanosomes survived better in goat blood than sheep blood. Similar studies with other species of trypanosomes and livestock would be worthwhile.

Murray M., Murray P. K., McIntyre W. I. M.: An improved parasitological technique for the diagnosis of African trypanosomiasis. *Trans. roy. Soc. trop. Med. Hyg.* 71, 325–326 (1977).

Murray M., Trail J. C. M., Turner D. A., Wissocq Y.: Livestock productivity and trypanotolerance. Network training manual, p. 6–28. ILCA, Addis Ababa 1983.

Correspondence: Prof. B. O. Ikede, Department of Veterinary Pathology, University of Ibadan, Ibadan, Nigeria

Paris J., Murray M., McOdimba F.: A comparative evaluation of the parasitology techniques currently available for the diagnosis of African trypanosomiasis in cattle. *Acta trop. (Basel)* 39, 307–316 (1982).

Table 1. Darkground/phase contrast buffy coat parasitaemia scoring system (Paris et al., 1982)

Score	Trypanosomes/field*	Estimated parasitaemia trypanosomes/ml
6+	Swarming >100	$>5 \times 10^6$
5+	>10	$>5 \times 10^5$
4+	1–10	10^4 – 5×10^5
3+	1 per 2 fields to 1 per 10 fields	5×10^3 – 5×10^4
2+	1–10 per preparation	10^3 – 10^4
1+	1 per preparation	10^2 – 10^3

* magnification = 250

Table 2. Mean trypanosome scores with time in refrigerated blood samples from sheep and goats naturally infected with *T. vivax* (Experiment 1)

	Hours after blood collection		
	2	16	24
Sheep (n = 9)	3.8 ± 1.2	3.1 ± 1.2	2.2 ± 1.1
Goat (n = 9)	4.7 ± 1.1	3.7 ± 1.4	2.7 ± 1.3

Table 3. Mean trypanosome scores with time in refrigerated blood samples from sheep and goats naturally infected with *T. vivax* (Experiment 2)

Hours after blood collection	Sheep (mean \pm SD) (n = 3)	Goat (mean \pm SD) (n = 4)
2	4.3 ± 1.9	4.5 ± 1.1
8	4.3 ± 1.9	4.5 ± 1.1
14	4.3 ± 1.3	4.5 ± 1.1
20	4.3 ± 1.3	4.5 ± 1.1
26	3.6 ± 1.3	4.5 ± 1.1
32	3.6 ± 1.0	3.5 ± 1.1
38	3.3 ± 1.0	3.5 ± 1.1
44	3.3 ± 1.0	3.5 ± 1.1
50	2.3 ± 1.0	3.3 ± 1.5
56	2.9 ± 0.8	3.0 ± 1.2
62	1.3 ± 0.4	2.5 ± 1.1
68	1.0 ± 0.4	2.5 ± 1.1

SD = Standard deviation