

Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	43 (1986)
Heft:	1
Artikel:	Case report : non-ulcerative amebiasis of rectum : short communication
Autor:	George, G.H. / Baird, J.K. / Thorpe, R.G.
DOI:	https://doi.org/10.5169/seals-313616

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Department of Infectious and Parasitic Disease Pathology, Armed Forces Institute of Pathology,
Washington, D.C., USA
Karawa Mission Hospital, Ubangi, Zaire

Case report: Non-ulcerative amebiasis of rectum¹

Short communication

G. H. GEORGE², J. K. BAIRD³, R. G. THORPE⁴, D. H. CONNOR

Studies of man and animals indicate that lysis of intestinal epithelium precedes movement of *Entamoeba histolytica* into the lamina propria (Prathap and Gilman, 1970; Pittman et al., 1973; Takeuchi and Phillips, 1975; Takeuchi et al., 1977; Chadee and Meerovitch, 1985). Here we describe trophozoites of *E. histolytica* in rectal tissue without detectable lytic or other injury to any of the tissues, including intestinal epithelium, lamina propria, and muscularis mucosae.

A 69-year-old black Zairian woman living in the Ubangi territory was imprisoned for sorcery and died suddenly after 2 days of punitive starvation. Autopsy at Karawa Mission Hospital revealed no significant gross lesions. Microscopically, there was a myocarditis of undetermined cause, but this was mild and not thought to be the cause of death. Sections of rectum showed minimal autolysis characterized by separation of epithelial cells from the basement membrane. Trophozoites of *E. histolytica* were in most sections of the rectum studied. They measured 20 µm in maximum diameter and contained a nucleus with a prominent central karyosome. Their cytoplasm was amphophilic, granular, and sometimes vacuolated (Fig. 1). One trophozoite contained a red blood cell. The trophozoites occurred singly and in clusters of up to 4 or


¹ The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Navy, the Department of the Army, or the Department of Defense.

² Dr. George was on a research fellowship to the AFIP.


³ The military status of J. K. Baird is LTJG, MSC, USNR.

⁴ Karawa Mission Hospital, Ubangi, Zaire.

Correspondence: Dr. Daniel H. Connor, Chairman, Department of Infectious and Parasitic Disease Pathology, Armed Forces Institute of Pathology, Washington, D.C. 20306-6000, USA



1



2

Fig. 1. Three trophozoites (arrows) of *E. histolytica* in lamina propria and muscularis mucosae of rectum. One lies between epithelium and basement membrane and appears to be constricting the adjacent epithelial cells. Hematoxylin and eosin, $\times 415$, AFIP MIS # 85-7403.

Fig. 2. A cluster of 7 *E. histolytica* trophozoites in the space between the basement membrane and epithelium. Adjacent tissue is normal or possibly slightly inflamed, but the inflammation cannot be related to the invading trophozoites. Hematoxylin and eosin, $\times 165$, AFIP MIS # 85-7398.

more (Fig. 2). They were in the lamina propria and muscularis mucosae (Fig. 1). Many were in the space between the epithelium and basement membrane (Figs. 1 and 2). The characteristic lytic changes of colonic amebiasis were not present. Elsewhere in the rectum there was a slight diffuse infiltrate of the lamina propria by plasma cells, lymphocytes, and eosinophils, but these were no more than usually is seen in colons of Africans, and none of the inflammatory cells was concentrated around or near the invading trophozoites. This mild inflammation therefore was not a response to the invading trophozoites. Addi-

tional sections, taken routinely, of colon, ileum, liver, lung, and pericardium revealed no trophozoites or necrotizing lesions that suggested amebiasis.

The source of the trophozoites remains unclear. It seems likely, however, that they migrated to and invaded the rectum – either from the lumen of the large intestine where they had not caused lesions or from inapparent lesions in the colon that we did not detect. But in the rectum there is invasion of intestinal mucosa and lamina propria without lytic activity and without inflammation and ulceration. We think this is because the trophozoites penetrated the rectal mucosa post mortem. If so, the absence of lysis and necrosis could be explained by the lack of host response. Takeuchi and Phillips (1975) suggested that lysis of host tissue may be a consequence of infiltrating polymorphonuclear leukocytes. They showed experimentally that invading trophozoites penetrate epithelial tissue between cell junctions and that necrosis followed contact between amebae and infiltrating neutrophils. Direct contact between trophozoites and neutrophils resulted in immediate lysis of the neutrophils with spillage of hydrolytic enzymes. The invasion of trophozoites in our present case appears to support the view that lysis requires a reaction by the host. This case also reveals that trophozoites of *E. histolytica* are capable of non-lytic penetration of intestinal epithelium. In summary, we believe the absence of necrosis in the rectum of this corpse is a consequence of the absence of a neutrophilic response.

Chadee K., Meerovitch E.: *Entamoeba histolytica*: early progressive pathology in the cecum of the gerbil (*Meriones unguiculatus*). Amer. J. trop. Med. Hyg. 34, 283–291 (1985).

Pittman F. E., El-Hashimi W. K., Pittman J. C.: Studies of human amebiasis. II. Light and electron microscopic observations of colonic mucosa and exudate in acute amebic colitis. Gastroenterology 65, 588–603 (1973).

Prathap K., Gilman R.: The histopathology of acute intestinal amebiasis. A rectal biopsy study. Amer. J. Path. 60, 229–245 (1970).

Takeuchi A., Phillips B. R.: Electron microscope studies of experimental *Entamoeba histolytica* infection in the guinea pig. I. Penetration of the intestinal epithelium by trophozoites. Amer. J. trop. Med. Hyg. 24, 34–48 (1975).

Takeuchi A., Jervis H. R., Phillips B. R.: Electron microscope studies of experimental *Entamoeba histolytica* infection of the guinea pig. III: Histolysis of the cecum. Virchows Arch. B Cell Path. 24, 263–277 (1977).