**Zeitschrift:** Acta Tropica

**Herausgeber:** Schweizerisches Tropeninstitut (Basel)

**Band:** 44 (1987)

Heft: 1

**Artikel:** Stage-specific homocytotropic antibody response of "Mastomys

natalensis" to "Dipetalonema viteae" infection: short communication

**Autor:** Singh, D.P. / Misra, Shailja / Chatterjee, R.K.

**DOI:** https://doi.org/10.5169/seals-313825

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 19.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Division of Parasitology, Central Drug Research Institute, Lucknow, India

# Stage-specific homocytotropic antibody response of *Mastomys natalensis* to *Dipetalonema viteae* infection<sup>1</sup>

### Short communication

## D. P. Singh, Shailja Misra, R. K. Chatterjee, A. B. Sen

Attempt was made in the present study to investigate whether filaria-specific homocytotropic antibody (HcAb) level of host was uniform throughout or related to prevalence of specific life-stage of the parasite, using *Dipetalonema viteae* in *Mastomys natalensis* as a working model and assaying antibody by passive cutaneous anaphylaxis (PCA).

Male mastomys (6 weeks old, GRA Giessen strain) were infected by inoculating 50 infective larvae (L<sub>3</sub>) of *D. viteae* isolated from infected ticks, *Ornithodoros moubata* (Worms et al., 1961). Animals were also infected by implantation of adult male and female worms (5–8 numbers) and microfilariae (0.2 million) subcutaneously. Soluble somatic antigen was prepared separately from adult male and female worms, L<sub>3</sub> and microfilariae (mf) by the method of Singh et al. (1985). Protein content of the preparations was estimated by the technique of Lowry et al. (1951).

Microfilariaemia/5 mm<sup>3</sup> of tail blood of animals was assessed at weekly or fortnightly interval starting from day 60 post L<sub>3</sub> exposure or days 4 and 7 postimplantation. Blood for serum was collected from retro-orbital plexus of animal on days 20 and 30 of infection and thereafter at monthly interval along with matched uninfected controls.

PCA test was performed in homologous, healthy recipients (12 weeks old) following the method of Soulsby et al. (1977). After initial evaluation, a quantum of 250  $\mu$ g of antigen protein was found optimal for positive PCA reaction. PCA test was also performed after heating sera samples at 56°C for 1 h to exclude the possibility of interference of IgG type of reagin in the test.

The study showed the presence of HcAb against all the three important life stages e.g. mf,  $L_3$  and adults of D. viteae (Fig. 1).  $L_3$ -specific HcAb did not last longer possibly due to short persistence of this life-stage in the host. The peak level was detected within 40 days and total disappearance occurred before day 150 post exposure (p.e.). HcAb response of host to adult stage was detected on day 20 and persisted for longer period i.e. up to day 240 p.e. The persistence of antibody to adult parasites even after their death could possibly be due to cross-reactions with other developing forms. High adult-specific HcAb titre as detected on day 60 coinciding with the appearance of mf in circulation could also be due to enhanced common antigenic stimuli from mf and adults. However, this does not explain why adult-specific antibody disappeared on day 240 p.e. when mf still persisted. The plausible reason could be that major antigenic stimuli (excretory and secretory product) specific to adults slowly disappeared with the death of worms and the quantum of 250  $\mu$ g of adult antigen in PCA test was insufficient to detect cross reacting mf antibody during that period. Almost similar pattern of

Correspondence: Dr. R. K. Chatterjee, Division of Parasitology, Central Drug Research Institute, P.O. Box 173, Lucknow 226001, India

<sup>&</sup>lt;sup>1</sup> C.D.R.I. Communication No. 3704




Fig. 1. Stage-specific homocytotropic antibody response of Mastomys during the course of D. viteae infection. Fig. 2. Homocytotropic antibody of *Mastomys* to implanted stage-specific infection.

HcAb response with adult antigen was observed earlier by Benjamin and Soulsby (1976) in mastomys infected with Brugia pahangi. In spite of very long persistence of mf, anti-mf titre never reached to a very high level and low titre persisted up to the end of present observation period (i.e. 300 days) when amicrofilaraemia was set in. Apparently mf appear to be more antigenic and their sensitization effects last longer. Wederhof and Wenk (1975) reported stage-specific protection using live mf, though it was not clear from their study whether IgE played any role in that. However, Haque et al. (1981) reported IgE-mediated in vitro killing of D. viteae mf by macrophage-rich peritoneal exudate cells. Detection of mf-specific PCA titre as early as day 40 p.e. could be due to antigenic stimulus from uterine secretions of gravid female worms. Nevertheless, some cross-reactivity with other life-stages can not be ruled out. The nature of stage-specific HcAb response as observed with adult male, female or mf-implanted infection was by and large similar to L<sub>3</sub>-induced infection (Fig. 2). Apparently male worms were found to be more potent in inducing HcAb response than their female counterparts which might be due to earlier death of the latter sex (thus lacking E-S products).

Apart from eliciting hypersensitivity reaction, the role of filaria-specific IgE in acquired resistance against filarial infection remains unanswered. Though there are certain reports regarding its role in functional immunity to helminth parasites (Ogilvie et al., 1966; Sadun and Gore, 1970; Gusmao et al., 1981) its involvement in protective immunity in filariasis is yet to be conclusively proved. The present study shows that stage-specific HcAb develop during filarial infection but their association in eliminating any particular life-stage or infection as a whole is not clear. The homocytotropic antibody detected in the present study was heat labile.

```
Benjamin D. B., Soulsby E. J. L.: Amer. J. trop. Med. Hyg. 25, 266-272 (1976).
Gusmao R. D. A., Stanley A. M., Ottesen E. A.: Exp. Parasit. 52, 147-149 (1981).
Haque A., Ouaissi A., Joseph M., Capron M., Capron A.: J. Immunol. 127, 716-725 (1981).
Lowry O. H., Rosebrough A. C., Farr R. L., Randall R. J.: J. biol. Chem. 193, 265-275 (1951).
Ogilvie B. M., Smithers S. R., Terry R. J.: Nature (Lond.) 209, 1221–1223 (1966).
Sadun E. H., Gore R. W.: Exp. Parasit. 28, 435–449 (1970).
Singh D. P., Rathore S., Misra S., Chatterjee R. K., Ghatak S., Sen A. B.: Trop. Med. Parasit. 36, 21–24
  (1985).
Soulsby E. J. L., Zahner H., Weidner E., Lammler G.: Pathophysiol. Parasit. Infect. p. 149–159 (1977).
```

Wederhof P. H., Wenk P.: Parasitology 71, XIX-XX (1975).

Worms M. J., Terry R. J., Terry A.: J. Parasit. 47, 963–970 (1961).