**Zeitschrift:** Acta Tropica

**Herausgeber:** Schweizerisches Tropeninstitut (Basel)

**Band:** 44 (1987)

Heft: 4

**Artikel:** Competent metabolic utilization of hydrogen peroxide by trypanosomes

: short communication

**Autor:** Penketh, P.G. / Kennedy, W.P.K. / Patton, C.L.

**DOI:** https://doi.org/10.5169/seals-313876

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

## Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 24.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

## Competent metabolic utilization of hydrogen peroxide by trypanosomes

Short communication

P. G. Penketh<sup>1</sup>, W. P. K. Kennedy<sup>2</sup>, C. L. Patton<sup>1</sup>, A. C. Sartorelli<sup>1</sup>

Many trypanosomatids have been reported to be deficient in their ability to metabolize  $H_2O_2$  and be particularly sensitive to  $H_2O_2$  oxidant stress (Docampo and Moreno, 1984) due to their lack of the glutathione peroxidase and catalase. Recently, however, it has been demonstrated that *Trypanosoma brucei* possesses a novel trypanothione-dependent enzymic system for metabolizing  $H_2O_2$  (Penketh, 1986; Penketh and Klein, 1986). This finding has led us to question the generally accepted idea that trypanosomes are particularly sensitive to  $H_2O_2$  oxidant stress.

T. brucei rhodesiense bloodstream trypomastigotes were purified as described by Lanham (1968). T. cruzi strain epimastigotes and trypomastigotes were prepared and purified as described by Widmer (1986) and Piras et al. (1982), respectively. Levels of parasitemia were determined using a Neubauer haemocytometer at a magnification of 200×. Erythrocyte morphology was examined in thin wet smears at magnifications of 200–1000×. The toxicity of H<sub>2</sub>O<sub>2</sub> to mice and to T. b. rhodesiense was determined as follows. Female CD-1 25–35 g mice were injected intravenously (i.v.) with glucose oxidase (GO) in 50 μl of phosphate buffered saline. The LD<sub>50</sub> dose of glucose oxidase was determined to be 3.8–4.5 mg/kg (133 units/mg; Sigma). When catalase (180 mg/kg, 11,000 units/mg; Sigma) was co-administered, mice tolerated doses of 45.0 mg/kg of GO with no deaths (0/4). In the absence of catalase, however, doses of 5.5 mg/kg were 100% lethal (8/8). This finding indicates that H<sub>2</sub>O<sub>2</sub> production is the major toxic product of GO treatment. Mice given an LD<sub>50</sub> dose of GO generally died within 24 h or they survived.

Mice were infected with T. b.  $rhodesiense~(2\times10^6~cells~i.p.)$  and the doubling time of the organisms was determined by following the increase in parasitemia. A mean doubling time of 6.8~h~(n=24, SD=0.6) was observed and all of the mice died approximately 3 days after infection with parasitemias of greater than  $10^9~cells/ml$ . In a further experiment 14 mice were infected with  $2\times10^6~parasites$  and after 24 h were injected with 3.8 mg/kg of GO. Parasitemia levels were then followed in these animals. Approximately 70% (9/14) of the animals survived the GO treatment. However, they all died approximately 3 days after infection with maximal parasitemias. A mean doubling time of 7.1 h (n = 9, SD = 1.3) was calculated. At all times the parasites were of normal appearance. Since both

Correspondence: Dr. P. G. Penketh, Department of Pharmacology, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06510, USA

<sup>&</sup>lt;sup>1</sup> MacArthur Center for Parasitology and Tropical Medicine, Yale University School of Medicine, New Haven, CT 06510, USA

<sup>&</sup>lt;sup>2</sup> Wolfson Molecular Biology Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1 E7HT, U.K.

trypanosomes (in acutely infected mice) and erythrocytes reside in the bloodstream, both cell types should be exposed to equivalent steady state  $H_2O_2$  concentrations. Although trypanosomes appeared to be unaffected by GO administration, erythrocytes were adversely affected unless catalase, 180 mg/kg, was co-administered. Within 12 h of GO treatment, virtually all of the erythrocytes were of echinocyte I–IV morphology indicating major cytoskeletal changes had occurred within these cells.  $H_2O_2$  has been previously shown to cause cytoskeletal damage (Ragu et al., 1986) and inhibit cellular volume-regulation (Rosenberg and Mathews, 1973). This suggests that the oxidant defense mechanisms of the erythrocytes had been compromised even though these cells are not normally considered to be deficient in defenses against  $H_2O_2$ .

The quantities of  $H_2O_2$  metabolized in 1 min by T. b. rhodesiense trypomastigotes (5×10 $^7$  cells/ml), T. cruzi epimastigotes (2×10 $^7$  cells/ml) and trypomastigotes (2×10 $^7$  cells/ml), using an initial  $H_2O_2$  concentration of 20  $\mu$ M, were measured as described previously (Penketh and Klein, 1986). Rates of  $H_2O_2$  metabolism of 3.5 nmol/10 $^8$  cells/min (n = 5, SD = 0.6), 18.3 nmol/10 $^8$  cells/min (n = 3, SD = 1.5) and 48.2 nmol/10 $^8$  cells/min (n = 3, SD = 1.9), respectively, were obtained. Rates of  $H_2O_2$  metabolism in 11 other trypanosomatids have been determined (Penketh et al., 1987), and rates comparable to or greater than those found in T. b. rhodesiense were observed.

In view of these findings, trypanosomes cannot be considered deficient in their ability to metabolize  $H_2O_2$  or particularly sensitive to  $H_2O_2$  stress. The very high rate of  $H_2O_2$  metabolism in *T. cruzi* trypomastigote forms compared to epimastigote forms probably has a pathophysiological function, and correlates with their previously reported greater resistance to the phagocytic killing mechanism and the oxidant stress in the form of  $H_2O_2$  (Tanaka et al., 1983).

Declaration. All animal procedures described are in compliance with USA regulations.

Acknowledgments. P.G.P., C.L.P., and A.C.S. acknowledge support from USPHS Grant AI15742 and a grant from the MacArthur Foundation. – W.P.K. K. acknowledges support from the Wolfson Foundation.

- Docampo R., Moreno S. N. J.: Free-radical intermediates in the antiparasitic action of drugs and phagocytic cells. In: Free radical's in biology, ed. by W. A. Pryor, p. 263–288. Academic Press, New York 1984.
- Lanham S. M.: Separation of trypanosomes from the blood of infected rats and mice by anion-exchange. Nature (Lond.), 218, 1273–1274 (1968).
- Penketh P. G.: Nitrofurazone and hydrogen peroxide metabolism in *Trypanosoma brucei*. Ph.D. thesis, University of Cambridge 1986.
- Penketh P. G., Klein R. A.: Hydrogen peroxide metabolism in *Trypanosoma brucei*. Mol. Biochem. Parasit. *20*, 111–121 (1986).
- Penketh P. G., Kennedy W. P. K., Patton C. L., Sartorelli A. C.: Trypanosomatid hydrogen peroxide metabolism. FEBS Lett. *221*, 427–431 (1987).
- Piras M. M., Piras R., Henriquez D.: Changes in morphology and infectivity of cell culture-derived trypomastigotes of *Trypanosoma cruzi*. Mol. Biochem. Parasit. 6, 67–81 (1982).
- Ragu G., Striker L., Harlan J., Gown A., Striker G.: Cytoskeletal changes as an early event in hydrogen peroxide-induced cell injury: a study in A 549 cells. Brit. J. exp. Path. 67, 105–112 (1986).
- Rosenberg H. M., Mathews E.: Short-term effects of ionizing radiation on volume-regulation of murine lymphoma cells in vitro. Evidence for the involvement of hydrogen peroxide. Int. J. Radiat. Biol. 23, 91–94 (1973).
- Tanaka Y., Tanowitz H., Bloom B. R.: Growth of *Trypanosoma cruzi* in a cloned macrophage cell line and a variant defective in oxygen metabolism. Infect. Immun. 41, 1322–1331 (1983).
- Widmer G. A.: Biochemical and epidemiological study of enzyme polymorphism in *Trypanosoma cruzi*. Ph.D. thesis, London School of Hygiene and Tropical Medicine 1986.