**Zeitschrift:** Acta Tropica

**Herausgeber:** Schweizerisches Tropeninstitut (Basel)

**Band:** 45 (1988)

Heft: 2

**Artikel:** Infection rates in sterile males of "morsitans", "palpalis" and "fusca"

groups "Glossina" for pathogenic "Trypanosoma" species from East and

West Africa

Autor: Moloo, S.K. / Kutuza, S.B. / Desai, J. DOI: https://doi.org/10.5169/seals-314067

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 24.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

International Laboratory for Research on Animal Diseases (ILRAD), Nairobi, Kenya

# Infection rates in sterile males of morsitans, palpalis and fusca groups Glossina for pathogenic Trypanosoma species from East and West Africa

S. K. Moloo, S. B. Kutuza, J. Desai

# **Summary**

Infection rates in sterile male Glossina morsitans centralis, G. austeni, G. palpalis, palpalis, G. p., gambiensis, G. fuscipes fuscipes, G. tachinoides and G. brevipalpis for Trypanosoma vivax, T. congolense and T. brucei isolated from East and West Africa, were studied. Five groups of the sterile males, together with the five groups of sexually fertile males, of each of the respective species and subspecies were allowed to feed for 24 days on a Boran calf or goats infected with T. vivax or T. congolense, or with T. brucei for 34 days, after which they were dissected. The results showed that the infection of the pathogenic Trypanosoma species became better established in some tsetse species than in others. Also, the infection rates of T. vivax, T. congolense and T. brucei for sterile and sexually fertile males of any of the above Glossina did not differ significantly. These results indicate that releases of sterile male tsetse in the tsetse control programme will potentially increase the risk of trypanosomiasis during the period of tsetse releases in the affected areas, unless in the areas with low tsetse density the sterile male tsetse are rendered refractory to trypanosome infection prior to their releases while in the areas with medium to high tsetse densities, the resident tsetse populations are initially reduced with insecticides, traps and/or targets.

**Key words:** Glossina morsitans centralis; G. austeni; G. palpalis palpalis; G. p. gambiensis; G. fuscipes fuscipes; G. tachinoides; G. brevipalpis; gamma-irradiation; sterile males; infection rates; Trypanosoma vivax; T. congolense; T. brucei; ox; goats.

Correspondence: Dr. S. K. Moloo, International Laboratory for Research on Animal Diseases (ILRAD), P.O. Box 30709, Nairobi, Kenya

10 Acta Tropica 145

## Introduction

Sterile insect release (SIR) is a practical method for eradication or control of tsetse populations (Simpson, 1958; Dame and Schmidt, 1970). This genetic method was used to eradicate Glossina palpalis gambiensis from an area in Burkina Faso (Cuisance et al., 1980; Politzar and Cuisance, 1982) and G. morsitans morsitans from Mkwaja Ranch in Tanzania (Williamson et al., 1983). In both these attempts to eradicate tsetse, the resident tsetse populations were initially reduced with the aerial application of insecticides. The SIR method was also used in combination with biconical traps and insecticide-impregnated targets for the eradication of G. p. palpalis in a 1500-km<sup>2</sup> area of central Nigeria (Takken et al., 1986). However, it has been shown that sterile male G. m. morsitans, G. m. centralis, G. austeni and G. tachinoides are as efficient vectors of pathogenic Trypanosoma species as sexually fertile males (Moloo, 1982; Moloo and Kutuza, 1984). This paper reports on further investigations on the infection rates in gamma-irradiated sterile male tsetse for T. vivax, T. congolense and T. brucei isolated from East and West Africa. Seven species and subspecies of Glossina belonging to the three main taxonomic groups, morsitans, palpalis and fusca, were studied.

#### Materials and Methods

The morsitans group tsetse used were G. m. centralis originating from mainland Tanzania and G. austeni from Zanzibar; the palpalis group were G. p. palpalis from Nigeria, G. p. gambiensis from Burkina Faso, G. fuscipes fuscipes from the Central African Republic and G. tachinoides from Chad; the fusca group tsetse used was G. brevipalpis from Kenya. All seven tsetse species and subspecies were from the production colonies bred at ILRAD. They were maintained under controlled climatic conditions at 25°C, with 12 h of subdued indirect light during the day and 12 h of darkness at night. Tsetse in the morsitans group were kept at 70% relative humidity, and the others at 85%. All the breeding tsetse colonies were allowed to feed on lop-eared rabbits daily except at weekends.

T. vivax IL 2241 is a derivative of KETRI 2375 which was isolated from a naturally infected cow at Likoni, Coast Province, Kenya in 1978 (Wellde et al., 1983). T. vivax IL 1392 is a derivative of Zaria Y486 which was isolated from a Zebu cow in Nigeria in 1973 (Leeflang et al., 1976). T. congolense IL 2047 is a derivative of STIB 212 which was isolated from a lion in Tanzania in 1971 (Geigy and Kauffmann, 1973). T. congolense IL 2281 is a cloned stock of TREU 1290 (Luckins and Gray, 1979) which was derived from Zaria/67/LUMP/69 isolated from a cow in Nigeria in 1967 (Luckins et al., 1986). T. brucei IL 923 is a derivative of STIB 247 isolated from a Coke's hartebeest in Tanzania in 1971 (Geigy and Kauffmann, 1973). T. brucei IL 2380 is a derivative of TREU 1317 isolated from a naturally infected pig in Nigeria in 1962 (Killick-Kendrick and Godfrey, 1963).

A one-year old Boran calf, reared at ILRAD and without previous exposure to haemoparasitic infections, as well as adult East African Galla crossbred goats, weighing 20–25 kg, we used. The goats were obtained from an area of Kenya which is free of trypanosomiasis and were screened for antibodies to trypanosomes by the indirect immunofluorescent antibody test (Wilson, 1969) and shown to be negative. All the animals were housed in maximum insect-proof isolation units, and were allowed access to water and hay with concentrate ration.

Teneral male G. m. centralis, G. austeni, G. p. palpalis, G. p. gambiensis, G. f. fuscipes, G. tachinoides and G. brevipalpis were exposed to 12 krad  $\pm 2.5\%$  (dose rate, 580 rad/min) in a <sup>137</sup>Cs radiation source under ambient conditions. The irradiated teneral males and about the same number of unirradiated teneral control males of the respective species were allowed to feed concurrently on the

Boran calf infected with *T. vivax* IL 2241, daily except Sundays. On day 25 post-emergence, all the surviving males were dissected to determine the infection rates. Four additional groups of irradiated and of unirradiated teneral males of each of the above seven tsetse species and subspecies were treated as above, but on different days, and their infection rates determined. This experimental procedure was followed to determine the infection rates of gamma-irradiated sterile males and the unirradiated control males of all the seven species and subspecies of *Glossina*. They were fed on the goats infected with *T. vivax* IL 1392, *T. congolense* IL 2047, *T. congolense* IL 2281, *T. brucei* IL 923 or *T. brucei* IL 2380; five different goats were used for each stock but the test and the control groups of each batch were fed on the same goat and at the same time. Tsetse infected with *T. congolense* were dissected on day 25 while those infected with *T. brucei* were dissected on day 35 after emergence.

## Results

Table 1 shows that the infection rates of *T. vivax* stocks IL 2241 isolated from Likoni, Kenya, were not significantly different between the sexually sterile and fertile males in each of the seven *Glossina* species and subspecies. However, the mature infection rates for this *T. vivax* stock were very high in G. m. centralis (55.9%, 61.1%) and *G. brevipalpis* (69.7%, 75.3%) compared to the rates in *G. austeni* (4.2%, 1.8%), *G. p. palpalis* (0%, 0%), *G. p. gambiensis* (1.8%, 1.3%), *G. f. fuscipes* (0.3%, 0.5%) and *G. tachinoides* (0%, 0.3%). Table 2 shows that the infection rates of *T. vivax* IL 1392 from Nigeria also were not significantly different between the sterile and fertile males of any of the seven tsetse species and subspecies. In contrast to the Kenyan *T. vivax*, the mature infection rates for the Nigerian *T. vivax* were very high in all the seven tsetse species and subspecies examined (range 39.9–81.6%).

Table 3 shows that there were no differences in the infection rates of *T. congolense* IL 2047 from Tanzania, between the sterile and fertile males in any of the seven tsetse species and subspecies. However, the mature infection rates were very high in *G. m. centralis* (44.0%, 35.5%), lower in *G. brevipalpis* (18.3%, 15.7%) and very low in *G. austeni* (0%, 2.0%), *G. p. palpalis* (0.5%, 0.5%), *G. p. gambiensis* (0%, 0.4%), *G. f. fuscipes* (1.6%, 6.0%) and *G. tachinoides* (0.3%, 0.2%). A similar pattern of infection rates was observed for *T. congolense* IL 2281 from Nigeria (Table 4).

Table 5 shows that there were some differences in the mature infection rates of *T. brucei* in the sterile and fertile males of the seven tsetse species and subspecies but the differences were small and inconsistent and probably not significant. Again, there were species and subspecies differences in the infection rates. In *G. m. centralis*, the mature *T. brucei* infection rates were very high (50.9%, 40.4%) whereas in the other six tsetse species and subspecies, they were extremely low (range 0–2.1%). A similar pattern for the mature infection rates was observed for *T. brucei* IL 2380 from Nigeria; the infection did not mature to metacyclics in the salivary glands of *G. austeni*, *G. f. fuscipes* and *G. brevipalpis* (Table 6).

Table 1. Infection rates of *Trypanosoma vivax* IL 2241 from East Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number infected	Number dissected	Infection rates (%) $\pm$ SE	
		mected	dissected	Labrum	Hypopharynx
G. m. centralis	Irradiated	500	338	58.0±12.6	55.9±13.0
	Control	500	379	66.9±9.3	61.1±7.8
G. austeni	Irradiated	500	360	4.9±4.2	4.2±3.6
	Control	500	374	2.5±1.6	1.8±1.2
G. p. palpalis	Irradiated	500	414	0.2	0.0
	Control	500	344	0.0	0.0
G. p. gambiensis	Irradiated	500	370	2.5±1.4	1.8±1.5
	Control	500	416	3.4±1.9	1.3±1.0
G. f. fuscipes	Irradiated	490	342	0.5	0.3
	Control	490	332	0.8	0.5
G. tachinoides	Irradiated	500	369	0.4	0.0
	Control	500	389	0.5	0.3
G. brevipalpis	Irradiated	435	354	76.8±8.2	69.7±8.6
	Control	436	391	81.1±6.3	75.3±6.8

The results given as mean  $\pm$  SE are from five independent experiments.

Table 2. Infection rates of *Trypanosoma vivax* IL 1392 from West Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number	Number	Infection rates (%) $\pm$ SE	
		infected	dissected	Labrum	Hypopharynx
G. m. centralis	Irradiated	500	386	66.0±10.4	49.5±6.2
	Control	500	403	63.1±8.9	46.3±8.4
G. austeni	Irradiated	500	410	62.7±7.7	50.0±4.9
	Control	500	427	70.8±4.5	54.8±4.2
G. p. palpalis	Irradiated	500	455	52.4±8.5	45.3±7.8
	Control	500	437	45.3±10.2	39.9±9.7
G. p. gambiensis	Irradiated	500	406	64.0±7.2	58.1±6.7
	Control	500	436	69.8±5.0	62.8±4.0
G. f. fuscipes	Irradiated	486	384	47.7±7.5	42.2±6.5
	Control	486	403	46.3±5.1	41.2±4.9
G. tachinoides	Irradiated	500	423	50.6±11.9	41.7±12.3
	Control	500	457	51.5±9.4	41.5±9.0
G. brevipalpis	Irradiated	385	344	86.8±8.3	75.4±12.6
	Control	385	348	92.1±5.7	81.6±7.5

The results given as mean  $\pm$  SE are from five independent experiments.

Table 3. Infection rates of *Trypanosoma congolense* IL 2047 from East Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number infected	Number dissected	Infection rates (%) ±SE		
				Gut	Labrum	Hypopharynx
G. m. centralis	Irradiated	500	324	57.4±8.1	45.6±8.6	44.0±8.7
	Control	500	374	49.3±8.6	36.5±5.6	35.3±5.7
G. austeni	Irradiated	500	352	14.5±5.3	0.5	0.0
	Control	500	377	14.4±4.0	3.4±0.6	2.0±1.0
G. p. palpalis	Irradiated	447	362	8.8±4.2	0.8	0.5
	Control	447	340	10.2±4.7	0.8	0.5
G. p. gambiensis	Irradiated	493	315	6.9±2.6	0.2	0.0
	Control	493	339	5.9±2.9	0.4	0.4
G. f. fuscipes	Irradiated	434	253	13.8±5.0	$2.5\pm1.2$	1.6±0.5
	Control	434	273	13.2±5.5	$6.7\pm3.4$	6.0±3.4
G. tachinoides	Irradiated	445	313	8.6±5.7	0.3	0.3
	Control	446	357	7.2±2.7	0.2	0.2
G. brevipalpis	Irradiated	425	349	30.5±5.8	20.1±5.5	18.3±6.1
	Control	424	354	33.8±5.0	17.6±5.0	15.7±5.4

The results given as mean  $\pm$  SE are from five independent experiments.

Table 4. Infection rates of *Trypanosoma congolense* IL 2281 from West Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number infected	Number	Infection rates (%) ±SE		
			dissected	Gut	Labrum	Hypopharynx
G. m. centralis	Irradiated	500	403	69.1±3.6	56.4±3.4	55.8±3.8
	Control	500	421	63.5±5.4	49.9±4.2	49.2±4.0
G. austeni	Irradiated	489	408	14.0±4.9	3.1±0.5	2.1±0.6
	Control	490	419	14.6±6.7	3.4±1.2	3.0±1.0
G. p. palpalis	Irradiated	388	284	9.5±3.4	1.4±0.6	1.0±0.3
	Control	388	286	18.7±12.4	0.7±0.4	0.7±0.3
G. p. gambiensis	Irradiated	485	395	16.3±11.7	0.5±0.3	0.3
	Control	485	363	15.6±12.2	2.2±0.6	0.3
G. f. fuscipes	Irradiated	445	316	16.1±3.7	6.8±2.6	1.5±0.5
	Control	444	301	9.2±2.7	3.2±1.1	0.9±0.7
G. tachinoides	Irradiated	483	360	11.1±9.4	1.2±0.9	0.8±0.5
	Control	482	401	9.8±4.3	2.5±0.7	2.0±0.5
G. brevipalpis	Irradiated	459	369	42.0±7.8	17.1±4.9	8.8±3.3
	Control	459	331	40.3±11.4	15.4±4.0	6.3±2.0

The results given as mean  $\pm$  SE are from five independent experiments.

Table 5. Infection rates of *Trypanosoma brucei* IL 923 from East Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number infected	Number dissected	Infection rates (%) ±SE	
			dissected	Gut	Salivary glands
G. m. centralis	Irradiated	482	208	80.5±5.1	50.9±3.7
	Control	481	357	80.7±4.9	40.4±2.7
G. austeni	Irradiated	500	427	10.2±2.1	1.0±0.6
	Control	500	426	5.4±1.9	0.2
G. p. palpalis	Irradiated	426	350	$5.5\pm0.9$	0.4
	Control	427	353	$3.7\pm1.3$	0.4
G. p. gambiensis	Irradiated	481	413	$2.0\pm0.6$	0.0
	Control	481	436	$2.3\pm0.3$	0.4
G. f. fuscipes	Irradiated	371	329	$6.0\pm1.2$	1.6
	Control	371	302	$5.6\pm1.1$	2.1±0.8
G. tachinoides	Irradiated	425	313	$5.0\pm1.2$	1.5
	Control	421	355	$3.3\pm0.3$	1.0
G. brevipalpis	Irradiated	481	342	64.8±2.3	0.0
	Control	482	402	59.3±3.4	0.2

The results given as mean  $\pm$  SE are from five independent experiments.

Table 6. Infection rates of *Trypanosoma brucei* IL 2380 from West Africa in irradiated and unirradiated control males of seven different *Glossina* species and sub-species

Tsetse species	Treatment	Number infected	Number dissected	Infection rates (%) ±SE		
				Gut	Salivary glands	
G. m. centralis	Irradiated	494	216	45.0±9.6	12.3±3.4	
	Control	494	440	36.4±5.2	6.8±1.7	
G. austeni	Irradiated	482	385	0.3	0.0	
	Control	486	419	0.9	0.0	
G. p. palpalis	Irradiated	386	319	1.2±0.6	0.4	
	Control	385	353	1.5±0.6	0.0	
G. p. gambiensis	Irradiated	459	379	0.5	0.2	
	Control	459	391	0.4	0.0	
G. f. fuscipes	Irradiated	345	289	0.0	0.0	
	Control	350	279	0.8	0.0	
G. tachinoides	Irradiated	427	284	$5.0\pm1.4$	1.3±0.6	
	Control	418	372	$2.2\pm0.6$	0.7±0.4	
G. brevipalpis	Irradiated	500	386	23.5±5.6	0.0	
	Control	500	402	18.1±5.0	0.0	

The results given as mean  $\pm$ SE are from five independent experiments.

## Discussion

The present study has demonstrated that male G. m. centralis, G. austeni, G. p. palpalis, G. p. gambiensis, G. f. fuscipes, G. tachinoides and G. brevipalpis rendered sexually sterile by gamma-irradiation are as susceptible to T. vivax, T. congolense and T. brucei, as sexually fertile corresponding males irrespective of whether the trypanosome isolate was from East or West Africa. These results confirm and extend our earlier observations that the use of gamma-irradiation to sterilize male Glossina prior to infection does not interfere with subsequent cyclical development of these parasites. By analogy with what has previously been shown for G. m. morsitans (Moloo, 1982), it is likely that the transmission characteristics of T. vivax, T. congolense and T. brucei to susceptible hosts by sterile males of the above tsetse species and subspecies remain unchanged by sterilization.

The results also demonstrated that some tsetse species are better vectors of the three pathogenic *Trypanosoma* species than others, but this selectivity is also unimpaired by gamma-irradiation.

The SIR method has been used in three previous campaigns to eradicate tsetse; G. p. gambiensis (Politzar and Cuisance, 1982) from an area in Burkina Faso, G. m. morsitans (Williamson et al., 1983) from Mkwaja Ranch in Tanzania, and G. p. palpalis (Takken et al., 1986) in an area in central Nigeria. In each of these campaigns, the tsetse populations were initially reduced by the application of insecticides prior to the sterile male releases. Consequently fewer sterile insects were required for release and so there was no associated increase in the risk of trypanosomiasis in the above areas. However, it has recently been shown that the incorporation of a trypanocidal drug, isometamidium chloride ('Samorin', May & Baker Ltd., Dagenham, England) in a bloodmeal completely suppressed the cyclical development within G. m. centralis of T. vivax, T. congolense or T. brucei. This was achieved by feeding the sterile teneral male tsetse on a bloodmeal containing 12 µg ml<sup>-1</sup> Samorin prior to a feed on infected goats (Moloo and Kamunya, 1987). Hence, inclusion in a SIR programme for tsetse eradication of a pre-release bloodmeal containing Samorin for the sterile teneral males would achieve two objectives. Firstly, is could effectively suppress the disease transmission by such tsetse and, secondly, in areas with medium or low tsetse densities it could reduce or eliminate the need for initial reduction of the resident tsetse populations with insecticides. However, in areas with high tsetse densities, the initial reduction of native tsetse populations will certainly have to be incorporated in SIR programme for tsetse eradication.

### Acknowledgments

We thank Mr. C.O. Machika and Mr. F. K. Masasi for excellent technical assistance. We are grateful to Dr. V. M. Nantulya and Dr. D. D. Whitelaw for helpful comments on the manuscript. – This is ILRAD Publication No. 560.

- Cuisance D., Politzar H., Clair M., Sellin E., Taze Y., Bourdoiseau G., Fevrier J.: La lutte contre *Glossina palpalis gambiensis* Vanderplank par lâchers de mâles irradiés en Haute-Volta. Etude de paramètres opérationnels. Proceedings of the IAEA/FAO Symposium, Vienna, 7–11 May 1979, p. 249–264 (1980).
- Dame D. A., Schmidt C. H.: The sterile-male technique against tsetse flies, *Glossina* species. Bull. ent. Soc. Amer. *16*, 24–30 (1970).
- Geigy R., Kauffmann M.: Sleeping sickness survey in the Serengeti area (Tanzania) 1971: examination of large mammals for trypanosomes. Acta trop (Basel) 30, 12–23 (1973).
- Killick-Kendrick R., Godfrey D. G.: Observations on a close association between *Glossina tachi-noides* and domestic pigs near Nsukka, Eastern Nigeria. 1. *Trypanosoma congolense* and *T. brucei* infections in the pigs. Ann. trop. Med. Parasit. 57, 225–231 (1963).
- Leeflang P., Buys J., Blotkamp C.: Studies on *Trypanosoma vivax*: infectivity and serial maintenance of natural bovine isolates in mice. Int. J. Parasit. 6, 413–417 (1976).
- Luckins A. G., Frame I. A., Gray M. A., Crowe J. S., Ross C. A.: Analysis of trypanosome variable antigen types in cultures of metacyclic and mammalian form of *Trypanosoma congolense*. Parasitology *93*, 99–109 (1986).
- Luckins A. G., Gray A. R.: Observations on the antigenicity and serological relationships of stocks of *Trypanosoma congolense* from East and West Africa. Parasitology 79, 337–347 (1979).
- Moloo S. K.: Cyclical transmission of pathogenic *Trypanosoma* species by gamma-irradiated sterile male *Glossina morsitans morsitans*. Parasitology 84, 289–296 (1982).
- Moloo S. K., Kamunya G. W.: Suppressive action of Samorin on the cyclical development of pathogenic *Trypanosoma* species in *Glossina morsitans centralis*. Med. Vet. Ent. 1, 285–287 (1987).
- Moloo S. K., Kutuza S. B.: Vectorial capacity of gamma-irradiated sterile male *Glossina morsitans* centralis, G. austeni and G. tachinoides for pathogenic Trypanosoma species. Insect. Sci. Applic. 5, 411–414 (1984).
- Politzar H., Cuisance D.: SIT in the control and eradication of *Glossina palpalis gambiensis*. Proceedings of IAEA/FAO Symposium, Neuherberg, 29 June to 3 July 1981 (1982).
- Simpson H. R.: The effect of sterilized males on a natural tsetse fly population. Biometrics 14, 159–173 (1958).
- Takken W., Oladunmade M. A., Dengwat L., Feldmann H. U., Onah J. A., Tenabe S. O., Hamann H. J.: The eradication of *Glossina palpalis palpalis* (Robineau-Desvoidy) (Diptera: Glossinidae) using traps, insecticide- impregnated targets and the sterile insect technique in central Nigeria. Bull. ent. Res. 76, 275–286 (1986).
- Wellde B. T., Chumo D. A., Adoyo M., Kovatch R. M., Mwongela G. N., Opiyo E. A.: Haemorrhagic syndrome in cattle associated with *Trypanosoma vivax* infection. Trop. Anim. Hlth Prod. *15*, 95–102 (1983).
- Williamson D. L., Dame D. A., Gates D. B., Cobb P. E., Bakuli B., Warner P.: Integration of insect sterility and insecticides for control of *Glossina morsitans morsitans* Westwood (Diptera: Glossindae) in Tanzania. V. The impact of sequential releases of sterilised tsetse flies. Bull. ent. Res. 73, 391–404 (1983).
- Wilson A. J.: Value of the indirect fluorescent antibody test as serological aid to diagnosis of *Glossina*-transmitted bovine trypanosomiasis. Trop. Anim. Hlth Prod. 1, 89–95 (1969).