

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 45 (1988)
Heft: 2

Artikel: Trypanosome infection rates in tsetse midguts using a short-term in vitro culture technique : short communication
Autor: Mehltz, D. / Tietjen, U.
DOI: <https://doi.org/10.5169/seals-314072>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Bernhard-Nocht-Institut für Schiffs- und Tropenkrankheiten, Abteilung für Veterinärmedizin,
Hamburg, Federal Republic of Germany and
Liberia Research Unit of the Tropical Institute Hamburg (LRU), Monrovia, Liberia

Trypanosome infection rates in tsetse midguts using a short-term in vitro culture technique

Short communication

D. MEHLITZ, U. TIETJEN

To assess the risk of trypanosome infection or 'challenge' which tsetse flies present to animal or human hosts, it is first necessary to determine the infection rates in the flies. Trypanosome infection rates in tsetse have customarily been assessed by dissection and microscopical examination of the proboscis, midgut and salivary glands (Lloyd and Johnson, 1924). Using this dissection technique infection rates in wild flies have usually been found to be very low (Jordan, 1974); this reflects both the inherent refractoriness of most flies to infection (Maudlin et al., 1986) and the insensitivity of dissection as a diagnostic technique.

In attempting to isolate trypanosomes from wild flies we placed tsetse midguts in culture medium, and during the course of these experiments, established procyclic cultures from flies which were found to be uninfected by the dissection method. Following these findings, we surveyed the infection rates in wild flies using the two diagnostic techniques – dissection and culture. We present here the results of this survey carried out in Liberia.

Glossina palpalis palpalis (n = 962), *G. nigrofusca* (n = 122) and *G. pallicera* (n = 28) were caught in the Liberian rain forest using biconical traps (Challier and Laveissière, 1973). Flies were dissected in the laboratory of the LRU within 24 h of capture. Dissection took place in a clean environment with flame sterilized instruments, on sterile slides in a few drops of sterile dissection and isolation medium for procyclics (DIMP). This solution was a mixture of 10–15% foetal calf serum (FCS), 100 µg/ml gentamycin and 50 µg/ml mycostatin in 50% Cunningham's Medium (Cunningham, 1977) and 50% SDM-79 (Brun and Schönenberger, 1979).

The proboscis (hypopharynx, labrum, labium), midgut and salivary glands was examined under a Zeiss-microscope (128× and 320× magnification). Immediately after microscopic examination each midgut was transferred into one well of a 96-microwell dish (Nunclon) containing approx. 100 µl DIMP. The guts were incubated for 3 days at 27°C and examined daily with a Wild-invertoscope (200× magnification).

For all three *Glossina* species the overall mature infection rates determined by dissection were: 1.3% for *Nannomonas* and 1.7% for *Duttonella*, *Trypanozoon* were not detected (Table 1). Immature (midgut) trypanosomes were diagnosed in 44 (4%) of the 1112 flies examined. In contrast, in vitro cultivation of the midguts from these same flies recovered 107 (9.6%) procyclic midgut infections (6.05% on day 0, 2.2% on day 1, 0.7% on day 2, 0.6% on day 3 and 0.1% after day 3 of cultivation). Various fungal contaminations in the cultures prevented a longer observation period.

Correspondence: Dr. D. Mehlitz, Bernhard-Nocht-Institut, Bernhard-Nocht-Strasse 74,
D-2000 Hamburg 4

Table 1. Tsetse fly infection rates with trypanosomes determined by dissection and short-term in vitro culture technique

<i>Glossina</i> species	No. ex- am- ined	Dissection		In vitro culture	
		mature		immature	
		infections with			
		<i>N.</i> ¹	<i>D.</i> ²	<i>N.</i> or <i>T.</i> ³	<i>N.</i> or <i>T.</i>
<i>G. p. palpalis</i>	962	13 (1.4%)	12 (1.2%)	36 (3.7%) ⁴	86 (8.9%) ⁴
<i>G. nigrofusca</i>	122	2 (1.6%)	6 (4.9%)	5 (4.1%) ⁴	17 (13.9%) ⁴
<i>G. pallicera</i>	28	0	1 (3.6%)	3 (10.7%) ⁴	4 (14.3%) ⁴
Total	1112	15 (1.3%)	19 (1.7%)	44 (4.0%) ⁴	107 (9.6%) ⁴

¹*N.* = *Nannomonas*, ²*D.* = *Duttonella*, ³*T.* = *Trypanozoon*

⁴Flies with mature infections (*Nannomonas*) included

These findings show that the in vitro culture technique is significantly ($p < 0.001$) more sensitive than the dissection method in detecting immature *Nannomonas* or *Trypanozoon* infections; the ratio of mature: immature infections being increased from 1:1.9 (dissection) to 1:6.1 (culture). Examination of flies by dissection alone may lead to an underestimation of the numbers of immature midgut infections present in the fly population. The short-term in vitro culture technique described here could be used routinely to more accurately determine infection rates. The detection and discrimination of *Trypanosoma* spp. in *Glossina*, particularly of *T. congolense*, *T. simiae*, and human-infective and non-infective *T. brucei*, would probably reflect the variety of the entire trypanosome population in the host-fly-host cycle better than analysis of bloodstream forms derived from mammals.

Acknowledgments. This study was supported by the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases. The authors wish to thank the Ministry of Health and Social Welfare and the Ministry of Agriculture, Monrovia, Liberia, for support and cooperation.

Brun R., Schönerberger M.: Cultivation and in vitro cloning of procyclic culture forms of *Trypanosoma brucei* in semi-defined medium. *Acta trop.* (Basel) 36, 289–292 (1979).

Challier A., Laveissière C.: Un nouveau piège pour la capture des glossines (*Glossina*: *Diptera*, *Muscidae*): description et essais sur le terrain. *Cah. ORSTOM, sér. Ent. méd. Parasit.* 11, 251–262 (1973).

Cunningham I.: New culture medium for maintenance of tsetse tissues and growth of trypanosomatids. *J. Protozool.* 24, 325–329 (1977).

Jordan A. M.: Recent developments in the ecology and methods of control of tsetse flies (*Glossina* spp.; *Dipt. Glossinidae*) – a review. *Bull. ent. Res.* 63, 361–399 (1974).

Lloyd L., Johnson W. B.: The trypanosome infections of tsetse flies in Northern Nigeria and a new method of estimation. *Bull. ent. Res.* 14, 265–288 (1924).

Maudlin I., Dukes P., Luckins A. G., Hudson K. M.: Extrachromosomal inheritance of susceptibility to trypanosome infection in tsetse flies. II. Susceptibility of selected lines of *Glossina morsitans morsitans* to different stocks and species of trypanosome. *Ann. trop. Med. Parasit.* 80, 97–105 (1986).