

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 45 (1988)
Heft: 3

Artikel: Trypanosome-induced ovarian dysfunction : evidence of higher residual fertility in trypanotolerant small East African goats
Autor: Mutayoba, B.M. / Gombe, S. / Kaaya, G.P.
DOI: <https://doi.org/10.5169/seals-314080>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

¹ Reproductive Biology Unit, NCCR, University of Nairobi, P.O. Box 30197, Nairobi, Kenya

² International Centre of Insect Physiology and Ecology, P.O. Box 30772, Nairobi, Kenya

Trypanosome-induced ovarian dysfunction

Evidence of higher residual fertility in trypanotolerant small East African goats

B. M. MUTAYOBA¹, S. GOMBE¹, G. P. KAAAYA², E. N. WAINDI¹

Summary

Changes in the length of oestrous cycles, plasma progesterone and oestradiol-17 β levels were monitored for 6 months in *Trypanosoma congolense*-infected normocyclic small East African goats obtained from three tsetse-endemic areas and one tsetse-free area of East Africa. Irregular oestrous cycles were observed in all infected goats, before cessation at the second cycle post-infection in the more susceptible and fourth cycle in the more resistant goat groups. A significant decline in the progesterone and oestradiol-17 β parameters were observed. The decline in hormonal values was, however, less in the more resistant than in the susceptible goat groups at least in the first 2 months post-infection. Resumption of the ovarian cycle were observed in few resistant goats after 5 months of the infection. It is concluded that clinical tolerance is correlated with residual fertility, i.e., the greater the tolerance the higher the retention of fertility.

Key words: ovarian dysfunction; residual fertility; trypanotolerance; *Trypanosoma congolense*.

Introduction

The use of trypanotolerant livestock is now considered to be an important alternative strategy for increasing animal numbers in tsetse-endemic areas of Africa where other conventional disease control measures have proved uneconomical and/or ineffective. With the increasing reports of the occurrence of severe degenerative lesions of the reproductive organs in trypanosusceptible

Correspondence: B. M. Mutayoba, Sokoine University of Agriculture, Faculty of Veterinary Medicine, P.O. Box 3017, Morogoro, Tanzania

animals (Ikede and Losos, 1972, 1975; Isoun et al., 1975; Ikede, 1979; Anosa and Isoun, 1980; Kaaya and Oduor-Okelo, 1980; Waindi et al., 1986), it has recently become necessary to examine the degree of gonadal dysfunction in trypanotolerant livestock before wide exploitation of these animals is encouraged. To what extent trypanotolerant animals retain their fertility following trypanosome infection is not well known. Since, trypanosome-infected animals develop varying degrees of reproductive impairments, the term "residual fertility" is used in this paper to denote the resulting fertility which potential remain following the infection.

Murray et al. (1981) reported that N'dama cattle can survive and retain a higher degree of residual fertility in tsetse-endemic areas where Zebu perish. Recently, Mutayoba et al. (1987) showed that Small East African breed of goats obtained from Morogoro, a tsetse-endemic area in Tanzania, were more tolerant to experimental chronic *Trypanosoma congolense* infection than goats obtained from Arusha in Tanzania and Lambwe Valley in Kenya (also tsetse-endemic areas). Goats obtained from Imbo, a tsetse-free enclave in Central Nyanza, Kenya, however, were much more susceptible as assessed by parasitaemic and haematic parameters, body weight change and mortality rates. The present paper reports on the changes in reproductive function in the same groups of goats (for which previous data was obtained) following the infection. The relationship between trypanotolerance and higher residual fertility is discussed.

Materials and Methods

A total of 52 normocyclic Small East African female goats (Mutayoba et al., 1987) aged between 2–3 years were used. Ten female goats obtained from each of the tsetse-endemic areas (Morogoro in Tanzania and Lambwe Valley in Kenya) and tsetse-free (Imbo, Kenya) and 9 goats from Arusha, Tanzania (tsetse-endemic area) were experimentally inoculated via the jugular vein with approximately 2.5×10^4 *T. congolense* (EATRO-1753) obtained from Kenya Trypanosomiasis Research Institute (KETRI).

Thirteen uninfected goats from all experimental groups served as controls.

Prior to the infection, all goats were acclimatized for 1½ months and screened for haemoprotozoa using Giemsa-stained smears and were found negative. At the end of the acclimatization period, male bucks with deviated penises were introduced to monitor overt oestrus. At the same time, all females were bled (5 ml/goat) for two consecutive cycles between 9.00–10.00 a.m. every other day to provide plasma for baseline pre-infection self-control hormonal levels. The bleeding continued throughout the infection period at the same rate except during oestrus when it was done at 6 hourly intervals. The plasma obtained was stored at -20°C until needed for hormonal analyses. The infection period lasted 6 months.

Plasma concentrations of progesterone and oestradiol-17 β were determined in 200 μl and 500 μl plasma aliquots, respectively, by dextran-charcoal radioimmunoassay according to the WHO (1984/85 matched reagent manual) and adopted for use with goat plasma (O'hara et al., 1985; Katongole and Gombe, 1985).

The antisera to progesterone and oestradiol-17 β were supplied by WHO and had been raised against progesterone 3-(o-carboxymethyl) oximino-BSA and oestradiol-6CMO-BSA, respectively. The ^3H -tracers were obtained from the Radiochemical Centre, Amersham, England.

Table 1. Estimated parasitaemia scores in the 4 groups of Small East African breed of goats during the first 2 months of trypanosome infection

	Parasitaemia scores*						Samples examined	
	0	1+	2+	3+	4+	5+		6+
Lambwe Valley	5 (7.2%)	9 (13%)	16 (23.2%)	21 (30.4%)	12 (17.4%)	5 (7.2%)	1 (1.4%)	69
Arusha	12 (16.9%)	16 (22.5%)	13 (18.3%)	18 (25.4%)	9 (12.7%)	3 (4.2%)	0	71
Morogoro	21 (27.3%)	16 (20.7%)	14 (18.2%)	13 (16.9%)	10 (13%)	3 (3.9%)	0	77
Imbo	3 (4.8%)	7 (11.3%)	11 (17.7%)	17 (27.4%)	12 (19.4%)	10 (16.1%)	2 (3.2%)	62

* Parasitaemia scores were obtained using the Darkground/Phase Contrast Buffy Method described by Paris et al. (1982).

Three quality control (QC) goat plasma pools containing different concentration of each hormone were included at three places in each assay to monitor the assay drifts and interassay coefficient of variations (CV's). The interassay CV's for progesterone assay were 9.8% (n = 36) at 12.7 ± 0.3 (SEM) nmol/l; 10.4% (n = 12) at 6.4 ± 0.3 nmol/l and 13.3% (n = 21) at 2.7 ± 0.1 nmol/l. The CV's for oestradiol-17 β assays were 7.6% (n = 36) at 998.9 ± 17.7 pmol/l; 10.1% (n = 36) at 623.4 ± 10.5 pmol/l and 12.8% (n = 35) at 120.3 ± 3.5 pmol/l. The intraassay CV's were calculated from 25 similar aliquots assayed together for each QC pool and were below 8% for both hormones. The sensitivity of the assays were 0.05 nmol/l (n = 21) and 11.8 pmol/l (n = 29) for progesterone and oestradiol-17 β assays, respectively.

The respective hormonal values were averaged on monthly basis for each goat and each group and were analysed by one-way analysis of variance.

The difference between group means were compared by a modified Duncan's New Multiple Range Test (Kramer, 1956) as described by Steel and Torrie (1960).

Results

Clinical observations

Comparative detailed clinical observations in the same experimental and control groups have been described recently (Mutayoba et al., 1987). Morogoro goats were found to be more tolerant followed by the Arusha, Lambwe Valley and Imbo goats in that order. All infected goats become parasitaemic within 5–8 days and thereafter, the Morogoro goats had intermittent scanty parasitaemias than other groups. Thus, out of 77 samples examined for the presence of trypanosomes during the first two months of the infection in the Morogoro group (Table 1) 21 (27.3%) were negative, whilst, only 5 (7.2%), 12 (16.9%) and 3 (4.8%) of the 69, 71 and 62 samples examined in the Lambwe, Arusha and Imbo goats, respectively, were negative. This was significantly different ($p < 0.01$). This pattern persisted throughout the course of the infection.

The Morogoro goats had also less severe anaemia (indicated by PCV changes), weight losses and lowest mortality rate compared to the Imbo goats (Table 2). The Lambwe and Arusha goats were intermediate between the resistant (Morogoro) and susceptible (Imbo) goats.

Oestrous cycles

The lengths of the oestrous cycles between the experimental and control goats prior to the infection period were not markedly different (Table 3) and a length of 20.1 ± 0.2 days (n = 104) being recorded. The cycle lengths were determined as an interval in days between two behavioural oestrus with the aid of teaser bucks and supplemented by hormonal assays. Following the onset of parasitaemia, irregular and increasingly shorter ($p < 0.05$) overt and silent cycles (detected by hormonal analyses) were noted in all infected groups. In the more susceptible Imbo goats, cyclicity ceased at the second cycle post-infection, whilst in the more resistant Morogoro goats, shorter 3rd and 4th cycles were recorded before cessation. Ovarian cycles ceased at the end of the 3rd cycle in

Table 2. Changes in the mean monthly PCV% and body weights (BW kg) of the infected and control goats

Duration of infection	Parameters (means \pm SEM)	Lambwe Valley	Arusha	Morogoro	Imbo	Control	F ratio
Preinfection means	PCV	28.1 \pm 0.6	29.2 \pm 0.6	28.9 \pm 0.5	28.8 \pm 0.7	28.9 \pm 0.7	
	BW	22.8 \pm 1.5 (10)	19.9 \pm 1.0 (9)	19.3 \pm 1.2 (10)	21.7 \pm 1.2 (10)	21.3 \pm 0.9 (13)	
June	PCV	-5.8 ^a \pm 0.4	-4.9 ^a \pm 0.5	-4.9 ^a \pm 0.3	-7.7 ^b \pm 0.3	+0.2 ^c \pm 0.7	10.19***
	BW	-1.8 ^b \pm 0.3 (8)	-1.4 ^b \pm 0.2 (9)	-0.6 ^a \pm 0.2 (10)	-0.7 ^a \pm 0.3 (8)	+0.1 ^c \pm 0.1 (13)	3.08**
July	PCV	-5.8 ^a \pm 0.3	-5.7 ^a \pm 0.5	-4.8 ^a \pm 0.3	-8.5 ^b \pm 0.6	+1.5 ^c \pm 0.3	14.81***
	BW	-3.1 ^d \pm 0.5 (8)	-2.5 ^{ad} \pm 0.5 (7)	-0.9 ^{cb} \pm 0.3 (9)	-1.8 ^{ab} \pm 0.4 (6)	+0.1 ^c \pm 0.3 (13)	5.41***
August	PCV	-7.0 ^{bc} \pm 0.4	-6.3 ^{ab} \pm 0.6	-4.8 ^a \pm 0.5	-9.0 ^c \pm 0.7	+1.8 ^d \pm 0.4	9.05***
	BW	-3.0 ^a \pm 0.5 (6)	-2.3 ^a \pm 0.6 (7)	-1.4 ^a \pm 0.4 (7)	-2.7 ^a \pm 0.9 (4)	+0.5 ^b \pm 0.4 (13)	1.62 ns
September	PCV	-6.7 ^b \pm 0.5	-6.8 ^b \pm 0.6	-4.3 ^a \pm 0.3	-9.1 ^c \pm 0.9	+0.3 ^d \pm 0.1	12.29***
	BW	-4.2 ^b \pm 0.5 (6)	-3.7 ^{ab} \pm 0.7 (6)	-2.3 ^a \pm 0.4 (7)	-4.0 ^{ab} \pm 1.1 (4)	+0.8 ^c \pm 0.5 (13)	2.10 ns
October	PCV	-5.8 ^b \pm 0.4	-6.4 ^b \pm 0.5	-4.4 ^a \pm 0.3	-8.3 ^c \pm 0.3	+0.5 ^d \pm 0.2	10.74***
	BW	-4.5 ^b \pm 0.6 (6)	-3.6 ^{ab} \pm 0.8 (6)	-2.4 ^a \pm 0.6 (7)	-3.7 ^{ba} \pm 1.1 (4)	+1.3 ^c \pm 0.4 (13)	1.57 ns
November	PCV	-5.1 ^a \pm 0.3	-5.0 ^a \pm 0.6	-4.0 ^a \pm 0.3	-6.3 ^b \pm 0.3	+0.6 ^c \pm 0.1	4.70***
	BW	-3.5 ^b \pm 0.5 (6)	-2.5 ^{ba} \pm 0.9 (5)	-1.5 ^a \pm 0.5 (7)	-1.5 ^a \pm 0.9 (3)	+1.4 ^c \pm 0.5 (13)	2.02 ns

- Values from June onwards denote loss (-) or gain (+) from the preinfection values.

- ns, ** *** denote nonsignificance, significance at 5% and 1%, respectively in the 4 infected groups.

- Means bearing the same superscripts (a b c d) in the same row do not differ significantly from each other (p < 0.05); control values are significant at (p < 0.01).

- Numbers in brackets denote surviving goats.

Table 3. Changes in the length of oestrous cycles (days) of infected and control goats

Approximate period in months	Cycles before (-) and after (+) infection	Lambwe Valley	Arusha	Morogoro	Imbo	Control	Level of significance F ₁ ratio	Level of significance F ₂ ratio
May to	-2	20.0±0.3 (10)	19.2±0.4 (9)	19.5±0.5 (10)	19.1±0.1 (10)	19.8±0.4 (13)	0.85 ns	0.71 ns
Mid-June	-1	19.8±0.4 (10)	19.1±0.8 (9)	19.4±0.3 (10)	19.9±0.7 (9)	20.6±0.4 (13)	0.41 ns	1.32 ns
Mid-June ↓	1	17.0 ^{ab} ±2.3 (8)	17.2 ^{ab} ±1.6 (9)	19.6 ^{bc} ±1.7 (9)	15.5 ^a ±2.0 (8)	20.9 ^c ±0.4 (13)	0.91 ns	2.21 ^{**}
	2	13.3 ^a ±1.9 (8)	15.3 ^a ±2.0 (7)	15.9 ^a ±2.2 (9)	11.8 ^a ±2.0 (4)	20.7 ^b ±0.9 (12)	0.74 ns	5.56 ^{***}
	3	9.7 ^b ±1.0 (6)	14.0 ^{ab} ±2.8 (5)	15.7 ^a ±1.4 (7)	-	20.6 ^c ±0.7 (10)	2.62 ^{**}	6.21 ^{***}
	4	-	-	7.5±0.5 (3)	-	21.3±0.7 (6)	-	-
	5	-	-	-	-	23.0±1.0 (3)	-	-
Mid-October	6	-	-	-	-	-	-	-
Mid-October to November	7	-	-	19.0±0.5 (2)	20.5±0.5 (2)	21.3±0.5 ⁿ (9)	-	-
	8	-	16.0±0.5 ^c (2)	15 (1)	-	19.8±0.5 ⁿ (8)	-	-

- Values given as means ± SEM

- Numbers in brackets denote cycling goats. Goats not included are either acyclic or dead (compare with Table 2).

- F₁ and F₂ denote F values between the infected groups, infected and control groups, respectively.

- ns, ** *** = nonsignificance, significance at 5% and 1% level of confidence, respectively.

- Means bearing the same superscripts (a b c d) in the same row do not differ significantly (p < 0.05).

- ^{e-n}One cycling Arusha and 2 control goats, respectively, became pregnant and their values excluded.

Lambwe and Arusha goats. Resumption of the oestrous cycles were recorded in six infected goats following the spell of short rains in mid-October.

Seasonal ovarian activity was observed in the control goats, with highest cyclicity (100%) in May and June and least (0%) in September. Cyclicity was once again high at the start of short rains in late October. Eleven control goats came back on heat within 1½ weeks after the onset of short rains but resumption of cyclicity in the infected goats (two from each of the Arusha, Morogoro and Imbo groups) was delayed to an average of 3 weeks after the onset of rains. However, two of the control and one infected (Arusha) goats became pregnant thereafter.

Plasma progesterone levels

No marked regional (group) differences were noted in the monthly and peak luteal progesterone values prior to trypanosome infection (Fig. 1, Table 4). However, a significant difference in the mean monthly and luteal progesterone values were recorded within a month and 2nd cycle postinfection, respectively, between the infected and control goats and between infected groups. Whereas, the control goats maintained high mean progesterone levels (10.7 ± 0.8 nmol/l) in July, one month postinfection (Fig. 1), the mean monthly progesterone values of the infected groups (3.0 ± 0.5 nmol/l; 4.7 ± 0.6 nmol/l; 7.1 ± 0.8 nmol/l and 2.2 ± 0.3 nmol/l in the Lambwe, Arusha, Morogoro and Imbo groups, respectively) recorded during the same month, were significantly lower ($p < 0.001$).

The decline was, however, least in the Morogoro goats ($p < 0.001$) when compared to other infected groups. This pattern was maintained up to the end of August when all infected and control goats became acyclic. The mean progesterone levels of the infected groups were still slightly low in September (nonsignificant) and in October ($p < 0.01$) during the acyclic period. Luteal progesterone values (Table 4) followed the same trend between the infected and control goats from the 2nd cycle postinfection. The differences in hormonal levels among the infected groups were, however, equivocal.

Seasonal ovarian activity in the control goats were marked by a decline in the mean monthly progesterone values in August ($p < 0.01$) and September ($p < 0.001$). Following the onset of short rains, the hormonal levels started to rise. However, the mean monthly progesterone values of the pregnant infected Arusha goat was significantly lower (6.9 nmol/l, $p < 0.001$) than the pregnant control goat values (> 20 nmol/l).

Plasma oestradiol-17 β levels

The preinfection monthly oestradiol levels and corresponding preovulatory peaks were not significantly different between the experimental and control groups in May (Fig. 1, Table 5).

A significant difference in the monthly ($p < 0.01$) and peak ($p < 0.01$) values




Fig. 1. Mean monthly progesterone and oestradiol-17β in the infected and control goat groups.

Table 4. Mean luteal plasma progesterone levels (nmol/l) of infected and control goats

Approximate period in months	Cycles before (-) and after (+) infection	Lambwe Valley	Arusha	Morogoro	Imbo	Control	Level of significance F ₁ ratio	Level of significance F ₂ ratio
May to	-2	21.3±0.7 (10)	20.2±0.8 (9)	20.8±0.7 (10)	20.4±0.9 (10)	21.8±0.8 (13)	0.46 ns	0.82 ns
Mid-June	-1	21.8±0.8 (10)	21.7 ^b ±1.1 (9)	19.0 ^a ±0.7 (10)	19.6 ^{ab} ±0.8 (9)	19.3 ^a ±0.6 (13)	2.82**	2.79**
Mid-June ↓	1	19.6 ^a ±1.1 (8)	16.7 ^a ±1.5 (9)	18.3 ^a ±1.2 (9)	16.2 ^a ±2.0 (8)	19.8 ^a ±0.8 (13)	1.08 ns	1.56 ns
	2	5.4 ^a ±0.5 (8)	10.0 ^b ±1.2 (7)	13.8 ^c ±1.3 (9)	10.1 ^{bc} ±1.5 (4)	18.3 ^d ±0.9 (12)	9.89***	22.69***
	3	4.6 ^a ±0.6 (6)	8.7 ^b ±1.5 (5)	10.2 ^b ±0.9 (7)	-	18.8 ^c ±1.0 (10)	5.93***	27.46***
	4	-	-	5.6±1.3 (3)	-	19.3±0.9 (6)	-	-
	5	-	-	-	-	12.5±4.2 (3)	-	-
Mid-October	6	-	-	-	-	-	-	-
Mid-October	7	-	-	7.6±2.7 (2)	8.8±2.1 (2)	16.3±1.2 (9)	-	-
to November	8	-	6.7±2.4 (2)	9.7 (1)	-	16.9±0.8 (8)	-	-

- Values given as means ± SEM.

- F₁ and F₂ are F values between infected groups, infected and control groups, respectively.

- Means bearing the same superscripts (a b c d) in the same row do not differ significantly at (p < 0.05); control values in the 2nd and 3rd cycle are significant at (p < 0.001).

- ns, ** *** denote nonsignificance, significance at 5% and 1% level of confidence, respectively.

Table 5. Mean peak plasma oestradiol-17 β levels (pmol/l) of infected and control goats

Approximate period in months	Cycles before (-) and after (+) infection	Lambwe Valley	Arusha	Morogoro	Imbo	Control	Level of significance F ₁ ratio	Level of significance F ₂ ratio
May to	-2	896.7 \pm 29.8 (10)	899.6 \pm 51.0 (9)	808.9 \pm 51.2 (10)	785.2 \pm 32.4 (10)	847.5 \pm 23.5 (13)	2.00 ns	1.76 ns
Mid-June	-1	814.1 ^{ab} \pm 24.3 (10)	741.2 ^b \pm 32.3 (7)	870.8 ^a \pm 23.6 (9)	737.3 ^b \pm 28.4 (9)	786.6 ^{ab} \pm 17.3 (13)	5.59***	4.72***
Mid-June ↓	1	611.3 ^a \pm 55.1 (8)	582.5 ^a \pm 62.2 (9)	605.9 ^a \pm 55.1 (9)	576.1 ^a \pm 72.5 (8)	792.6 ^b \pm 22.4 (13)	0.08 ns	3.41**
	2	365.0 ^{ba} \pm 52.8 (8)	401.0 ^b \pm 36.1 (7)	404.9 ^b \pm 34.9 (9)	262.7 ^a \pm 18.8 (4)	679.1 ^c \pm 52.5 (12)	2.68**	11.15***
	3	274.4 ^a \pm 34.3 (5)	323.6 ^a \pm 80.5 (5)	365.4 ^a \pm 42.1 (4)	-	588.5 ^b \pm 29.8 (9)	0.59 ns	11.16***
	4	-	-	297.9 (1)	-	616.8 \pm 39.0 (6)	-	-
	5	-	-	-	-	506.0 \pm 93.7 (2)	-	-
Mid-October	6	-	-	-	-	-	-	-
Mid-October	7	-	428.2 (1)	340.9 \pm 22.3 (2)	285.9 \pm 59.5 (2)	790.3 \pm 27.3 (9)	-	-
to November	8	-	-	412.6 (1)	-	634.4 \pm 30.7 (8)	-	-

- Values given as means \pm SEM.

- F₁ and F₂ denote F values between the infected groups, infected and control groups, respectively.

- ns, ** *** denote nonsignificance, significance at 5% and 1% level of confidence, respectively.

- Means bearing the same superscripts (a b c) in the same row do not differ significantly at (p<0.05).

- Numbers in brackets denote cycling goats whose preovulatory oestradiol peaks were determined.

in June were attributed to the missed peak values in the Arusha (two), Imbo (one) and Morogoro (one) goats.

A significant decline in the monthly ($p < 0.01$) and peak values ($p < 0.05$; $p < 0.01$) were recorded in all infected groups within a month, and first and second cycle post-infection, respectively. The monthly levels remained low ($p < 0.01$) during the acyclic period in the infected groups. The Morogoro goats, however, maintained slightly higher monthly values in July and August ($p < 0.05$) than the Lambwe and Imbo goats. Changes in the preovulatory peaks were not significantly different among the infected groups.

Seasonality in the ovarian function in the control goats were marked by a significant decline in both monthly and peak oestradiol levels ($p < 0.01$) from July to October before increasing in November following resumption of the ovarian cycles. The monthly and peak levels were still significantly lower ($p < 0.01$) in the few infected goats which came back on heat in November.

Discussion

Chronic trypanosome infection resulted in progressive cessation of oestrous cycles and marked depression of plasma progesterone and oestradiol levels. Changes were initially more marked in the peak hormonal levels, but first decline occurred also in the basal values as the infection period progressed. Trypanosome-induced depression of plasma oestradiol- 17β and testosterone have recently been reported in goats (Kivihya et al., 1985; Waindi et al., 1986). Disruption of ovarian activity and prolonged elevation of progesterone levels due to the persistence of corpora lutea have also been reported recently in the British white goats and Boran cows chronically infected with *T. congolense* (Luckins et al., 1986). A close relationship in the fall of preovulatory oestradiol and luteal progesterone values observed in the infected goats in this study suggested the lack of normal corpora lutea (CL's) formation resulting from inhibited preovulatory oestradiol rise. Histologically, O'hara et al. (1985) and Mutayoba et al. (1988) reported the degeneration of partially lutenized tertiary follicles, absence of CL and lymphocytic infiltration in the degenerating ovarian stroma in the female goats infected with *T. congolense*. It seems from the above studies that trypanosomiasis causes variable reproductive anomalies in female animals ranging from luteolysis failure to impaired luteogenesis and degeneration of the entire ovarian stroma.

The plasma oestradiol levels obtained in this study were, however, higher than values reported in other breeds of goats but were closely similar to those previously reported in the Small East African goats from Uganda (Katongole and Gombe, 1985). Values ranging between 120–900 pmol/l were reported during normal oestrous cycles of the goats in that study.

The rate and degree of ovarian depression in the infected goats seemed to depend on the susceptibility of individual goats and in resistant goats, they also

depended on duration of the infection. The resistant Morogoro goats were better in maintaining the ovarian cycles and higher hormonal values than other infected groups, at least in the first 2 months of the infection. It was, however, observed from this study, that, chronic parasitaemia even at low levels may exacerbate the underlying ovarian lesions and reduce the fertility of resistant animals even further as evidenced by the loss of ovarian cycles in the Morogoro goats and levelling of hormonal values between infected groups in subsequent months.

Resumption of ovarian activity was observed in some chronically infected goats during the spell of short rains.

Although, this could have been due to the seasonal effect as also evidenced in the control goats and reported in other studies in the Small East African goats (Katongole and Gombe, 1985), it was an indication that trypanotolerant animals were potentially capable of regaining their ovarian function during the course of the infection. Indeed, one infected goat became pregnant. Quantitatively all goats which survived a six month infection period irrespective of the group were viewed as having been more resistant (trypanotolerant) than those which succumbed. These were 3 (Imbo), 5 (Arusha), 6 (Lambwe) and 7 (Morogoro) goats. The surviving Morogoro goats were in good body condition and had scanty parasitaemias than other infected groups. Their residual fertility by the time the study was terminated was still markedly reduced as indicated by low hormonal values even in the goat which became pregnant. These results, however, help to explain the high level of herd infertility and pregnancy failures often associated with trypanosomiasis in the field (Parkne and Dhake, 1972; Kanyari et al., 1983).

It was also noted in this study that trypanosomiasis delayed the resumption of ovarian cycles in the infected goats following the onset of short rains. In the field conditions, this delay is more likely to be worsened by an increase in the tsetse population and rate of trypanosome infections during the rainy season. To what extent does it affect the onset of goat breeding season in tsetse-endemic areas during rainy seasons is poorly documented and needs to be investigated. It suffices to speculate from the observations obtained from this study that there might be a considerable interference in the breeding season even in resistant infected animals so that young ones are born late when pastures are deteriorating. In this regard, other conventional disease control methods (namely tsetse-control and chemotherapy) are quite essential especially during the rains so as to exploit the genetic superiority of resistant livestock in improving their reproductive efficiency.

Acknowledgments

We wish to thank Messrs. J. Osaso, D. Ndegwa, L. M'rewa, F. Kariba and H. Odongo for their expert technical assistance and Miss J. Mbena for typing this manuscript. The Research project was partly financed by the German Academic Exchange Service (DAAD/AFAA) in collaboration with Sokoine University of Agriculture, Tanzania.

- Anosa V. O., Isoun T. T.: Further observations on the testicular pathology in *Trypanosoma vivax* infection of sheep and goats. *Res. Vet. Sci.* 28, 151–160 (1980).
- Ikede B. O.: Genital lesions in experimental chronic *Trypanosoma brucei* infection in rams. *Res. Vet. Sci.* 26, 145–151 (1979).
- Ikede B. O., Losos G. J.: Pathology of the disease in sheep produced experimentally by *Trypanosoma brucei*. *Vet. Path.* 9, 278–289 (1972).
- Ikede B. O., Losos G. J.: Pathogenesis of *Trypanosoma brucei* infection in sheep. III. Hypophysial and other endocrine lesions. *J. comp. Path.* 85, 37–44 (1975).
- Isoun T. T., Akpokodje J. U., Anosa V. O.: Testicular changes in White Fulani Zebu (Bunaji) cattle experimentally infected with *Trypanosoma vivax*. A preliminary report. *J. Nig. Vet. Med. Ass.* 4, 107–108 (1975).
- Kaaya G. P., Oduor-Okelo D.: The effects of *Trypanosoma congolense* infection on the testis and epididymis of the goat. *Bull. Anim. Hlth Prod. Afr.* 28, 1–5 (1980).
- Kanyari P. W. N., Allonby E. W., Wilson A. J., Munyua W. K.: Some economic effects on trypanosomiasis in goats. *Trop. Anim. Hlth Prod.* 15, 153–160 (1983).
- Katongole C. B., Gombe S.: A hormonal study of indigenous goats of Uganda. Symposium of Small Ruminants and Camel Production. ILCA, Addis Ababa, Ethiopia 1985.
- Kivihya L., Makawiti D. W., Gombe S.: The measurement of oestradiol-17 β by chemiluminescent immunoassay in goats with trypanosomiasis. In: Recent advances in the management and control of infections in Eastern Africa. Proc. 6th Ann. Med. Sci. Conf., Nairobi, Kenya (KEMRI-KETRI), p. 119–124 (1985).
- Kramer C. Y.: Extension of multiple range tests to group means with unequal number of replication. *Biometrics* 12, 307–310 (1956).
- Luckins A. G., Llewelyn C., Munro C. D., Murray M.: Effects of pathogenic trypanosomes on the mammalian reproductive system. IAEA-SM-292/34, P. 351–367 (1986).
- Murray M., Clifford D. J., Gettinby G., Snow F., McIntyre W. I. M.: Susceptibility to African trypanosomiasis of N'dama and Zebu cattle in areas of *Glossina morsitans submorsitans* challenge. *Vet. Rec.* 109, 503–510 (1981).
- Mutayoba B. M., Gombe S., Waindi E. N., Kaaya G. P.: Comparative trypanotolerance of the Small East African breed of goats from different localities to *Trypanosoma congolense* infection. *Vet. Parasit.* (1987, in press).
- Mutayoba B. M., Gombe S., Kaaya G. P., Waindi E. N.: Effect of chronic experimental *Trypanosoma congolense* infection on the ovaries, pituitary, thyroid and adrenal glands in female goats. *Res. Vet. Sci.* 44, 140–146 (1988).
- O'hara H. B., Gombe S., King G. P.: Responsiveness of pituitary to LRH and adrenals to ACTH following chronic *Trypanosoma congolense* infection in the female goat. VIIth International Congress of Protozoology, Nairobi, Kenya, June 22nd–29th, 1985.
- Paris J., Murray M., McOdimba F.: A comparative evaluation of the parasitological techniques currently available for the diagnosis of African trypanosomiasis in cattle. *Acta trop. (Basel)* 39, 307–316 (1982).
- Parkne D. J., Dhake R. P.: Abortions due to *Trypanosoma evansi* in the buffalo (*Bubalus bubalis*). *Ind. Vet. J.* 49, 1091–1092 (1972).
- Steel R. G. D., Torrie J. H.: Principles and procedures of statistics with special references to the biological sciences. McGraw-Hill Book Co. Inc., New York 1960.
- Waindi E. N., Gombe S., Oduor-Okelo D.: Plasma testosterone in *Trypanosoma congolense*-infected Toggenburg goats. *Arch. Endr.* 17, 9–17 (1986).

