

Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	45 (1988)
Heft:	3
Artikel:	Inhibition of "Plasmodium vinckeii"-malaria in mice by recombinant murine interferon- : short communication
Autor:	Bienzle, U. / Fritsch, K.-G. / Hoth, G.
DOI:	https://doi.org/10.5169/seals-314086

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Landesinstitut für Tropenmedizin, Berlin, FRG

Inhibition of *Plasmodium vinckeii*-malaria in mice by recombinant murine interferon- γ

Short communication

U. BIENZLE, K.-G. FRITSCH, G. HOTH, E. ROZDZINSKI, K. KÖHLER,
M. KALINOWSKI, P. KREMSNER, F. ROSENKAIMER, H. FELDMEIER

The activation of effector cells is an important aspect of immunity to malaria. Macrophages and natural killer cells are stimulated by interferon- γ (IFN- γ) to generate and release reactive superoxide (O_2^-) and its oxygen intermediates (Allison and Eugui, 1983). Rodent malaria infection was found to increase interferon production (Clark and Cowden, 1985) and oxidative killing of parasitized red blood cells (Ockenhouse and Shear, 1983, 1984; Ockenhouse et al., 1984). The release of tumor necrosis factor (TNF) by activated macrophages during malaria infection has also been described, but its role remains obscure (Grau et al., 1987). In this communication we report the effect of recently developed recombinant murine IFN- γ and TNF in mice infected with *Plasmodium vinckeii*. Ten-week-old inbred male BALB/c-mice were infected with the rodent malaria parasite *P. vinckeii* (kindly provided by Dr. Büngener, Tropeninstitut Hamburg). Blood was drawn from the tail of parasitized mice into 20 μ l heparinized capillary tubes, diluted with phosphate buffered saline (PBS) to final concentration of 10^3 – 10^5 parasitized erythrocytes and then injected intraperitoneally (i.p.). Control animals received the same volume of PBS.

Recombinant murine interferon- γ (r-IFN- γ), derived from *E. coli* (Genentech, USA), was supplied and tested for activity by Boehringer Ingelheim, FRG (Gray and Coeddel, 1983). The specific activity was $1-2 \times 10^7$ U/mg protein. r-IFN- γ was stored at 4°C and dissolved immediately before application in PBS containing 2% mouse albumin. 100 μ l were injected i.p. opposite to the injection site of the malaria parasites. Control animals received the same volume and concentration of PBS buffered mouse albumin. Recombinant murine TNF (supplied by Boehringer Ingelheim, FRG) was injected i.p. in concentrations of 0.01, 0.1, 1 and 10 μ g/100 μ l PBS containing 2% mouse albumin.

Survival time, haematological and parasitological parameters were recorded in 3 series of experiments to 1. standardize the *P. vinckeii*-inoculum, the r-IFN- γ concentration and the TNF concentration, 2. determine the effect of r-IFN- γ application before and after parasite inoculation, and 3. determine the effect of the combined r-IFN- γ and TNF application. Test groups were always matched with control groups of untreated mice. The analysis was performed by non-parametric statistics (Wilcoxon-test). Haematological and parasitological findings will be published elsewhere.

After establishing a suitable course of *P. vinckeii*-infection and the optimal concentration of r-IFN- γ , the survival times of 4 groups of mice were compared: one untreated control group, two other groups which received r-IFN- γ either before or after infection, and a fourth group in which prophylactic and therapeutic applications of r-IFN- γ were combined (Table 1). The results demonstrated that r-IFN- γ given after infection only slightly, but significantly, prolonged the median survival time. In mice receiving r-IFN- γ before malaria infection, the median survival time was significantly longer. If the prophylactic and therapeutic regimens were combined, the effect on the survival period was even more pronounced (Table 1).

In another set of experiments a therapeutic effect of TNF in infected mice could not be demonstrated, but TNF treated mice showed an increase in the mononuclear blood cell count compared to untreated controls. Moreover there was no difference in the course of the malaria

Correspondence: Prof. Dr. U. Bienzle, Landesinstitut für Tropenmedizin, Königin-Elisabeth-Strasse 32, D-1000 Berlin 19, FRG

Table 1. Three of four groups of mice infected with 10^4 parasitized erythrocytes received r-IFN- γ in a) a prophylactic regimen: 3×10^4 U/100 μ l PBS, 2% mouse albumin at 26 h, 16 h and 6 h before infection; b) a therapeutic regimen: 10 4 U/100 μ l PBS, 2% mouse albumin daily for 7 days after infection; c) a combination of a and b

r-IFN- γ (U/100 μ l)	Number of mice (n)	Median survival time (days)	Wilcoxon-test (p)
Control group	5	10.25	—
a) 3×10^4 prophylactic	10	11.00	<0.01
b) 1×10^4 7 days post infection	10	10.75	<0.05 >0.01
c) 3×10^4 prophylactic and 7 days post infection	10	12.25	<0.01

infection between groups receiving only a prophylactic regimen of r-IFN- γ and receiving a prophylactic regimen of r-IFN- γ plus a therapeutic regimen of TNF.

In rodent malaria, lymphokine activation of macrophages leads to enhanced phagocytosis of parasitized erythrocytes and to the production of intra- and extracellular oxygen intermediates (Dockrell and Playfair, 1983). In addition, IFN- γ stimulates the secretion of a factor, possibly TNF, which was thought, to inhibit parasite multiplication (Haidaris and Haynes, 1983).

Based on these reports, *P. vinckei*-infected BALB/c-mice were treated with r-IFN- γ and TNF. The rapidly fatal course of murine *P. vinckei*-malaria, which precludes interference of disease development by B-cell mediated immunity was regarded as an advantage of this animal model, allowing the study of lymphokine activity during malaria infection.

r-IFN- γ was presented in three different ways: a prophylactic and a therapeutic regimen, and a combination of both. Increasing doses of the prophylactic regimen effectively slowed the progress of the disease. The therapeutic regimen was less efficient, r-IFN- γ being effective too late to appreciably inhibit parasite multiplication. The combination of prophylactic and therapeutic applications acted in a cumulative way. The macrophage system, already activated before infection, was subjected to prolonged stimulation by daily r-IFN- γ doses. However, therapeutic applications of TNF did not further increase the effect of r-IFN- γ . Our results showed, that TNF did not influence the course of murine malaria infection in any way.

The animals died from malaria in all the experiments regardless of the time and amount of r-IFN- γ given. However, the findings raise the possibility that r-IFN- γ may play a role as a supportive therapeutic agent in the treatment of animal and human malaria, especially in the case of life threatening *P. falciparum*-infection, or against parasite strains which are partially or totally resistant against chemotherapeutic drugs.

Acknowledgment. The study was supported by Boehringer Ingelheim, FRG. Data of this communication are part of doctoral theses by G. H., K. K., M. K.

Allison A. C., Eugui E. M.: The role of cell mediated immune responses in resistance to malaria, with special reference to oxidant stress. *Ann. Rev. Immunol.* 1, 361–392 (1985).

Clark I. A., Cowden W. B., in: "Oxidative stress", p. 131. Academic Press, London 1985.

Dockrell H. M., Playfair J. H. L.: Killing of blood stage murine malaria parasites by hydrogen peroxide. *Infect. Immun.* 35, 456–459 (1983).

Grau G. E., Fajardo L. F., Piquet P. F., Allet B., Lambert B. A., Vasalli P.: Tumor necrosis factor (Cachectin) as an essential mediator in murine cerebral malaria. *Science* 237, 1210–1212 (1987).

Gray W. P., Coeddel D. V.: Cloning and expression of murine IFN- γ c DNA. *Proc. nat. Acad. Sci. USA* 80, 5842–5846 (1983).

Haidaris C. G., Haynes J.: Serum containing tumor necrosis factor is cytotoxic for the human malaria parasite *Plasmodium falciparum*. *Infect. Immun.* 42, 385–393 (1983).

Ockenhouse C. F., Shear H. L.: Malaria induced lymphokines stimulation of macrophages for enhanced phagocytosis. *Infect. Immun.* 42, 733–737 (1983).

Ockenhouse C. F., Shear H. L.: Oxidative killing of the intraerythrocytic malaria parasite *Plasmodium yoelii* by activated macrophages. *J. Immunol.* 132, 424–431 (1984).

Ockenhouse C. F., Schulman S., Shear H. L.: Induction of crisis forms in the human malaria parasite *Plasmodium falciparum* by γ -Interferon activated, monocyte-derived macrophages. *J. Immunol.* 133, 1601–1608 (1984).