

Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	45 (1988)
Heft:	4
Artikel:	Quantitation of antibodies to infective larvae in "Wuchereria bancrofti" filariasis : short communication
Autor:	Das, M.K. / Dash, A.P. / Ravindran, B.
DOI:	https://doi.org/10.5169/seals-314097

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Regional Medical Research Centre (Indian Council of Medical Research), P.O. Sainik School, Bhubaneswar 751005, India

Quantitation of antibodies to infective larvae in *Wuchereria bancrofti* filariasis

Short communication

M. K. DAS, A. P. DASH, B. RAVINDRAN, N. M. PATTNAIK, V. R. SUBRAMANYAM

A host of immunoassays have been devised in filariasis with the objective that the immunological laboratory parameters could be correlated with the disease status and eventually will provide sensitive and specific assay system for identifying infection. Antibody assays against microfilarial (mf) stages of *W. bancrofti*, namely somatic or excretory-secretory (ES) antigens and against related parasite antigens have been performed by a range of methods (Taylor and Denham, 1986). However, there are few studies directed at L₃ stages of *W. bancrofti*. In the present communication, an attempt has been made to evaluate L₃ specific antibody levels by ELISA in an infected population.

Materials and Methods

L₃ were dissected from adult female *Culex quinquefasciatus* caught in Puri district (*W. bancrofti* endemic region) of Orissa. A soluble extract of somatic antigens was prepared by homogenizing, sonicating and finally centrifuging the larval extract. The solution was then adsorbed with insolubilized rabbit antiserum against *C. quinquefasciatus* in order to remove any contaminating *Culex* materials since these are known to be immunogenic in man (Das and Dash, 1986). Human filarial sera (Das et al., 1987) were analyzed (at 300 fold dilution) by ELISA for antibody responses (IgG + M + A) against L₃ antigens. Serum from healthy individuals living in hilly regions, not endemic for filariasis, of Orissa served as non-endemic control.

Results and Discussion

Anti-L₃ antibody level in filarial sera was shown in Table 1. Taking mean A492 ± 3 SD of non-endemic normals as the reference point, the increased antibody level in each category of filariasis may be evaluated. Thus about 14% of endemic normals, 50% of asymptomatic carriers, 65% of chronic filarial and

Correspondence: Dr. M. K. Das, Regional Medical Research Centre (Indian Council of Medical Research), P.O. Sainik School, Bhubaneswar 751005, India

Table 1. L₃-specific antibody level in filarial sera as measured by ELISA

Group	Mean A492 ± SD (range)	No. positive/ No. tested	Positive %
Non-endemic normals	0.10 ± 0.06 (0.02 – 0.25)	0/22	0%
Endemic normals	0.21 ± 0.13 (0.04 – 0.60)	4/29	13.8%
Asymptomatic carriers	0.32 ± 0.16 (0.08 – 0.64)	12/25	48.0%
Chronic	0.38 ± 0.19 (0.10 – 0.80)	19/30	63.3%
TPE	0.56 ± 0.27 (0.20 – 1.40)	43/48	89.6%

90% of TPE patients could be designated as positive by L₃ antibody assay. Although anti-L₃ response was found to be minimal in endemic normals, some sera in this group nevertheless have high level of L₃ antibodies.

Acknowledgment

We thank Indian Council of Medical Research for providing necessary facilities.

- Das M. K., Dash A. P.: Detection of antibody to *Culex quinquefasciatus* in man. IRCS med. Sci. 14, 1190–1191 (1986).
 Das M. K., Subramanyam V. R., Ravindran B., Pattnaik N. M.: A study of the antigen, antibody and immune complex levels in *Wuchereria bancrofti* filariasis with reference to clinical status. J. trop. Med. Hyg. 90, 135–141 (1987).
 Taylor A. E. R., Denham D. A.: Immunology and diagnosis of filariasis. Trop. Dis. Bull. 83, R1–R20 (1986).