

**Zeitschrift:** Archives des sciences [1948-1980]  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 12 (1959)  
**Heft:** 8: Colloque Ampère : Maxwell-Ampère conference

**Artikel:** Dielectric saturation and molecular structure  
**Autor:** Piekara, A. / Kielich, S. / Chelkowski, A.  
**DOI:** <https://doi.org/10.5169/seals-739095>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 05.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Dielectric Saturation and Molecular Structure

by A. PIEKARA, S. KIELICH and A. CHELKOWSKI

Institute of Physics, Polish Academy of Sciences;  
A. Mickiewicz University, Poznan, Poland

---

## Résumé.

La saturation diélectrique positive consiste en l'augmentation de la constante diélectrique d'un liquide sous l'action d'un champ électrique. Cet effet a été trouvé dans le nitrobenzène et reconnu comme effet intermoléculaire. D'autre part, l'effet de saturation positive a été trouvé dans certains liquides dont les molécules présentent la rotation interne de groupes ayant chacun un moment électrique. Dans ce dernier cas, l'effet est intramoléculaire. La théorie de cet effet présentée par les auteurs admet l'existence des structures isomériques *droite* et *gauche*. Cette théorie explique l'existence de l'effet positif de la saturation et conduit à un terme additionnel dans la polarisation diélectrique, terme qui rend compte d'un nouveau mécanisme de cette polarisation consistant dans le fait que le champ électrique extérieur provoque des transitions *droite*  $\rightarrow$  *gauche*, de sorte que l'équilibre entre ces deux formes se déplace quelque peu vers la structure *gauche* ayant un moment électrique plus grand que celui de la structure *droite*.

The inverse or positive saturation effect in dielectric liquids consists in an increase  $\Delta\epsilon$  in the dielectric permittivity under the influence of an external electric field  $E$  [1, 2, 3, 4]. The effect is dependent on the quantity  $\frac{\partial}{\partial E} \langle m_E \rangle_E$ , where  $m_E$  denotes the projection of the electric moment of the molecule on the direction of the field vector  $E$ . With the local field  $F$ , the mean value of  $m_E$  is given by the formula

$$\langle m_E \rangle_E = \frac{\mu^2 F}{3kT} R_p - \frac{\mu^4 F^3}{45k^3 T^3} R_s, \quad (1)$$

provided the polarizability is due only to the orientational effect of the rigid dipoles  $\mu$  [5]. If this is the case, the quantities  $R_p$  and  $R_s$  are *correlation factors* for the polarisation and the saturation, respectively. For polar gases, these factors become unity, leading to the normal or negative saturation effect. However, in dense media such as condensed gases and liquids,  $R_p$  and  $R_s$  are strongly dependent on the intermolecular directional forces. In some cases, as in liquid nitrobenzene,  $R_s$  becomes negative

giving rise to the positive saturation effect. The graphs in Fig. 1 show the quantities  $R_p$  and  $R_s$  plotted *versus* the interaction energy in dimensionless units  $Y = \frac{W}{kT}$  of a system of two neighbouring dipoles tending to anti-parallel configuration. It is seen from the curve labelled  $R_s$  that, for the




Fig. 1.

Correlation factors  $R_p$ , for dielectric polarisation, and  $R_s$ , for dielectric saturation as functions of the interaction energy,  $Y = \frac{W}{kT}$ , expressed in dimensionless units. The case of pair coupling tending to an antiparallel configuration (e.g. nitrobenzene).

weak interaction existing in sufficiently dilute solutions of nitrobenzene in non-polar solvents, the normal saturation effect appears. Thus, the positive saturation effect was first recognized as due to molecular interaction. This point of view has received support from cryoscopic measurements [4, 6].

Now, some liquids present a positive saturation effect of a different kind. These are liquids the molecules of which exhibit internal rotation [2, 3, 7, 8, 9]. In certain liquids belonging to this group, an external field

produces a rise in the dielectric permittivity,  $\Delta\epsilon > 0$ ; this occurs both in the pure liquid and in solutions thereof in non-polar solvents at arbitrary concentrations. The mechanism of this kind of effect is clearly of an intramolecular nature. It is, nevertheless, similar to the mechanism of the intermolecular effect appearing in nitrobenzene. Here, however, the two interacting electric dipole moments are those of two rotating groups within the molecule. In such molecules the *trans* configuration is the privileged one. The external field acts to orient both moments in its direction and thus tends to diminish the angle between them. This results in a minute increase in the effective moment of the molecule and a corresponding increase in the dielectric permittivity. If the potential barrier is sufficiently high, this effect prevails over the normal saturation effect (which consists in a fall in  $\epsilon$ ) and the positive saturation effect appears.

If we follow the assumption made by Mizushima [10] for computing the dielectric polarisation, which is that of a single cosine potential barrier hindering the internal rotation, we again obtain formula (1). In the present case,  $\mu$  denotes the effective dipole moment of a molecule with two identical rotating groups, provided rotation is entirely free.  $R_p$  and  $R_s$  are factors reducing the polarisation term and saturation term, respectively, as accounting for hindered rotation. Both *reducing factors* are functions of the interaction energy given in dimensionless units,  $Z = \frac{V_0}{kT}$ . The amount  $2V_0$  is the difference in energy between the *cis* and *trans* positions. Thus, Mizushima's expression for  $R_p$  is obtained as follows:

$$R_p = 1 - \frac{I_1(Z)}{I_0(Z)} \quad (2)$$

whereas the reducing factor for dielectric saturation is given by the formula:

$$R_s = \frac{1}{4} - 2 \frac{I_1(Z)}{I_0(Z)} + \frac{5}{2} \left\{ \frac{I_1(Z)}{I_0(Z)} \right\}^2 - \frac{3}{4} \frac{I_2(Z)}{I_0(Z)}, \quad (3)$$

wherein  $I_0(Z)$ ,  $I_1(Z)$  and  $I_2(Z)$  are Bessel functions. It is easily seen by a survey of Fig. 2 that  $R_s$  can change its sign for  $Z \geq 0.30$ , thus explaining the inverse saturation effect consisting in an increase of the dielectric permittivity in all cases when the potential barrier is sufficiently high.

This, indeed, is the case of the 1,2-dihalogenoethanes experimentally investigated [2, 4, 9]. The measured value of the increase  $\Delta\epsilon$  in dielectric permittivity produced by an external field  $E$  allows to compute  $R_s$ . The theory using the model with single potential barrier yields qualitative

agreement with the experimental results. The degree of quantitative agreement is, however, unsatisfactory.

For this reason we resolved to test a theory assuming the existence of two rotational isomeres, to see whether such a theory would lead to better agreement with the measurements.




Fig. 2.

Reducing factors  $R_p$  and  $R_s$ , for dielectric polarisation and saturation, respectively, in 1, 2-dihalogenoethanes in gaseous form, assuming a single potential barrier (of height  $2Z$  expressed in  $kT$ ).

In considering liquid di-halogenated ethanes, let us assume, in agreement with various authors [10, 11, 12], that there is thermal equilibrium between three forms of various stability, the most stable of which is a trans structure, the remaining ones being two equivalent *gauche* structures of lesser stability. Since the dielectric saturation effect in these liquids is a positive one, it seems reasonable to suggest that the electric field can

stimulate transitions from the *trans* to the *gauche* forms, causing an increase in the effective electric moment. Following the procedure of Boltzmann statistics, a general expression for  $\langle m_e \rangle_E$  is obtained. For the case that




Fig. 3.

Plot of factors  $R_p^g$  and  $R_s^g$  expressing the contribution of the *gauche* form to the dielectric polarisation and saturation, respectively, vs. the energy difference between the *gauche* and *trans* structures.

the electric moment of the *trans* structure  $\mu_t$  is zero, the following formula holds:

$$\langle m_e \rangle_E = \frac{\mu_g^2 F}{3kT} R_p^g - \frac{\mu_g^4 F^3}{45k^3 T^3} R_s^g ; \quad (4)$$

here,  $\mu_g$  denotes the electric moment of the *gauche* structure.  $R_p^g$  and  $R_s^g$  are factors accounting for dielectric polarisation and dielectric saturation, respectively, and are given by

$$R_p^g = \frac{1}{1 + r_t}, \quad (5)$$

and

$$R_s^g = \frac{1 - \frac{3}{2} r_t}{(1 + r_t)^2}; \quad (6)$$

$r_t = \frac{N_t(0)}{N_g(0)}$  is the ratio of the numbers of molecules of *trans* and *gauche* structure at zero field, and is given by

$$r_t = \frac{1}{\omega} e^{\frac{\Delta U}{kT}}, \quad (7)$$

wherein  $\omega = 1,9$  [10], with regard to the presence of two *gauche* forms, and  $\Delta U = U_g - U_t$  is the difference between the energy values corresponding to the *gauche* and *trans* forms of the molecule.

The factors  $R_p^g$  and  $R_s^g$  are plotted *versus*  $\frac{\Delta U}{kT}$  in Fig. 3. Throughout the range of small differences in energy  $\Delta U$ , the factor  $R_s^g$  is positive, and we have the normal saturation effect,  $\Delta\varepsilon < 0$ . However, if  $\Delta U$  exceeds 0,24  $kT$ , the factor  $R_s^g$  becomes negative, yielding the inverse, that is positive effect  $\Delta\varepsilon > 0$ . This is the case of 1,2-dichloro- and 1,2-dibromoethane. For both these liquids, we obtained reasonable values for  $\mu_g$  and  $\Delta U$  (for the liquid state), as is shown in the following table:

Compound	Vapour			Liquid	
	$\Delta U$ from other investiga- tions [10] ( $k$ cal/mole)	$\mu_g$ computed according to the present theory from measured dipole moments [10] (D)	$\mu_g'$ from bond moments [12, 13] (D)	$\Delta U$ computed according to the present theory from measure- ments of $\Delta\varepsilon$ [2, 4, 9] and using $\mu_g$ ( $k$ cal/mole)	$\Delta U$ from other investiga- tions [10] ( $k$ cal/mole)
ClH <sup>2</sup> C — CH <sup>2</sup> Cl .	1,2	2,45	2,4	0,16	0
BrH <sup>2</sup> C — CH <sup>2</sup> Br .	1,4	ca. 2,3	2,3	ca. 0,4	$\geq 0,65$ , very disper- sed

The values obtained for  $\mu_g$  fit well with those computed for  $\mu'_g$  from the bond moments C — Cl or C — Br and the geometry of the molecule. As to the energy difference  $\hat{E}U$  for the liquid state, we consider the value 0 obtained from other experiment (see [10]) is too low, for the reason that it would yield a negative saturation effect which is contrary to experiment.

I think there is just one more point that is worth stressing. If an external electric field is present, then, according to our assumption, in addition to collision-stimulated transitions leading to thermal equilibrium between the *trans* and *gauche* forms, there occur field-stimulated *trans* → *gauche* transitions which shift the equilibrium somewhat towards the *gauche* forms possessing an increased electric moment. In this way, rotational isomerism leads to a *new mechanism of (collective) polarizability*, i.e. "field stimulated transition polarizability" due to the appearing of structures possessing a greater electric dipole moment than the vanishing structures they replace. That is why the dielectric saturation effect becomes positive.

#### RÉFÉRENCES

1. PIEKARA, A. and B. PIEKARA, *C. R. Ac. Sc.*, **203**, 552, 1058, 1936.
2. —— and A. CHEŁKOWSKI, *Journ. of Chem. Phys.*, **25**, 794, 1956.
3. CHEŁKOWSKI, A., *Journ. of Chem. Phys.*, **28**, 1249, 1958.
4. PIEKARA, A., A. CHEŁKOWSKI and S. KIELICH, *Z. Phys. Chemie*, **206**, 375, 1957.
5. —— *Proc. Roy. Soc., A* **172**, 360, 1939; *Acta Phys. Polonica*, **10**, 37, 107, 1950.
6. —— *Acta Phys. Polonica*, **11**, 99, 1951.
7. —— *Nuovo Cimento*, Supplemento al Vol. 9, Sez. X, 192, 1958.
8. —— *Acta Physica Polonica*, **18**, 1959, in print.
9. CHEŁKOWSKI, A., *Acta Physica Polonica*, in preparation.
10. MIZUSHIMA, San-Ichiro, *Structure of Molecules and Internal Rotation*, Acad. Press Inc., New York, 1954.
11. CABANNES, J., *Reports on Progress in Physics*, **X**, 378, 1946.
12. SMYTH, C. P., *Dielectric Behaviour and Structure*, Mc Graw-Hill Book Comp., New York, 1955.
13. STUART, H. A., *Die Structur des freien Moleküls*, Springer-Verlag, Berlin, 1952.