

Zeitschrift: Archives des sciences [1948-1980]
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 13 (1960)
Heft: 9: Colloque Ampère

Artikel: Nuclear magnetic resonance spectra of crystals rotated macroscopically
: fine structure of the spectrum of phosphorus pentachloride
Autor: Andrew, E.R. / Bradbury, A. / Eades, R.G.
DOI: <https://doi.org/10.5169/seals-738600>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Nuclear Magnetic Resonance Spectra of Crystals Rotated Macroscopically: Fine Structure of the Spectrum of Phosphorus Pentachloride

by E. R. ANDREW, A. BRADBURY, R. G. EADES and G. J. JENKS

Department of Physics, University College of North Wales, Bangor,
Caernarvonshire, Great Britain

Summary.

It is pointed out that the dipolar broadening of the nuclear magnetic resonance spectrum of a solid may effectively be removed by rotating the sample at a high speed about an axis inclined with an angle $54^{\circ} 44'$ to the direction of the field. Using this technique the doublet fine structure of the ^{31}P spectrum of solid phosphorus pentachloride has been recorded. The two lines are ascribed to the different chemical shifts in the $(\text{PCl}_4)^+$ and $(\text{PCl}_6)^-$ ions of which the solid is composed. The two shifts relative to aqueous phosphoric acid are found to be $+ 281$ and $- 96$ p.p.m.

When a solid whose n.m.r. spectrum is determined predominantly by nuclear dipolar interactions is rotated at high speed its spectrum changes. (Andrew, Bradbury and Eades 1958 *a, b*, Lowe 1959). It now consists of a central line narrower than the original static line, flanked by satellite lines set at integral multiples of the rotation frequency on either side of the central line. When the rotation rate exceeds the static linewidth, the satellites are resolved from the narrowed central line. As the rotation rate is increased the satellites move out and their intensity is diminished in such a way as to preserve the invariance of the second moment of the whole spectrum. When the rotation rate becomes much greater than the static linewidth the satellites are unobservably weak so that the narrowed central line alone is recorded. A detailed discussion of the form of the spectrum is given in the paper by Clough which follows. The width of the narrowed central line is given in first approximation by the factor $\frac{1}{2} (3 \cos^2 \gamma - 1)$ times the width of the static line, where γ is the angle between the axis of rotation and the direction of the applied magnetic field. We have found that the width of the central line follows this angular dependence quite well (Andrew, Bradbury and Eades 1959). In particular when $\gamma = \cos^{-1} 1/\sqrt{3} = 54^{\circ} 44'$ the central linewidth is found, as predicted,

to be quite small. For this orientation of the axis the static dipolar interaction is zero, and at high speeds of rotation the dipolar breadth of the central line should be small as in a liquid.

Fine structure of the type encountered in the high-resolution spectra of liquids is not normally observed in the n.m.r. spectra of solids, since it is masked by the stronger dipolar interactions. However, high-speed rotation of solids at the angle $54^{\circ} 44'$ offers the possibility of removing the dipolar broadening and thus of revealing fine structure.

PCl₅ SPECTRUM




Figure Caption. ³¹P magnetic resonance spectrum of polycrystalline phosphorus pentachloride rotated at high speed. The resonance frequency of the reference compound (85% aqueous phosphoric acid) is also indicated; this frequency was determined from recordings taken immediately before and after.

An example of the application of this method is provided by phosphorus pentachloride. The ³¹P resonance spectrum of a polycrystalline sample of phosphorus pentachloride rotated in an airtight container at the angle $54^{\circ} 44'$ is shown in the figure. The sidebands are scarcely distinguishable from the noise background, and the central spectrum consists of two sharp lines of approximately equal intensity. It is known from an X-ray diffraction study by Clark, Powell and Wells (1942) that in the solid state phosphorus pentachloride consists of equal numbers of tetrahedral (PCl₄)⁺

and octahedral $(\text{PCl}_6)^-$ ions packed in a tetragonal lattice. It is therefore reasonable to identify the two spectral lines with phosphorus nuclei in the two types of ion, the separation between the lines being ascribed to their different chemical shifts. In order to be sure that the separation does in fact have its origin in different chemical shifts measurements are in hand at different field strengths.

The doublet separation corresponds to a difference in chemical shift of 377 ± 10 parts per million. The chemical shifts* of each line have also been measured relative to the usual phosphorus standard, namely aqueous phosphoric acid, and are found to be $+ 281$ and $- 96$ p.p.m. These shifts fall within the range $+ 448$ to $- 222$ p.p.m. recorded for various liquid specimens (see for example Pople, Schneider and Bernstein 1959), though the difference between the shifts for the two ions is remarkably large.

RÉFÉRENCES

- ANDREW, E. R., A. BRADBURY and R. G. EADES (1958a), *Arch. Sci.*, *11*, 223.
 — A. BRADBURY and R. G. EADES (1958b), *Nature*, *182*, 1659.
 — A. BRADBURY and R. G. EADES (1959), *Nature*, *183*, 1802.
 CLARK, D., H. M. POWELL and A. F. WELLS (1942), *J. Chem. Soc.*, *1942*, 642.
 LOWE, I. J. (1959), *Phys. Rev. Letters*, *2*, 285.
 POPLE, J. A., W. G. SCHNEIDER and H. J. BERNSTEIN (1959), *High Resolution Nuclear Magnetic Resonance*, McGraw-Hill, p. 347.

Résumé.

SPECTRES DE RÉSONANCE MAGNÉTIQUE NUCLÉAIRE DE CRISTAUX
 TOURNÉS MACROSCOPIQUEMENT;
 STRUCTURE FINE DU SPECTRE DU PENTACHLORURE DE PHOSPHORE

On a montré que l'élargissement dipolaire du spectre de résonance magnétique d'un solide peut être supprimé de façon effective par rotation de l'échantillon à grande vitesse autour d'un axe incliné d'un angle de $54^\circ 44'$ sur la direction du champ. En employant cette technique la structure fine du doublet de P^{31} du pentachlorure de phosphore solide a été enregistrée.

Les deux lignes sont dues aux shifts différents des ions $(\text{PCl}_4)^+$ et $(\text{PCl}_6)^-$ dont est composé le solide. Les deux shifts sont trouvés, vis-à-vis d'une solution aqueuse d'acide phosphorique à $+ 281$ et $- 96$ p.p.m.

* The spectra were measured using a fixed field and a variable radiofrequency. The shift is therefore defined as $(\nu_r - \nu_c)/\nu_r$, where ν_c and ν_r are the frequencies of the compound and the reference respectively. This is equivalent to the usual definition $(H_c - H_r)/H_r$ when the frequency is held constant and the field is varied.