

**Zeitschrift:** Archives des sciences [1948-1980]  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 13 (1960)  
**Heft:** 9: Colloque Ampère

**Artikel:** Long range chemical shifts in aceto-acetic ester and acetylaceton  
**Autor:** Ranft, J.  
**DOI:** <https://doi.org/10.5169/seals-738631>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 29.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Long range chemical shifts in aceto-acetic ester and acetylaceton

by J. RANFT

Physikalisches Institut der Universität, Leipzig

---

Because of the small electron density near the protons in molecules one must consider the possibility, that large numbers of electrons in groups relatively far from the proton will produce significant shielding effects. McConnell [1] showed, that these long range shielding effects may be estimated in terms of the diamagnetic anisotropies of these distant electron groups. He gives the following formula for the part of the shielding tensor due to this group

$$\delta_d = \frac{1}{L_0} \left[ \frac{\chi}{R^3} - \frac{3\chi \cdot \mathfrak{R} \mathfrak{R}}{R^5} \right].$$


This gives for axial symmetry of the group

$$\delta_d = \frac{\Delta\chi}{3L_0 R^3} (1 - 3 \cos^2 \theta_z).$$

$\theta_z$  is the angle between  $\mathfrak{R}$ , the vector from the induced dipole to the regarded proton, and the symmetry axis of the distant group.

Guy and Tillieu [3] and Baudet, Tillieu and Guy [4] gave the theoretical values for the molar diamagnetic susceptibilities of some bonds. Narasimhan and Rogers [2] have calculated the diamagnetic anisotropy of the C=O bond with the aid of the chemical shifts in formamide and dimethylformamide.

It is possible with the above formulas to calculate the relative chemical shifts of the CH<sub>3</sub> lines in substances with keto enol tautomerism. The two groups have the forms



The screening constant of the  $\text{CH}_3$  protons must be averaged over the possible spatial arrangements. The location of the induced magnetic dipole has been taken at the middle of the bonds for  $\text{C} = \text{C}$  and  $\text{C} - \text{C}$  and at the oxygen atom for  $\text{C} = \text{O}$  and  $\text{C} - \text{O}$  [2].

TABLE 1.

| <i>Diamagnetic susceptibilities . <math>10^6</math></i> |     |                                                |     |
|---------------------------------------------------------|-----|------------------------------------------------|-----|
| $\Delta\chi$ ( $\text{C} - \text{C}$ ) = 1,16           | [3] | $\Delta\chi$ ( $\text{C} - \text{O}$ ) = 0,97  | [4] |
| $\chi_{xx}$ ( $\text{C} = \text{O}$ ) = — 6.77          | [2] | $\chi_{xx}$ ( $\text{C} = \text{C}$ ) = — 6.90 | [3] |
| $\chi_{yy}$ ( $\text{C} = \text{O}$ ) = 2.64            | [2] | $\chi_{yy}$ ( $\text{C} = \text{C}$ ) = — 5.03 | [3] |
| $\chi_{zz}$ ( $\text{C} = \text{O}$ ) = 1.09            | [2] | $\chi_{zz}$ ( $\text{C} = \text{C}$ ) = — 4.13 | [3] |

In table 1 are given the values of the diamagnetic susceptibilities. The evaluation gives

$$\delta_d (\text{CH}_3 \text{ keto}) = -0.17 \cdot 10^{-6}, \quad \delta_\alpha (\text{CH}_3 \text{ enol}) = -0.06 \cdot 10^{-6},$$

$$\Delta\delta = 0.11 \cdot 10^{-6} = 2.75 \text{ c/s at } 25 \text{ Mc/s}.$$

The experimental values are

Aceto-acetic ester:  $\Delta\delta = 6.8 \text{ c/s}$   
 $\Delta\delta = 6.0 \text{ c/s at } \infty \text{ dilution in isotropic solv.}$   
 Acetylacetone:  $\Delta\delta = 4.3 \text{ c/s}$   
 $\Delta\delta = 3.5 \text{ c/s at } \infty \text{ dilution in isotropic solv.}$

In view of the crude susceptibility values and the approximative method the results seems to be good.

I am greatly indebted to Dr Weissenfels for preparation of the substances.

1. McCONNELL, H. M., *J. Chem. Phys.*, **27**, 226, 1957.
2. NARASIMHAN, P. T., M. T. ROGERS, *J. Phys. Chem.*, **63**, 1388, 1959.
3. GUY, J., J. TILLIEU, *J. Chem. Phys.*, **24**, 1117, 1956.
4. BAUDET, J., J. TILLIEU, J. GUY, *C. R.*, **244**, 2920, 1957.